

# Dissolution Effect of Gastric and Intestinal pH for a BCS class II drug, Pioglitazone: New *in vitro* Dissolution System to Predict *in vivo* Dissolution

Yasuhiro Tsume\*, Gordon L Amidon and Susumu Takeuchi

University of Michigan, Ann Arbor, MI 48109, USA

**Keywords:** Dissolution; *In vitro*; Gastric pH; Intestinal pH; ASD; GIS

## Introduction

The U.S. Food and Drug Administration (FDA) released guidelines based on the Biopharmaceutics Classification System (BCS), which classifies drugs into 4 groups for the pharmaceutical industry in 2000 [1]. The European Medicines Agency (EMA) and FDA have released guidances for the bioequivalence (BE) and the requirements of the BCS-based bioequivalence (BE) and the requirements of the BCS-based bioequivalence (BE) and the requirements of the BCS-based bioequivalence (BE) for immediate release (IR) drug products in BCS class I [1,2]. The expansion of bioequivalence (BE) to some BCS class II drugs has been evaluated; particularly weak acidic drugs such as ibuprofen, ketoprofen and naproxen, are potential candidates for bioequivalence [3]. These drugs, which are highly permeable, can dissolve quickly at intestinal pH (6.5-7.0) and, therefore, behave like BCS class I drugs in the intestinal tract, even though they exhibit low solubility at acidic pH [4-6]. However, Alvarez and coworkers have been reported that it would be risky to bioequivalence (BE) BCS class II acidic drugs using just *in vitro* dissolution tests [7]. On the other hand, BCS class II weak basic drugs such as ketoconazole, dipyridamole, and carvedilol, easily dissolve in gastric pH and then may occur the precipitation or reach the supersaturation entering the duodenum due to higher environmental pH [8,9]. Even the limited solubility in intestinal pH, almost complete oral absorption of BCS class II drugs like ketoconazole, carvedilol, and pioglitazone, which dissolve in acidic pH but less dissolve in intestinal pH, has been reported [10-13]. The *in vitro* dissolution study with the USP apparatus II (paddle), which is a golden standard for *in vitro* dissolution study, for those drugs would not allow the observation of precipitation and supersaturation of those test drugs. Taken together, those reports and concerns raise questions of current *in vitro* dissolution systems and of the discrepancy of *in vitro*-*in vivo* dissolution. The development of new biopredictive dissolution systems, which emulates *in vivo* condition, would be required to predict the dynamic *in vivo* dissolution effects. Thus, new dissolution systems and techniques such as artificial stomach-duodenal (ASD) model and two-phase testing dissolution systems have been developed and assessed for their feasibilities and reliabilities to predict *in vivo* phenomena [14-19]. Those *in vitro* dissolution systems predict *in vivo* dissolution more accurately for BCS class II drugs than USP dissolution systems, indicating that the current *in vitro* dissolution systems may not be good enough to predict *in vivo* dissolution.

In this article, the new dissolution apparatus called gastro intestinal simulator (GIS) is constructed with three compartments to predict *in vivo* drug dissolution and to monitor the change of drug concentration in each compartment. This system would help to understand the drug dissolution stimulating *in vivo* environment and the transit of drug dissolution along with GI tract. We examine the dissolution profile of a BCS class II drug, pioglitazone, using the GIS and the feasibility of dissolution result by the GIS to predict *in vivo* dissolution of pioglitazone.

## Experimental

### Materials

Pioglitazone tablets were obtained from Takeda Pharmaceutical (Osaka, Japan). High-performance liquid chromatography (HPLC) grade acetonitrile was obtained from Fisher Scientific (St. Louis, MO). Trifluoroacetic acid (TFA), and other reagents and solvents were purchased from Aldrich Chemical Co. (Milwaukee, WI). All chemicals were either analytical or HPLC grade.

### GIS apparatus

The *in vitro* GIS model consists of three dissolution chambers representing the stomach, duodenum, and early jejunum with pH monitoring system in each chamber (Figure 1). The stomach and




Figure 1: The diagram of the GIS.

\*Corresponding author: Tsume Y, College of Pharmacy, The University of Michigan, 428 Church Street, Ann Arbor, MI 48109-1065, Phone: 734-647-2345; Fax: 734-763-6423; E-mail: [ytsume@umich.edu](mailto:ytsume@umich.edu)

Received May 27, 2013; Accepted September 23, 2013; Published September 30, 2013

Citation: Tsume Y, Amidon GL, Takeuchi S (2013) Dissolution Effect of Gastric and Intestinal pH for a BCS class II drug, Pioglitazone: New *in vitro* Dissolution System to Predict *in vivo* Dissolution. J Bioequiv Availab 5: 224-227. doi:10.4172/jbb.1000162

Copyright: © 2013 Tsume Y, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

duodenum chambers have capacities of 50-300 mL and 50-125 mL, respectively, and the fluid transit time between those chambers can be adjusted at the range of 1 - 40mL/minute by peristaltic pumps based on the simulation design of the gastric emptying time and GI transit time. The jejunal chamber is only the reservoir from the duodenal chamber and will provide the important information of drug concentration to predict the oral drug absorption of test compound. Those settings can be modified depending on the physiological characteristics of the species.

### Preparation of dissolution media

The following dissolution media were used; 0.01N HCl solution without enzymes (pH 2) and USP simulated intestinal fluid (SIF) without enzymes (pH 6.5).

### Dissolution Study of pioglitazone with USP dissolution apparatus II


Dissolution characteristics of a pioglitazone tablet were examined using the USP apparatus II (paddle), Hanson Research (Chatsworth, CA) Model SR6, Serial # 0698-1166 was used for dissolution studies at a rotational speed of 50 rpm at  $37.0 \pm 0.5^\circ\text{C}$  in 300 mL of the dissolution media; 50 mM phosphate buffer (simulated intestinal fluid: SIF) (pH 6.5). The test tablet was placed in the buffer media to start its dissolution study. Samples were drawn manually at 5, 10, 15, 20, 25, 30, 45, 60, 75, 90, 120, and 180 min and the same sample volume was replaced with the equal amount of blank medium tempered at  $37^\circ\text{C}$ . All samples were immediately spun at  $2,000 \times g$  for 20 seconds and the supernatant was diluted with the equal volume of methanol for HPLC analysis.

### Dissolution study of pioglitazone with GIS

In this experiment, 0.01N HCl solution (pH 2), and 50 mM phosphate buffer (simulated intestinal fluid: SIF) (pH 6.5) were used in each chamber. The initial volumes and pHs of the stomach and duodenum chambers were 300 mL at pH 2, representing 50 mL of stomach fluid plus 250 mL of dose volume, and 50 mL at pH 6.5, respectively. The test tablet was placed in the stomach chamber to start the dissolution study with the GIS. Samples were taken at the specific time points from each chamber over 65 - 120 minutes. The fluid was transferred at the range of 1-14 mL/minute to determine the dissolved drug concentration in each chamber. Those chambers were incubated at  $37^\circ\text{C}$  and pH changes were monitored by pH probes (Beckman Coulter, Brea, CA). Paddles were provided to mix fluid in each chamber. Fluid volume in duodenum was maintained at the constant volume. All samples were immediately spun at  $2,000 \times g$  for 20 seconds and the supernatant was diluted with the equal volume of methanol. The drug concentration was determined by HPLC analysis.

### Solubility study of a pioglitazone tablet

An excess amount of each drug, a pioglitazone tablet, was placed in 200 mL of each one of the following solutions to determine the saturated solubility of this pioglitazone product: 3.125 mM phosphate buffer containing 15.4 mM NaCl, 6.25 mM phosphate buffer containing 15.4 mM NaCl, 12.5 mM phosphate buffer containing 15.4 mM NaCl, 25 mM phosphate buffer containing 15.4 mM NaCl, and 50 mM phosphate buffer containing 15.4 mM NaCl (pH 6.5). The flasks were incubated at  $37^\circ\text{C}$  and samples were collected every 45 minutes for 4 hrs. All samples were centrifuged at  $12,000 \times g$  for 3 minutes and the supernatant was diluted with the equal volume of methanol. The drug solubility was determined by HPLC analysis.




### HPLC Analysis

The concentration of pioglitazone was determined on a Waters HPLC system (Waters, Inc., Milford, MA). The HPLC system consisted of two Waters pumps (model 515), a Waters autosampler (WISP model 712), and a Waters UV detector (996 photodiode array detector) controlled by Waters Millennium<sup>®</sup> 32 software (version 3.0.1). Samples were resolved in a Cadenza CD-C18 column (3  $\mu\text{m}$ , 3x100 mm) equipped with a guard column. The mobile phase consisted of 0.1% TFA/water (Solvent A) and 0.1% TFA/acetonitrile (Solvent B) with the solvent B gradient changing from 25 - 45% at a rate of 6.7%/minute during a 10 minute run. Standard curve generated for pioglitazone was

utilized for quantitation of integrated area under peaks. The detection wavelength was 269 nm.

## Results

In GIS, the dissolution of pioglitazone tablets with the initial transit time of 10 mL/minute exhibited the highest pioglitazone concentration ( $16.8 \pm 2.2 \mu\text{g/mL}$ ) in the duodenal chamber (pH 6.5). However, there is no significant difference in all dissolution profiles with the initial transit time of 5 - 10 mL/minute (Figure 2). The dissolution for a pioglitazone tablet in the USP dissolution apparatus II exhibited only 0.6% release (the range of concentration;  $0.56 \pm 0.04 \mu\text{g/mL}$ ) over 3 hours in 50 mM phosphate buffer (pH 6.5) (Figure 3). A pioglitazone tablet was quickly disintegrated and dissolved in the stomach chamber or in the USP vessel within 30 minutes regardless of transit time and dissolution methodology. The solubility studies of a pioglitazone tablet showed the solubility range of concentration 1.8 - 4.3  $\mu\text{g/mL}$  in different buffer strength (Table 1). This solubility range of pioglitazone was applied to Figure 2 to indicate the possible pioglitazone concentration in 3.125 - 50 mM phosphate buffer (pH 6.5). The highest concentrations of pioglitazone in duodenal and jejunal chambers were  $16.8 \mu\text{g/mL}$  and  $13.5 \mu\text{g/mL}$ , which were 7.5-fold and 9.25-fold higher than the concentration ( $1.8 \mu\text{g/mL}$ ) obtained by the solubility tests of pioglitazone in 50 mM phosphate buffer (pH 6.5). On the other hand, the highest concentration of pioglitazone in the USP dissolution apparatus II was  $0.6 \mu\text{g/mL}$ , which was 3.0-fold lower than the concentration obtained by the same tests.



Phosphate Conc. (mM)	Tablet
	Pioglitazone Conc. ( $\mu\text{g/mL}$ )
3.125	4.25
6.25	3.82
12.5	4.27
25	3.11
50	1.78

The pioglitazone tablet was placed in 200 mL of various concentration of phosphate buffer (pH 6.5). The flasks were incubated at  $37^\circ\text{C}$  for 4 hours. The concentration of pioglitazone was determined by HPLC analysis. Values presented are the mean (n=3).

**Table 1:** The solubility study of pioglitazone in the different concentration of phosphate buffer (pH 6.5).

## Discussion

Weak base drugs like ketoconazole and carvedilol might exhibit bioequivalence (BE) using existing *in vitro* dissolution methods like USP dissolution apparatus II. However, the progression of *in vivo* dissolution will be far more complicated than the results of *in vitro* dissolution studies with USP dissolution apparatus due to the pH changes in the GI tract causing the precipitation or the supersaturation of those BCS class II drugs. Consequently, the *in vitro* dissolution results may not predict *in vivo* dissolution profile. This type of discrepancy between *in vitro* dissolution profiles and *in vivo* dissolution results and, hence, *in vivo* drug absorption, especially for BCS class II and IV drugs, raises the questions regarding the current *in vitro* dissolution tests for bioequivalence. The gastro intestinal simulator (GIS) is the *in vitro* dissolution apparatus to understand the *in vivo* dissolution phenomena of those drugs. The buffer solution in each compartment of the GIS will be transferred from one chamber to the next chamber with various transit rates.

The duodenal and small-intestinal transit times in the fasted state are approximately 0.5 hour and 3 - 4 hours, respectively, and the gastric residence time is approximately 15 - 60 minutes [20-24]. The measured pH of human intestine in the fasted state is reportedly the range of 5.5 to 7.5 in the duodenum and 6.2 to 6.7 in the proximal small intestine [25-27]. Therefore, the middle of pH (pH 6.5) was adopted for this set of *in vitro* dissolution studies. A BCS class II drug product such as pioglitazone (pKa 5.8 and 6.4) would exhibit much lower solubility in the duodenum and jejunum in pH 6.5 than one in gastric pH due to its chemical characters. Therefore, the drug which has similar physicochemical characteristics to pioglitazone may be completely dissolved in the stomach but may occur the precipitation to inhibit its absorption or reach the supersaturation in small intestine to enhance its absorption.

The average volumes of human intestinal fluid in a fasted state are reportedly 184 mL for duodenum and 63 mL for jejunum [27]. With those volumes at the duodenum and jejunum in a fasted state and the solubility results in 50 mM phosphate buffer, less than 2% of pioglitazone (30 mg tablet) would be dissolved at those small intestinal segments at pH 6.5. The dissolution results of pioglitazone tablets in 50 mM phosphate buffer (pH 6.5) with USP dissolution apparatus II displayed even less drug dissolution over 3 hours, implying the less absorption of pioglitazone at the small intestine. However, the dissolution results of pioglitazone with the GIS exhibited 3.1- to 9.2-fold higher concentration than the obtained saturated concentration in duodenal and jejunal chambers, suggesting the supersaturation of pioglitazone in the proximal small intestinal region. As a result, higher *in vivo* absorption of pioglitazone will be anticipated. It has been reported that the orally dosed pioglitazone is well absorbed and its oral bioavailability is  $>80\%$  [11,12]. Less than 2% of *in vitro* dissolution profile may not explain well-absorbed oral dose of pioglitazone. This would be the discrepancy between current *in vitro* dissolution profiles and *in vivo* results.

The differences of the dissolution rate and solubility in between acidic pH and pH 6.5 were significant for weak basic drugs and the solubility of pioglitazone would be largely lowered at pH 6.5, which is closer to the one of pKa of pioglitazone. The dissolution rate of pioglitazone would be sensitive to the pH changes in the human intestine. This supersaturation phenomenon will not be observed in current dissolution systems like a USP apparatus II. As a result, it is extremely hard to predict *in vivo* dissolution. However, the GIS has the

capability to observe the supersaturation and, perhaps, precipitation and, hence, to predict better *in vivo* dissolution.

Buffer species and buffer capacity of dissolution medium along with buffer pH clearly have a significant effect on the dissolution rate for test drugs like ketoconazole and pioglitazone. Thus, the selection of dissolution media is crucial. It is likely that the concentration of phosphate buffer (50 mM) is too high for BE studies and may not reflect *in vivo* dissolution media, which is mainly bicarbonate [27,28]. The *in vitro* dissolution rate is clearly dependent on the pH, buffer species and buffer capacity of the medium for drugs products. The USP test does not reflect the human intestinal *in vivo* environment and may not be suitable for *in vitro* dissolution study to predict *in vivo* dissolution and, hence, to evaluate bioequivalence. The GIS has the potential to be a standard biopredictive *in vitro* dissolution method for the better prediction of *in vivo* dissolution and, therefore, bioequivalence.

## References

1. U.S. FDA Department of Health and Human Services (2000) Food and Drug Administration Center for Evaluation and Research., in Guidances for industry: Waiver of *in vivo* bioavailability and bioequivalence studies for immediate-release solid oral dosage forms based on a Biopharmaceutics Classification System.
2. Verbeeck RK, Musuamba FT (2012) The revised EMA guideline for the investigation of bioequivalence for immediate release oral formulations with systemic action. J Pharm Pharm Sci 15: 376-388.
3. Tubic-Grozdanis M, Bolger MB, Langguth P (2008) Application of gastrointestinal simulation for extensions for biowaivers of highly permeable compounds. AAPS J 10: 213-226.
4. Avdeef A, Box KJ, Comer JE, Gilges M, Hadley M, et al. (1999) pH-metric log P 11. pKa determination of water-insoluble drugs in organic solvent-water mixtures. J Pharm Biomed Anal 20: 631-641.
5. Potthast H, Dressman JB, Junginger HE, Midha KK, Oeser H, et al. (2005) Biowaiver monographs for immediate release solid oral dosage forms: ibuprofen. J Pharm Sci 94: 2121-2131.
6. Yazdani M, Briggs K, Jankovsky C, Hawi A (2004) The "high solubility" definition of the current FDA Guidance on Biopharmaceutical Classification System may be too strict for acidic drugs. Pharm Res 21: 293-299.
7. Alvarez C, Nunez I, Torrado JJ, Gordon J, Potthast H, et al. (2011) Investigation of the possibility of believers for ibuprofen. J Pharm Sci 100: 2343-2349.
8. Hsieh YL, Ilevbare GA, Van Eerdenbrugh B, Box KJ, Sanchez-Felix MV, et al. (2012) pH-Induced precipitation behavior of weakly basic compounds: determination of extent and duration of supersaturation using potentiometric titration and correlation to solid state properties. Pharm Res. 29: 2738-2753.
9. Psachoulas D, Vertzoni M, Goumas K, Kalioras V, Beato S, et al. (2011) Precipitation in and supersaturation of contents of the upper small intestine after administration of two weak bases to fasted adults. Pharm Res 28: 3145-3158.
10. Carver PL, Berardi RR, Knapp MJ, Rider JM, Kauffman CA, et al. (1994) *In vivo* interaction of ketoconazole and sucralfate in healthy volunteers. Antimicrob Agents Chemother 38: 326-329.
11. Eckland DA, Danhof M (2000) Clinical pharmacokinetics of pioglitazone. Exp Clin Endocrinol Diabetes 108: S234-S242.
12. Kadam R, Bourne D, Kompella U, Aquilante C (2013) Effect of Cytochrome P450 2C8\*3 on the Population Pharmacokinetics of Pioglitazone in Healthy Caucasian Volunteers. Biol Pharm Bull 36: 245-251.
13. Carino SR, Sperry DC, Hawley M (2006) Relative bioavailability estimation of carbamazepine crystal forms using an artificial stomach-duodenum model. J Pharm Sci 95: 116-125.
14. Carino SR, Sperry DC, Hawley M (2010) Relative bioavailability of three different solid forms of PNU-141659 as determined with the artificial stomach-duodenum model. J Pharm Sci 99: 3923-3930.
15. Mudie DM, Shi Y, Ping H, Gao P, Amidon GL, et al. (2012) Mechanistic analysis of solute transport in an *in vitro* physiological two-phase dissolution apparatus. Biopharm Drug Dispos 33: 378-402.
16. Shi Y, Gao P, Gong Y, Ping H (2010) Application of a biphasic test for characterization of *in vitro* drug release of immediate release formulations of celecoxib and its relevance to *in vivo* absorption. Mol Pharm 7: 1458-1465.
17. Vatieer J, Malikova-Sekera E, Vitre MT, Mignon M (1992) An artificial stomach-duodenum model for the in-vitro evaluation of antacids. Aliment Pharmacol Ther 6: 447-458.
18. Vatieer JL, Gao Z, Fu-Cheng XM, Vitre MT, Levy DA, et al. (1992) Evidence for the interaction between antacid and gastric mucosa using an "artificial stomach" model including gastric mucosa. J Pharmacol Exp Ther 263: 1206-1211.
19. Kaus LC, Fell JT (1984) Effect of stress on the gastric emptying of capsules. J Clin Hosp Pharm 9: 249-251.
20. Kortejärvi H, Shawahna R, Koski A, Malkki J, Ojala K, et al. (2010) Very rapid dissolution is not needed to guarantee bioequivalence for biopharmaceutics classification system (BCS) I drugs. J Pharm Sci 99: 621-625.
21. Mudie DM, Amidon GL, Amidon GE (2010) Physiological parameters for oral delivery and *in vitro* testing. Mol Pharm 7: 1388-1405.
22. Oberle RL, Chen TS, Lloyd C, Barnett JL, Owyang C, et al. (1990) The influence of the interdigestive migrating myoelectric complex on the gastric emptying of liquids. Gastroenterology 99: 1275-1282.
23. Reddy SM, Sinha VR, Reddy DS (1999) Novel oral colon-specific drug delivery systems for pharmacotherapy of peptide and nonpeptide drugs. Drugs Today (Barc) 35: 537-580.
24. Ibekwe VC, Fadda HM, McConnell EL, Khela MK, Evans DF, et al. (2008) Interplay between intestinal pH, transit time and feed status on the *in vivo* performance of pH responsive ileo-colonic release systems. Pharm Res 25: 1828-1835.
25. Lee KJ, Vos R, Janssens J, Tack J (2004) Influence of duodenal acidification on the sensorimotor function of the proximal stomach in humans. Am J Physiol Gastrointest Liver Physiol 286: G278-284.
26. Perez de la Cruz Moreno M, Oth M, Deferme S, Lammert F, Tack J, et al. (2006) Characterization of fasted-state human intestinal fluids collected from duodenum and jejunum. J Pharm Pharmacol 58: 1079-1089.
27. Karr WG, Osler AW, Sample AB (1935) Intubation Studies of the Human Small Intestine. IV. Chemical Characteristics of the Intestinal Contents in the Fasting State and as Influenced by the Administration of Acids, of Alkalies and of Water. J Clin Invest. 14: 893-900.
28. Sheng JJ, McNamara DP, Amidon GL (2009) Toward an *in vivo* dissolution methodology: a comparison of phosphate and bicarbonate buffers. Mol Pharm 6: 29-39.

**Citation:** Tsume Y, Amidon GL, Takeuchi S (2013) Dissolution Effect of Gastric and Intestinal pH for a BCS class II drug, Pioglitazone: New *in vitro* Dissolution System to Predict *in vivo* Dissolution. J Bioequiv Availab 5: 224-227. doi:[10.4172/jbb.1000162](https://doi.org/10.4172/jbb.1000162)