

Free Executive Summary Unequal Treatment: Confronting Racial and Ethnic Disparities in Health Care

Brian D. Smedley, Adrienne Y. Stith, and Alan R. Nelson, Editors, Committee on Understanding and Eliminating Racial and Ethnic Disparities in Health Care

ISBN: 0-309-08265-X, 782 pages, 6 x 9, hardback (2003)

This free executive summary is provided by the National Academies as part of our mission to educate the world on issues of science, engineering, and health. If you are interested in reading the full book, please visit us online at http://www.nap.edu/catalog/10260.html. You may browse and search the full, authoritative version for free; you may also purchase a print or electronic version of the book. If you have questions or just want more information about the books published by the National Academies Press, please contact our customer service department toll-free at 888-624-8373.

Racial and ethnic disparities in health care are known to reflect access to care and other issues that arise from differing socioeconomic conditions. There is, however, increasing evidence that even after such differences are accounted for, race and ethnicity remain significant predictors of the quality of health care received. In Unequal Treatment, a panel of experts documents this evidence and explores how persons of color experience the health care environment. The book examines how disparities in treatment may arise in health care systems and looks at aspects of the clinical encounter that may contribute to such disparities. Patients' and providers' attitudes, expectations, and behavior are analyzed. How to intervene? Unequal Treatment offers recommendations for improvements in medical care financing, allocation of care, availability of language translation, community-based care, and other arenas. The committee highlights the potential of cross-cultural education to improve provider-patient communication and offers a detailed look at how to integrate cross-cultural learning within the health professions. The book concludes with recommendations for data collection and research initiatives. Unequal Treatment will be vitally important to health care policymakers, administrators, providers, educators, and students as well as advocates for people of color.

ABSTRACT

Racial and ethnic minorities tend to receive a lower quality of healthcare than non-minorities, even when access-related factors, such as patients' insurance status and income, are controlled. The sources of these disparities are complex, are rooted in historic and contemporary inequities, and involve many participants at several levels, including health systems, their administrative and bureaucratic processes, utilization managers, healthcare professionals, and patients. Consistent with the charge, the study committee focused part of its analysis on the clinical encounter itself, and found evidence that stereotyping, biases, and uncertainty on the part of healthcare providers can all contribute to unequal treatment. The conditions in which many clinical encounters take place-characterized by high time pressure, cognitive complexity, and pressures for cost-containment—may enhance the likelihood that these processes will result in care poorly matched to minority patients' needs. Minorities may experience a range of other barriers to accessing care, even when insured at the same level as whites, including barriers of language, geography, and cultural familiarity. Further, financial and institutional arrangements of health systems, as well as the legal, regulatory, and policy environment in which they operate, may have disparate and negative effects on minorities' ability to attain quality care.

A comprehensive, multi-level strategy is needed to eliminate these disparities. Broad sectors—including healthcare providers, their patients, payors, health plan purchasers, and society at large—should be made aware of the healthcare gap between racial and ethnic groups in the United

States. Health systems should base decisions about resource allocation on published clinical guidelines, insure that physician financial incentives do not disproportionately burden or restrict minority patients' access to care, and take other steps to improve access-including the provision of interpretation services, where community need exists. Economic incentives should be considered for practices that improve provider-patient communication and trust, and reward appropriate screening, preventive, and evidence-based clinical care. In addition, payment systems should avoid fragmentation of health plans along socioeconomic lines.

The healthcare workforce and its ability to deliver quality care for racial and ethnic minorities can be improved substantially by increasing the proportion of underrepresented U.S. racial and ethnic minorities among health professionals. In addition, both patients and providers can benefit from education. Patients can benefit from culturally appropriate education programs to improve their knowledge of how to access care and their ability to participate in clinical-decision making. The greater burden of education, however, lies with providers. Cross-cultural curricula should be integrated early into the training of future healthcare providers, and practical, case-based, rigorously evaluated training should persist through practitioner continuing education programs. Finally, collection, reporting, and monitoring of patient care data by health plans and federal and state payors should be encouraged as a means to assess progress in eliminating disparities, to evaluate intervention efforts, and to assess potential civil rights violations.

Looking gaunt but determined, 59-year-old Robert Tools was introduced on August 21, 2001, as a medical miracle-the first surviving recipient of a fully implantable artificial heart. At a news conference, Tools spoke with emotion about his second chance at life and the quality of his care. His physicians looked on with obvious affection, grateful and honored to have extended Tools' life. Mr. Tools has since lost his battle for life, but will be remembered as a hero for undergoing an experimental technology and paving the way for other patients to undergo the procedure. Moreover, the fact that Tools was African American and his doctors were white seemed, for most Americans, to symbolize the irrelevance of race in 2001. According to two recent polls, a significant majority of Americans believe that blacks like Tools receive the same quality of healthcare as whites (Lillie-Blanton et al., 2000; Morin, 2001).

Behind these perceptions, however, lies a sharply contrasting reality. A large body of published research reveals that racial and ethnic minorities experience a lower quality of health services, and are less likely to receive even routine medical procedures than are white Americans. Relative to whites, African Americans-and in some cases, Hispanics-are less likely to receive appropriate cardiac medication (e.g., Herholz et al., 1996) or to undergo coronary artery bypass surgery (e.g., Ayanian et al., 1993; Hannan et al., 1999; Johnson et al., 1993; Petersen et al., 2002), are less likely to receive peritoneal dialysis and kidney transplantation (e.g., Epstein et al., 2000; Barker-Cummings et al., 1995; Gaylin et al., 1993), and are likely to receive a lower quality of basic clinical services (Ayanian et al., 1999) such as intensive care (Williams et al., 1995), even when variations in such factors as insurance status, income, age, co-morbid conditions, and symptom expression are taken into account. Significantly, these differences are associated with greater mortality among African-American patients (Peterson et al., 1997; Bach et al., 1999).

Study Charge and Committee Assumptions

These disparities prompted Congress to request an Institute of Medicine (IOM) study to assess differences in the kinds and quality of healthcare received by U.S. racial and ethnic minorities and non-minorities. Specifically, Congress requested that the IOM:

- Assess the extent of racial and ethnic differences in healthcare that are not otherwise attributable to known factors such as access to care (e.g., ability to payor insurance coverage);
- Evaluate potential sources of racial and ethnic disparities in healthcare, including the role of bias, discrimination, and stereotyping at the individual (provider and patient), institutional, and health system levels; and,
 - Provide recommendations regarding interventions to eliminate healthcare disparities.

This Executive Summary presents only abbreviated versions of the study committee's findings and recommendations. For the full findings and recommendations, and a more extensive justification of

each, the reader is referred to the committee report. Below, findings and recommendations are preceded by text summarizing the evidence base from which they are drawn. For purposes of clarity, some findings and recommendations are presented in a different sequence than they appear in the full report; however, their numeric designation remains the same.

Defining Racial and Ethnic Healthcare Disparities

The study committee defines *disparities* in healthcare as racial or ethnic differences in the quality of healthcare that are not due to access-related factors or clinical needs, preferences¹, and appropriateness of intervention (Figure S-1). The committee's analysis is focused at two levels: 1) the operation of healthcare systems and the legal and regulatory climate in which health systems function; and 2) discrimination at the individual, patient-provider level. Discrimination, as the committee uses the term, refers to differences in care that result from biases, prejudices, stereotyping, and uncertainty in clinical communication and decision-making. It should be emphasized that these definitions are not legal definitions. Different sources of federal, state and international law define discrimination in varying ways, with some focusing on intent and others emphasizing disparate impact.




FIGURE S-1 Differences, disparities, and discrimination: Populations with equal access to healthcare. SOURCE:

Gomes and McGuire, 2001.

¹ The committee defines patient preferences as patients' choices regarding healthcare that are based on a full and accurate understanding of treatment options. As discussed in Chapter 3 of this report, patients' understanding of treatment options is often shaped by the quality and content of provider-patient communication, which in turn may be influenced by factors correlated with patients' and providers' race, ethnicity, and culture. Patient preferences that are not based on a full and accurate understanding of treatment options may therefore be a source of racial and ethnic disparities in care. The committee recognizes that patients' preferences and clinicians' presentation of clinical information and alternatives influence each other, but found separation of the two to be analytically useful.

Evidence of Healthcare Disparities

Evidence of racial and ethnic disparities in healthcare is, with few exceptions, remarkably consistent across a range of illnesses and healthcare services. These disparities are associated with socioeconomic differences and tend to diminish significantly, and in a few cases, disappear altogether when socioeconomic factors are controlled. The majority of studies, however, find that racial and ethnic disparities remain even after adjustment for socioeconomic differences and other healthcare access-related factors (for more extensive reviews of this literature, see Kressin and Petersen, 2001; Geiger, this volume; and Mayberry, Mili, and Ofili, 2000).

Studies of racial and ethnic differences in cardiovascular care provide some of the most convincing evidence of healthcare disparities. The most rigorous studies in this area assess both potential underuse and overuse of services and appropriateness of care by controlling for disease severity using well-established clinical and diagnostic criteria (e.g., Schneider et al., 2001; Ayanian et al., 1993; Allison et al., 1996; Weitzman et al., 1997) or matched patient controls (Giles et al., 1995). Several studies, for example, have assessed differences in treatment regimen following coronary angiography, a key diagnostic procedure. These studies have demonstrated that differences in treatment are not due to clinical factors such as racial differences in the severity of coronary disease or overuse of services by whites (e.g., Schneider et al., 2001; Laouri et al., 1997; Canto et al., 2000; Peterson et al., 1997). Further, racial disparities in receipt of coronary revascularization procedures are associated with higher mortality among African Americans (Peterson et al., 1997).

Healthcare disparities are also found in other disease areas. Several studies demonstrate significant racial differences in the receipt of appropriate cancer diagnostic tests (e.g., McMahon et al., 1999), treatments (e.g., Imperato et al., 1996), and analgesics (e.g., Bernabei et al., 1998), while controlling for stage of cancer at diagnosis and other clinical factors. As is the case in studies of cardiovascular disease, evidence suggests that disparities in cancer care are associated with higher death rates among minorities (Bach et al., 1999). Similarly, African Americans with HIV infection are less likely than non-minorities to receive antiretroviral therapy (Moore et al., 1994), prophylaxis for pneumocystic pneumonia, and protease inhibitors (Shapiro et al., 1999). These disparities remain even after adjusting for age, gender, education, CD4 cell count, and insurance coverage (e.g., Shapiro et al., 1999). In addition, differences in the quality of HIV care are associated with poorer survival rates among minorities, even at equivalent levels of access to care (Bennett et al., 1995; Cunningham et al., 2000).

Racial and ethnic disparities are found in a range of other disease and health service categories, including diabetes care (e.g., Chin, Zhang, and Merrell, 1998), end-stage renal disease and kidney transplantation (e.g., Epstein et al., 2000; Kasiske, London, and Ellison, 1998; Barker-Cummings et al., 1995; Ayanian et al., 1999), pediatric care and maternal and child health, mental health, rehabilitative and nursing home services, and many surgical procedures. In some instances, minorities are *more* likely to receive certain procedures. As in the case of bilateral orchiectomy and amputation, however (which African Americans undergo at rates 2.4 and 3.6 times greater, respectively, than their white Medicare peers; Gornick et al., 1996), these are generally less desirable procedures.

Finding 1-1: Racial and ethnic disparities in healthcare exist and, because they are associated with worse outcomes in many cases, are unacceptable.

Recommendation 2-1: Increase awareness of racial and ethnic disparities in healthcare among the general public and key stakeholders.

Recommendation 2-2: Increase healthcare providers' awareness of disparities.

Racial Attitudes and Discrimination in the United States

By way of context, it is important to note that racial and ethnic disparities are found in many sectors of American life. African Americans, Hispanics, American Indians, and Pacific Islanders, and some Asian-American subgroups are disproportionately represented in the lower socioeconomic ranks, in lower quality schools, and in poorer-paying jobs. These disparities can be traced to many factors,

including historic patterns of legalized segregation and discrimination. Unfortunately, some discrimination remains. For example, audit studies of mortgage lending, housing, and employment practices using paired "testers" demonstrate persistent discrimination against African Americans and Hispanics. These studies illustrate that much of American social and economic life remains ordered by race and ethnicity, with minorities disadvantaged relative to whites. In addition, these findings suggest that minorities' experiences in the world outside of the healthcare practitioner's office are likely to affect their perceptions and responses in care settings.

Finding 2-1: Racial and ethnic disparities in healthcare occur in the context of broader historic and contemporary social and economic inequality, and evidence of persistent racial and ethnic discrimination in many sectors of American life.

Assessing Potential Sources of Disparities in Care

The studies cited above suggest that a range of patient-level, provider-level, and system-level factors may be involved in racial and ethnic healthcare disparities, beyond access-related factors.

Patient-Level Variables: The Role of Preferences, Treatment Refusal, and the Clinical Appropriateness of Care

Racial and ethnic disparities in care may emerge, at least in part, from a number of patient-level attributes. For example, minority patients are more likely to refuse recommended services (e.g., Sedlis et al., 1997), adhere poorly to treatment regimens, and delay seeking care (e.g., Mitchell and McCormack, 1997). These behaviors and attitudes can develop as a result of a poor cultural match between minority patients and their providers, mistrust, misunderstanding of provider instructions, poor prior interactions with healthcare systems, or simply from a lack of knowledge of how to best use healthcare services. However, racial and ethnic differences in patient preferences and care-seeking behaviors and attitudes are unlikely to be major sources of health care disparities. For example, while minority patients have been found to refuse recommended treatment more often than whites, differences in refusal rates are small and have not fully accounted for racial and ethnic disparities in receipt of treatments (Hannan et al., 1999; Ayanian et al., 1999). Overuse of some clinical services (i.e., use of services when not clinically indicated) may be more common among white than minority patients, and may contribute to racial and ethnic differences in discretionary procedures. Several recent studies, however, have assessed racial differences relative to established criteria (Hannan et al., 1999; Laouri et al., 1997; Canto et al., 2000; Peterson et al., 1997) or objective diagnostic information, and still find racial differences in receipt of care. Other studies find that overuse of cardiovascular services among whites does not explain racial differences in service use (Schneider et al., 2001).

Finally, some researchers have speculated that biologically based racial differences in clinical presentation or response to treatment may justify racial differences in the type and intensity of care provided. For example, racial and ethnic group differences are found in response to drug therapies such as enalapril, an angiotensin-converting-enzyme inhibitor used to reduce the risk of heart failure (Exner et al., 2001). These differences in response to drug therapy, however, are not due to "race" *per se* but can be traced to differences in the distribution of polymorphic traits between population groups (Wood, 2001), and are small in relation to the common benefits of most therapeutic interventions. Further, as noted above, the majority of studies document disparities in healthcare services and disease areas when interventions are equally effective across population groups-making the "racial differences" hypothesis an unlikely explanation for observed disparities in care.

Finding 4-2: A small number of studies suggest that racial and ethnic minority patients are more likely than white patients to refuse treatment. These studies find that differences in refusal rates are generally small and that minority patient refusal does not fully explain healthcare disparities.

Healthcare Systems-Level Factors

Aspects of health systems-such as the ways in which systems are organized and financed, and the availability of services-may exert different effects on patient care, particularly for racial and ethnic minorities. Language barriers, for example, pose a problem for many patients where health systems lack the resources, knowledge, or institutional priority to provide interpretation and translation services. Nearly 14 million Americans are not proficient in English, and as many as one in five Spanish-speaking Latinos reports not seeking medical care due to language barriers (The Robert Wood Johnson Foundation, 2001). Similarly, time pressures on physicians may hamper their ability to accurately assess presenting symptoms of minority patients, especially where cultural or linguistic barriers are present. Further, the geographic availability of healthcare institutions-while largely influenced by economic factors that are outside the charge of this study-may have a differential impact on racial and ethnic minorities, independent of insurance status (Kahn et al., 1994). A study of the availability of opioid supplies, for example, revealed that only one in four pharmacies located in predominantly non-white neighborhoods carried adequate supplies, compared with 72% of pharmacies in predominantly white neighborhoods (Morrison et al., 2000). Perhaps more significantly, changes in the financing and delivery of healthcare services-such as the shifts brought by cost-control efforts and the movement to managed care-may pose greater barriers to care for racial and ethnic minorities than for non-minorities (Rice, this volume). Increasing efforts by states to enroll Medicaid patients in managed care systems, for example, may disrupt traditional community-based care and displace providers who are familiar with the language, culture, and values of ethnic minority communities (Leigh, Lillie-Blanton, Martinez, and Collins, 1999). In addition, research indicates that minorities enrolled in publicly funded managed care plans are less likely to access services after mandatory enrollment in an HMO, compared with whites and other minorities enrolled in nonmanaged care plans (Tai-Seale et al., 2001).

Care Process-Level Variables: The Role of Bias, Stereotyping, Uncertainty

Three mechanisms might be operative in healthcare disparities from the provider's side of the exchange: bias (or prejudice) against minorities; greater clinical uncertainty when interacting with minority patients; and beliefs (or stereotypes) held by the provider about the behavior or health of minorities (Balsa and McGuire, 2001). Patients might also react to providers' behavior associated with these practices in a way that also contributes to disparities. Unfortunately, little research has been conducted to elucidate how patient race or ethnicity may influence physician decision-making and how these influences affect the quality of care provided. In the absence of such research, the study committee drew upon a mix of theory and relevant research to understand how clinical uncertainty, biases or stereotypes, and prejudice might operate in the clinical encounter.

Clinical Uncertainty

Any degree of uncertainty a physician may have relative to the condition of a patient can contribute to disparities in treatment. Doctors must depend on inferences about severity based on what they can see about the illness and on what else they observe about the patient (e.g., race). The doctor can therefore be viewed as operating with prior beliefs about the likelihood of patients' conditions, "priors" that will be different according to age, gender, socioeconomic status, and race or ethnicity. When these priors—which are taught as a cognitive heuristic to medical students—are considered alongside the information gained in a clinical encounter, both influence medical decisions.

Doctors must balance new information gained from the patient (sometimes with varying levels of accuracy) and their prior expectations about the patient to determine the diagnosis and course of treatment. If the physician has difficulty accurately understanding the symptoms or is less sure of the "signal"-the set of clues and indications that physicians rely upon to make diagnostic decisions-then he or she is likely to place greater weight on the "priors." The consequence is that treatment decisions and patients' needs are potentially less well matched.

The Implicit Nature of Stereotypes

A large body of research in psychology has explored how stereotypes evolve, persist, shape expectations, and affect interpersonal interactions. Stereotyping can be defined as the process by

which people use social categories (e.g., race, sex) in acquiring, processing, and recalling information about others. The beliefs (stereotypes) and general orientations (attitudes) that people bring to their interactions help to organize and simplify complex or uncertain situations and give perceivers greater confidence in their ability to understand a situation and respond in efficient and effective ways (Mackie, Hamilton, Susskind, and Rosselli, 1996).

Although functional, social stereotypes and attitudes also tend to be systematically biased. These biases may exist in overt, explicit forms, as represented by traditional bigotry. However, because their origins arise from virtually universal social categorization processes, they may also exist, often unconsciously, among people who strongly endorse egalitarian principles and truly believe that they are not prejudiced (Dovidio and Gaertner, 1998). In the United States, because of shared socialization influences, there is considerable empirical evidence that even well-meaning whites who are not overtly biased and who do not believe that they are prejudiced typically demonstrate unconscious implicit negative racial attitudes and stereotypes (Dovidio, Brigham, Johnson, and Gaertner, 1996). Both implicit and explicit stereotypes significantly shape interpersonal interactions, influencing how information is recalled and guiding expectations and inferences in systematic ways. They can also produce self-fulfilling prophecies in social interaction, in that the stereotypes of the perceiver influence the interaction with others in ways that conform to stereotypical expectations (Jussim, 1991).

Healthcare Provider Prejudice or Bias

Prejudice is defined in psychology as an unjustified negative attitude based on a person's group membership (Dovidio et al., 1996). Survey research suggests that among white Americans, prejudicial attitudes toward minorities remain more common than not, as over half to three-quarters believe that relative to whites, minorities-particularly African Americans-are less intelligent, more prone to violence, and prefer to live off of welfare (Bobo, 2001). It is reasonable to assume, however, that the vast majority of healthcare providers find prejudice morally abhorrent and at odds with their professional values. But healthcare providers, like other members of society, may not recognize manifestations of prejudice in their own behavior.

While there is no direct evidence that provider biases affect the quality of care for minority patients, research suggests that healthcare providers' diagnostic and treatment decisions, as well as their feelings about patients, are influenced by patients' race or ethnicity. Schulman et a1. (1999), for example, found that physicians referred white male, black male, and white female hypothetical "patients" (actually videotaped actors who displayed the same symptoms of cardiac disease) for cardiac catheterization at the same rates (approximately 90% for each group), but were significantly less likely to recommend catheterization procedures for black female patients exhibiting the same symptoms. Weisse et a1. (2001), using a similar methodology as that of Schulman, found that male physicians prescribed twice the level of analgesic medication for white "patients" than for black "patients." Female physicians, in contrast, prescribed higher doses of analgesics for black than for white "patients," suggesting that male and female physicians may respond differently to gender and/ or racial cues. In another experimental design, Abreu (1999) found that mental health professionals subliminally "primed" with African American stereotype-laden words were more likely to evaluate the same hypothetical patient (whose race was not identified) more negatively than when primed with neutral words. And in a study based on actual clinical encounters, van Ryn and Burke (2000) found that doctors rated black patients as less intelligent, less educated, more likely to abuse drugs and alcohol, more likely to fail to comply with medical advice, more likely to lack social support, and less likely to participate in cardiac rehabilitation than white patients, even after patients' income, education, and personality characteristics were taken into account. These findings suggest that while the relationship between race or ethnicity and treatment decisions is complex and may also be influenced by gender, providers' perceptions and attitudes toward patients are influenced by patient race or ethnicity, often in subtle ways.

Medical Decisions Under Time Pressure with Limited Information

Studies suggest that several characteristics of the clinical encounter increase the likelihood that stereotypes, prejudice, or uncertainly may influence the quality of care for minorities (van Ryn, 2002).

In the process of care, health professionals must come to judgments about patients' conditions and make decisions about treatment, often without complete and accurate information. In most cases, they must do so under severe time pressure and resource constraints. The assembly and use of these data are affected by many influences, including various "gestalts" or cognitive shortcuts. In fact, physicians are commonly trained to rely on clusters of information that functionally resemble the application of "prototypic" or stereotypic constellations. These conditions of time pressure, resource constraints, and the need to rely on gestalts map closely onto those factors identified by social psychologists as likely to produce negative outcomes due to lack of information, to stereotypes, and to biases (van Ryn, 2002).

Patient Response: Mistrust and Refusal

As noted above, the responses of racial and ethnic minority patients to healthcare providers are also a potential source of disparities. Little research has been conducted as to how patients may influence the clinical encounter. It is reasonable to speculate, however, that if patients convey mistrust, refuse treatment, or comply poorly with treatment, providers may become less engaged in the treatment process, and patients are less likely to be provided with more vigorous treatments and services. But these kinds of reactions from minority patients may be understandable as a response to negative racial experiences in other contexts, or to real or perceived mistreatment by providers. Survey research, for example, indicates that minority patients perceive higher levels of racial discrimination in healthcare than non-minorities (LaVeist, Nickerson, and Bowie, 2000; Lillie-Blanton et al., 2000). Patients' and providers' behavior and attitudes may therefore influence each other reciprocally, but reflect the attitudes, expectations, and perceptions that each has developed in a context where race and ethnicity are often more salient than these participants are even aware of. In addition, it is clear that the healthcare provider, rather than the patient, is the more powerful actor in clinical encounters. Providers' expectations, beliefs, attitudes, and behaviors are therefore likely to be a more important target for intervention efforts.

Finding 3-1: Many sources-including health systems, healthcare providers, patients, and utilization managers-may contribute to racial and ethnic disparities in healthcare.

Finding 4-1: Bias, stereotyping, prejudice, and clinical uncertainty on the part of healthcare providers may contribute to racial and ethnic disparities in healthcare. While indirect evidence from several lines of research supports this statement, a greater understanding of the prevalence and influence of these processes is needed and should be sought through research.

Interventions to Eliminate Racial and Ethnic Disparities in Healthcare

Legal, Regulatory, and Policy Interventions "De-Fragmentation" of Healthcare Financing and Delivery

Racial and ethnic minorities are more likely than whites to be enrolled in "lower-end" health plans, which are characterized by higher per capita resource constraints and stricter limits on covered services (Phillips et al., 2000). The disproportionate presence of racial and ethnic minorities in lower-end health plans is a potential source of health care disparities, given that efforts to control for insurance status in studies of healthcare disparities have not taken detailed account of variations among health plans. Such socioeconomic fragmentation of health plans engenders different clinical cultures, with different practice norms, tied to varying per capita resource constraints (Bloche, 2001).

Equalizing access to high-quality plans can limit such fragmentation.

Public healthcare payors such as Medicaid should strive to help beneficiaries access the same health products as privately-insured patients. This recommendation is also reflected in the IOM *Quality Chasm* report's strategies for focusing health systems on quality, in its call to "eliminate or modify payment practices that fragment the care system" (IOM, 2001, p.13).

Recommendation 5-1: Avoid fragmentation of health plans along socioeconomic lines.

Strengthening Doctor-Patient Relationships

Several lines of research suggest that the consistency and stability of the doctor-patient relationship is an important determinant of patient satisfaction and access to care. Having a usual source of care is associated, for example, with use of preventive care services (Agency for Healthcare Research and Quality, 2001). In addition, having a consistent relationship with a primary care provider may help to address minority patient mistrust of healthcare systems and providers, particularly if the relationship is with a provider who is able to bridge cultural and linguistic gaps (La Viest, Nickerson, and Bowie, 2000). Minority patients, however, are less likely to enjoy a consistent relationship with a provider, even when insured at the same levels as white patients (Lillie-Blanton, Martinez, and Salganicoff, 2001). This is due in part to the types of health systems in which they are enrolled and the relative lack of providers located in minority communities.

Health systems should attempt to ensure that every patient, whether insured privately or publicly, has a sustained relationship with an attending physician able to help the patient effectively navigate the healthcare bureaucracy. Federal and state performance standards for Medicaid managed care plans, for example, should include guidelines to ensure the stability of patients' assignments to primary care providers (and these providers' accessibility), reasonable patient loads per primary care physician, and time allotments for patient visits.

Recommendation 5-2: Strengthen the stability of patient-provider relationships in publicly funded health plans.

Patient and provider relationships will also be strengthened by greater racial and ethnic diversity in the health professions. Racial concordance of patient and provider is associated with greater patient participation in care processes, higher patient satisfaction, and greater adherence to treatment (Cooper-Patrick et al., 1999). In addition, racial and ethnic minority providers are more likely than their non-minority colleagues to serve in minority and medically underserved communities (Komaromy et al., 1996). The benefits of diversity in health professions fields are significant, and illustrate that a continued commitment to affirmative action is necessary for graduate health professions education programs, residency recruitment, and other professional opportunities.

Recommendation 5-3: Increase the proportion of underrepresented U.S. racial and ethnic minorities among health professionals.

Patient Protections

Much of the political focus on Capitol Hill in the summer of 2001 was devoted to managed care regulation. To one extent or another, the various bills debated would all extend protections to enrollees in private managed care organizations, providing avenues for appeal of care denial decisions, improving access to specialty care, requiring health plans to disclose information about coverage, banning physician "gag" clauses, and providing other legal remedies to resolve disputes. Publicly funded health plans, however, are not addressed in these legislative proposals. Given that many minorities are disproportionately represented among the publicly insured who receive care within managed care organizations, the same patient protections that apply to the privately insured should apply to those in publicly funded plans (Hashimoto, 2001).

Recommendation 5-4: Apply the same managed care protections to publicly funded HMO enrollees that apply to private HMO enrollees.

Civil Rights Enforcement

Enforcement of regulation and statute is also an important component of a comprehensive strategy to address healthcare disparities, but unfortunately has been too often relegated to low-priority status. The U.S. DHHS Office of Civil Rights (OCR) is charged with enforcing several relevant federal statutes and regulations that prohibit discrimination in healthcare (principally Title VI of the 1964 Civil Rights Act). The agency, however, has suffered from insufficient resources to investigate complaints of possible violations, and has long abandoned proactive, investigative strategies (Smith, 1999). Complaints to the agency declined in the early 1990s, but have increased in recent years, while funding has remained level in terms of appropriated dollars but lower in terms of spending power

after adjusting for inflation (U.S. Commission on Civil Rights, 2001). The agency should be equipped with sufficient resources to better address these complaints and carry out its oversight responsibilities.

Recommendation 5-5: Provide greater resources to the U.S. DHHS Office for Civil Rights to enforce civil rights laws.

Health Systems Interventions

A variety of interventions applied at the level of health systems may be effective as a part of a comprehensive, multi-level strategy to address racial and ethnic disparities in healthcare.

Evidence-Based Cost Control

In the current era of continually escalating healthcare costs, cost containment is an important goal of all health systems. To the extent possible, however, medical limit setting by health plans should be based on evidence of effectiveness. The application of evidence to healthcare delivery, such as through the use of evidence-based guidelines, can help to address the problem of potential underuse of services resulting from capitation or per case payment methods, as noted in the IOM *Quality Chasm* report (IOM, 2001). Evidence-based guidelines offer the advantages of consistency, predictability, and objectivity that general, discretionary advisory statements do not. In addition, because evidence-based guidelines and standards directly promote accountability, they also indirectly affect equity of care

In actual practice, however, a pragmatic balance must be sought between the advantages and limitations of guidelines, such as the tension between the goal of standardization versus the need for clinical flexibility. Disclosing health plans' clinical protocols offers one means of achieving this balance, as it would aid both private sector and public efforts in balancing the virtues of rules and discretion. To achieve this, private accrediting entities and state regulatory bodies could require that health plans publish their clinical practice protocols, along with supporting evidence, thereby opening these protocols to professional and consumer review (Bloche, 2001).

Recommendation 5-6: Promote the consistency and equity of care through the use of evidence-based guidelines.

Financial Incentives in Healthcare

Financial factors, such as capitation and health plan incentives to providers to practice frugally, can pose greater barriers to racial and ethnic minority patients than to white patients, even among patients insured at the same level. Low payment rates limit the supply of physician (and other healthcare provider) services to low-income groups, disproportionately affecting ethnic minorities (Rice, this volume). Inadequate supply takes the form of too few providers participating in plans serving the poor, and provider unwillingness to spend adequate time with patients. This time pressure may contribute to poor information exchange between physicians and members of minority groups.

If appropriately crafted, however, financial incentives to physicians can serve a positive role in efforts to reduce disparities in care. Economic rewards for time spent engaging patients and their families can help physicians to overcome barriers of culture, communication, and empathy. In addition, incentives that encourage physicians to adhere to evidence-based protocols for frugal practice and to engage in age- and gender-appropriate disease screening can promote efficient, quality care and penalize deviations, regardless of race or ethnicity. Further, financial incentives linked to favorable clinical outcomes, where reasonably measurable (e.g. control of diabetes, asthma, and high blood pressure) can also promote equity of care (Bloche, 2001). Again, this recommendation is consistent with the IOM *Quality Chasm* report, which calls for healthcare organizations, clinicians, purchasers, and other stakeholders to "align the incentives inherent in payment and accountability processes with the goal of quality improvement" (IOM, 2001, p.10).

Recommendation 5-7: Structure payment systems to ensure an adequate supply of services to minority patients and limit provider incentives that may promote disparities. Recommendation 5-8: Enhance patient-provider communication and trust by providing financial incentives for practices that reduce barriers and encourage evidence-based practice.

Interpretation Services

As noted above, many racial and ethnic minorities find that language barriers pose a significant problem in their efforts to access healthcare. Language barriers may affect the delivery of adequate care through poor exchange of information, loss of important cultural information, misunderstanding of physician instruction, poor shared decision making, or ethical compromises (e.g., difficulty obtaining informed consent; Woloshin et al., 1995). Linguistic difficulties may also result in decreased adherence with medication regimes, poor appointment attendance (Manson, 1988), and decreased satisfaction with services (Carrasquillo et al., 1999; David and Rhee, 1998; Derose and Baker, 2000).

Broader use of professional interpretation services has been hampered by a number of logistical and resource constraints. For example, in some regions of the country, few trained professional interpreters are available, and reimbursement for interpretation services via publicly funded insurance such as Medicaid is often inadequate. Greater resources are needed to support professional interpretation services, and more research and innovation should identify effective means to harness new technologies (e.g., simultaneous telephone interpretation) to aid interpretation.

Recommendation 5-9: Support the use of interpretation services where community need exists.

Community Health Workers

Community health workers-often termed lay health advisors, neighborhood workers, indigenous health workers, health aides, *consejera*, or *promotora-fulfill* multiple functions in helping to improve access to healthcare. Community health workers can serve as liaisons between patients and providers, educate providers about community needs and the culture of the community, provide patient education, contribute to continuity and coordination of care, assist in appointment attendance and adherence to medication regimens, and help to increase the use of preventive and primary care services (Brownstein et al., 1992; Earp and Flax, 1999; Jackson and Parks, 1997). In addition, some evidence suggests that lay health workers can help improve the quality of care and reduce costs (Witmer et al., 1995), and improve general wellness by facilitating community access to and negotiation for services (Rodney et al., 1998).

Recommendation 5-10: Support the use of community health workers.

Multidisciplinary Teams

Research demonstrates that multidisciplinary team approaches—including physicians, nurses, dietitians, and social workers, among others—can effectively optimize patient care. This effect is found in randomized controlled studies of patients with coronary heart disease, hypertension, and other diseases, and has extended to strategies for reducing risk behaviors and conditions such as smoking, sedentary lifestyle and obesity (Hill and Miller, 1996). Multidisciplinary teams coordinate and streamline care, enhance patient adherence through follow-up techniques, and address the multiple behavioral and social risks faced by patients. These teams may save costs and improve the efficiency of care by reducing the need for face-to-face physician visits and improve patients' day-to-day care between visits. Further, such strategies have proven effective in improving health outcomes of minorities previously viewed as "difficult to serve" (Hill and Miller, 1996). Multidisciplinary team approaches should be more widely instituted as strategy for improving care delivery, implementing secondary prevention strategies, and enhancing risk reduction.

Recommendation 5-11: Implement multidisciplinary treatment and preventive care teams.

Patient Education and Empowerment

Increasingly, researchers are recognizing the important role of patients as active participants in clinical encounters (Korsch, 1984). Patient education efforts have taken many forms, including the use of books and pamphlets, in-person instruction, CD-ROM-based educational materials, and internet-based information. These materials guide patients through typical office visits and provide information about asking appropriate questions and having their questions answered, communicating with the provider when instructions are not understood or cannot be followed, and being an active

participant in decision-making. While evaluation data are limited, particularly with respect to racial and ethnic minority patients, preliminary evidence suggests that patient education can improve patients' skills and knowledge of clinical encounters and improve their participation in care decisions.

Recommendation 5-12: Implement patient education programs to increase patients' knowledge of how to best access care and participate in treatment decisions.

Cross-Cultural Education in the Health Professions

Given the increasing racial and ethnic diversity of the U.s. population, the development and implementation of training programs for healthcare providers offers promise as a key intervention strategy in reducing healthcare disparities. As a result, cross-cultural education programs have been developed to enhance health professionals' awareness of how cultural and social factors influence healthcare, while providing methods to obtain, negotiate and manage this information clinically once it is obtained. Cross-cultural education can be divided into three conceptual approaches focusing on attitudes (cultural sensitivity/awareness approach), knowledge (multicultural/categorical approach), and skills (cross-cultural approach), and has been taught using a variety of interactive and experiential methodologies. Research to date demonstrates that training is effective in improving provider knowledge of cultural and behavioral aspects of healthcare and building effective communication strategies. Despite progress in the field, however, several challenges exist, including the need to define educational core competencies, reach consensus on approaches and methodologies, determine methods of integration into the medical and nursing curriculum, and develop and implement appropriate evaluation strategies. These challenges should be addressed to realize the potential of cross-cultural education strategies.

Summary of Findings

Finding 1-1: Racial and ethnic disparities in healthcare exist and, because they are associated with worse outcomes in many cases, are unacceptable.

Finding 2-1: Racial and ethnic disparities in healthcare occur in the context of broader historic and contemporary social and economic inequality, and evidence of persistent racial and ethnic discrimination in many sectors of American life.

Finding 3-1: Many sources—including health systems, healthcare providers, patients, and utilization managers—may contribute to racial and ethnic disparities in healthcare.

Finding 4-1: Bias, stereotyping, prejudice, and clinical uncertainty on the part of healthcare providers may contribute to racial and ethnic disparities in healthcare. While indirect evidence from several lines of research supports this statement, a greater understanding of the prevalence and influence of these processes is needed and should be sought through research.

Finding 4-2: A small number of studies suggest that racial and ethnic minority patients are more likely than white patients to refuse treatment. These studies find that differences in refusal rates are generally small and that minority patient refusal does not fully explain healthcare disparities.

Summary of Recommendations

General Recommendations

Recommendation 2-1: Increase awareness of racial and ethnic disparities in healthcare among the general public and key stakeholders.

Recommendation 2-2: Increase healthcare providers' awareness of disparities.

Legal, Regulatory, and Policy Interventions

Recommendation 5-1: Avoid fragmentation of health plans along socio-economic lines.

Recommendation 5-2: Strengthen the stability of patient-provider relationships in publicly funded health plans.

Recommendation 5-3: Increase the proportion of underrepresented U.S. racial and ethnic minorities among health professionals.

Recommendation 5-4: Apply the same managed care protections to publicly funded HMO enrollees that apply to private HMO enrollees.

Recommendation 5-5: Provide greater resources to the U.S. DHHS Office for Civil Rights to enforce civil rights laws.

Summary of Recommendations (continued)

Health Systems Interventions

Recommendation 5-6: Promote the consistency and equity of care through the use of evidence-based guidelines.

Recommendation 5-7: Structure payment systems to ensure an adequate supply of services to minority patients, and limit provider incentives that may promote disparities.

Recommendation 5-8: Enhance patient-provided communication and trust by providing financial incentives for practices that reduce barriers and encourage evidence-based practice.

Recommendation 5-9: Support the use of interpretation services where community need exists.

Recommendation 5-10: Support the use of community health workers.

Recommendation 5-11: Implement multidisciplinary treatment and preventive care teams.

Patient Education and Empowerment

Recommendation 5-12: Implement patient education programs to increase patients' knowledge of how to best access care and participate in treatment decisions.

Cross-Cultural Education in the Health Professions

Recommendation 6-1: Integrate cross-cultural education into the training of all current and future health professionals.

Data Collection and Monitoring

Recommendation 7-1: Collect and report data on health care access and utilization by patients' race, ethnicity, socioeconomic status, and where possible, primary language.

Recommendation 7-2: Include measures of racial and ethnic disparities in performance measurement.

Recommendation 7-3: Monitor progress toward the elimination of healthcare disparities.

Recommendation 7-4: Report racial and ethnic data by OMS categories, but use subpopulation groups where possible.

Research Needs

Recommendation 8-1: Conduct further research to identify sources of racial and ethnic disparities and assess promising intervention strategies.

Recommendation 8-2: Conduct research on ethical issues and other barriers to eliminating disparities.

Recommendation 6-1: Integrate cross-cultural education into the training of all current and future health professionals.

Data Collection and Monitoring

Standardized data collection is critically important in the effort to understand and eliminate racial and ethnic disparities in healthcare. Data on patient and provider race and ethnicity would allow researchers to better disentangle factors that are associated with healthcare disparities, help health plans to monitor performance, ensure accountability to enrolled members and payors, improve patient choice, allow for evaluation of intervention programs, and help identify discriminatory practices. Unfortunately, standardized data on racial and ethnic differences in care are generally unavailable. Federal and state-supported data collection efforts are scattered and unsystematic, and many health plans, with a few notable exceptions, do not collect data on enrollees' race, ethnicity, or primary language.

A number of ethical, logistical, and fiscal concerns present challenges to data collection and monitoring, including the need to protect patient privacy, the costs of data collection, and resistance from healthcare providers, institutions, plans and patients. In addition, health plans have raised significant concerns about how such data will be analyzed and reported. The challenges to data collection should be addressed, as the costs of failing to assess racial and ethnic disparities in care may outweigh new burdens imposed by data collection and analysis efforts.

Recommendation 7-1: Collect and report data on healthcare access and utilization by patients' race, ethnicity, socioeconomic status, and where possible, primary language.

Recommendation 7-2: Include measures of racial and ethnic disparities in performance measurement.

Recommendation 7-3: Monitor progress toward the elimination of healthcare disparities.

Recommendation 7-4: Report racial and ethnic data by federally defined categories, but use subpopulation groups where possible.

Recommendation 8-1: Conduct further research to identify sources of racial and ethnic disparities and assess promising intervention strategies.

Recommendation 8-2: Conduct research on ethical issues and other barriers to eliminating disparities.

Needed Research

While the literature that the committee reviewed provides significant evidence of racial and ethnic disparities in care, the evidence base from which to better understand and eliminate disparities in care remains less than clear. Several broad areas of research are needed to clarify how race and ethnicity are associated with disparities in the process, structure, and outcomes of care. Research must provide a better understanding of the contribution of patient, provider, and institutional characteristics on the quality of care for minorities. Research has been notably absent in other areas. More research is needed, for example, to understand the extent of disparities in care faced by Asian-American, Pacific-Islander, American Indian and Alaska Native, and Hispanic populations, and to better understand and surmount barriers to research on healthcare disparities, including those related to ethical issues in data collection.

References

Abreu JM. (1999). Conscious and nonconscious African American stereotypes: Impact on first impression and diagnostic ratings by therapists. *Journal of Consulting and Clinical Psychology* 67(3):387-93.

Agency for Healthcare Research and Quality. (2001). Addressing racial and ethnic disparities in healthcare. Fact sheet accessed from internet site www. ahrq.gov /research/ disparit.htm on December 18, 2001.

Allison JJ, Kiefe CI, Centor RM, Box JB, Farmer RM. (1996). Racial differences in the medical treatment of elderly Medicare patients with acute myocardial infarction. *Journal of General Internal Medicine* 11:736-43.

Ayanian JZ, Udvarhelyi IS, Gatsonis CA, Pasho, CL, Epstein AM. (1993). Racial differences in the use of revascularization procedures after coronary angiography. *Journal of the American Medical Association* 269:2642-6.

Ayanian JZ, Weissman JS, Chasan-Taber S, Epstein AM. (1999). Quality of care by race and gender for congestive heart failure and pneumonia. *Medical Care 37:1260-9*.

Bach PB, Cramer LD, Warren JL, Begg CB. (1999). Racial differences in the treatment of early-stage lung cancer. *New England Journal of Medicine 341:1198-205*.

Balsa A, McGuire TG. (2001). Prejudice, uncertainty and stereotypes as sources of health care disparities. Boston University, unpublished manuscript.

Barker-Cummings C, McClellan W, Soucie, JM, Krisher J. (1995). Ethnic differences in the use of peritoneal dialysis as initial treatment for end-stage renal disease. *Journal of the American Medical Association* 274(23):1858-1862.

Bennett CL, Horner RD, Weinstein RA, Dickinson GM, Dehovitz JA, Cohn SE, Kessler HA, Jacobson I, Goetz MB, Simberkoff M, Pitrak D, George WL, Gilman SC, Shapiro MF. (1995). Racial differences in care among hospitalized patients with pneumocyctis carinii pneumonia in Chicago, New York, Los Angeles, Miami, and Raleigh-Durham. *Archives of Internal Medicine* 155(15):1586-92.

Bernabei R, Gambassi G, Lapane K, et al. (1998). Management of pain in elderly patients with cancer. *Journal of the American Medical Association* 279:1877-82.

Bloche MG. (2001). Race and discretion in American medicine. *Yale Journal of Health Policy, Law, and Ethics* 1:95-131. Bobo LD. (2001). Racial attitudes and relations at the close of the twentieth century. In Smelser NJ, Wilson WI, and Mitchell F (Eds.), *America Becoming: Racial Trends and Their Consequences*. Washington, DC: National Academy Press.

Brogan D, Tuttle EP. (1988). Transplantation and the Medicare end-stage renal disease program [Letter]. *New England Journal of Medicine 319:55*.

Brownstein IN, Cheal N, Ackermann SP, Bassford TL, Campos-Outcalt D. (1992). Breast and cervical cancer screening in minority populations: A model for using lay health educators. *Journal of Cancer Education* 7(4):321-326.

Canto JG, AllisonJJ, Kiefe CI, Fincher C, Farmer R, Sekar P, Person S, Weissman NW. (2000).

Relation of race and sex to the use of reperfusion therapy in Medicare beneficiaries with acute myocardial infarction. *New England Journal of Medicine* 342:1094-1100.

Carrasquillo O, Orav EJ, Brennan TA, Burstin HR. (1999). Impact of language barriers on patient satisfaction in an emergency department. *Journal of General Internal Medicine* 14(2):82-7.

Chin MH, Zhang JX, Merrell K. (1998). Diabetes in the African-American Medicare population: Morbidity, quality of care, and resource utilization. *Diabetes Care* 21(7):1090-1095.

Cooper-Patrick L, Gallo JJ, Gonzales JJ, Vu HT, Powe NR, Nelson C, Ford DE. (1999). Race, gender, and partnership in the patient-physician relationship. *Journal of the American Medical Association* 282(6):583-9.

Cunningham WE, Mosen DM, Morales LS. (2000). Ethnic and racial differences in long-term survival from hospitalization for HIV infection. *Journal of Health Care for the Poor and Underserved* 11(2):163-178.

David RA, Rhee M. (1998). The impact of language as a barrier to effective health care in an underserved urban Hispanic community. *Mount Sinai Journal of Medicine* 65(5-6):393-397.

Derose KP, Baker DW. (2000). Limited English proficiency and Latinos' use of physician services. *Medical Care Research and Review* 57(1):76-91.

Dovidio JF, Brigham JC, Johnson BT, Gaertner SL. (1996). Stereotyping, prejudice, and discrimination: Another look. In Macrae N, Stangor C, and Hewstone M (Eds.), *Stereotypes and Stereotyping* 276-319. New York: Guilford.

Dovidio JF, Gaertner SL. (1998). On the nature of contemporary prejudice: The causes, consequences, and challenges of aversive racism. In Eberhardt J and Fiske ST (Eds.), *Confronting Racism: The Problem and the Response* 3-32. Thousand Oaks, CA: Sage.

Earp JL, Flax VL. (1999). What lay health advisors do: An evaluation of advisors' activities.

Cancer Practice 7(1):16-21.

Epstein AM, Ayanian JZ, Keogh JH, Noonan SJ, Armistead N, Cleary PD, Weissman JS, David-Kasdan JA, Carlson D, Fuller I, March D, Conti R. (2000). Racial disparities in access to renal transplantation. *New England Journal of Medicine* 343(21):1537-1544.

Exner DV, Dries DL, Domanski MJ, Cohn IN. (2001). Lesser response to angiotens in converting-enzyme inhibitor therapy in black as compared with white patients with left ventricular dysfunction. *New England Journal of Medicine* 344:1351-1357.

Gaylin DS, Held PI, Port FK, et al. (1993). The impact of comorbid and sociodemographic factors on access to renal transplantation. *Journal of the American Medical Association* 269:603-608.

Geiger J. (this volume). Racial and ethnic disparities in diagnosis and treatment: A review of the evidence and a consideration of causes.

Giles WH, Anda RF, Casper ML, Escobedo LG, Taylor HA. (1995). Race and sex differences in rates of invasive cardiac procedures in U.s. hospitals. *Archives of Internal Medicine* 155:318-24.

Gomes C, McGuire TG. (2001). Identifying the sources of racial and ethnic disparities in health care use. Unpublished manuscript.

Gornick ME, Egers PW, Reilly TW, Mentnech RM, Fitterman LK, Kucken LE, Via deck Be. (1996). Effects of race and income on mortality and use of services among Medicare beneficiaries. *New England Journal of Medicine* 335(11):791-799.

Hannan EL, Van Ryn M, Burke I, et al. (1999). Access to coronary artery bypass surgery by race/ethnicity and gender among patients who are appropriate for surgery. *Medical Care 37:68-77*.

Hashimoto DM. (2001). The proposed Patients' Bill of Rights: The case of the missing equal protection clause. *Yale Journal of Health Policy, Law, and Ethics 1:77-93*.

Herholz H, Goff DC, Ramsey DJ, Chan FA, Ortiz C, Labarthe DR, Nichaman MZ. (1996). Women and Mexican Americans receive fewer cardiovascular drugs following myocardial infarction than men and non-Hispanic whites: The Corpus Christi Heart Project, 1988-1990. Journal of Clinical Epidemiology 49(3):279-87.

Hill MN, Miller NH. (1996). Compliance enhancement. A call for multidisciplinary team approaches. *Circulation* 93(1):4-6

House JS, Williams D. (2000). Understanding and reducing socioeconomic and racial/ethnic disparities in health. In Smedley BD and Syme SL (Eds.), *Promoting Health: Intervention Strategies from Social and Behavioral Research*. Washington, DC: National Academy Press.

Imperato PI, Nenner RP, Will TO. (1996). Radical prostatectomy: Lower rates among African American men. *Journal of the National Medical Association 88*(9):589-94.

Institute of Medicine. (2001). Crossing the Quality Chasm. Washington, DC: The National Academy Press.

Jackson EJ, Parks CP. (1997). Recruitment and training issues from selected lay health advisor programs among African Americans: A 20-year perspective. *Health Education & Behavior* 24(4):418-431.

Johnson, PA, Lee TH, Cook EF, Rouan GW, Goldman 1. (1993). Effect of race on the presentation and management of patients with acute chest pain. *Annals of Internal Medicine* 118:593-601.

Jussim, L. (1991). Social perception and social reality: A reflection-construction model. *Psychological Review* 98:54-73. Kahn KL, Pearson ML, Harrison ER, Desmond KA, Rogers WH, Rubenstein LV, Brook RH, Keeler EB. (1994). Health care for black and poor hospitalized Medicare patients. *Journal of the American Medical Association* 271(15):1169-1174.

Kaplan GA, Everson SA, Lynch JW. (2000). The contribution of social and behavioral research to an understanding of the distribution of disease: A multilevel approach. In Smedley BD and Syme SL (Eds.), *Promoting Health: Intervention Strategies from Social and Behavioral Research*. Washington, DC: National Academy Press.

Kasiske B, London W, Ellison MD. (1998). Race and socioeconomic factors influencing early placement on the kidney transplant waiting list. *Journal of the American Society of Nephrology* 9(11):2142-2147.

Kjellstrand CM. (1988). Age, sex, and race inequality in renal transplantation. Archives of Internal Medicine 148(6):1305-

Komaromy M, Grumbach K, Drake M, Vranizan K, Lurie N, Keane D, Bindman AB. (1996).

The role of black and Hispanic physicians in providing health care for underserved populations. *New England Journal of Medicine 334:1305-1310*.

Korsch BM. (1984). What do patients and parents want to know? What do they need to know? *Pediatrics* 74(5 Pt 2):917-919.

Kressin NR, Petersen LA. (2001). Racial differences in the use of invasive cardiovascular procedures: Review of the literature and prescription for future research. *Annals of Internal Medicine* 135(5):352-366.

Laouri M, Kravitz RL, French WI, Yang I, Milliken JC, Hilborne L, Wachsner R, Brook RH. (1997). Underuse of coronary revascularization procedures: Application of a clinical method. *Journal of the American College of Cardiology* 29:891-897.

La Veist TA, Nickerson KJ, Bowie JV. (2000). Attitudes about racism, medical mistrust, and satisfaction with care among African American and white cardiac patients. *Medical Care Research and Review* 57(Supplement 1):146-61.

 $Leigh\ WA, Lillie-Blanton\ M,\ Martinez\ RM,\ Collins\ KS.\ (1999).\ Managed\ care\ in\ three\ states:$

Experiences of low-income African-Americans and Hispanics. *Inquiry* 36(3):318-31.

Lillie-Blanton M, Brodie M, Rowland D, Altman D, McIntosh, M. (2000). Race, ethnicity, and the health care system: Public perceptions and experiences. *Medical Care Research and Review 57(1):218-235*.

Lillie-Blanton M, Martinez RM, Salganicoff A. (2001). Site of medical care: Do racial and ethnic differences persist? *Yale Journal of Health Policy, Law, and Ethics* 1(1):1-17.

Lowe RA, Chhaya S, Nasci K, Gavin LJ, Shaw K, Zwanger ML, Zeccardi JA, Daisey WC, Abbuhl SB, Feldman H, Berlin JA. (2001). Effect of ethnicity on denial of authorization for emergency department care by managed care gatekeepers. *Academic Emergency Medicine* 8(3):259-66.

Mackie DM, Hamilton DL, Susskind I, Rosselli F. (1996). Social psychological foundations of stereotype formation. In Macrae N, Stangor C, and Hewstone M (Eds.), *Stereotypes and stereotyping* (pp.41-78). New York: Guilford Press.

Manson A. (1988). Language concordance as a determinant of patient compliance and emergency room use in patients with asthma. *Medical Care* 26(12):1119-1128.

Mayberry RM, Mili F, Ofili E. (2000). Racial and ethnic differences in access to medical care.

Medical Care Research and Review 57(1):108-45.

McMahon LF, Wolfe RA, Huang S, Tedeschi P, Manning W, Edlund MJ. (1999). Racial and gender variation in use of diagnostic colonic procedures in the Michigan Medicare population. *Medical Care* 37(7):712-7.

Mitchell JB, McCormack LA. (1997). Time trends in late-stage diagnosis of cervical cancer:

Differences by race/ethnicity and income. Medical Care 35(12):1220-4.

Moore RD, Stanton D, Gopalan R, Chaisson RE. (1994). Racial differences in the use of drug therapy for HIV disease in an urban community. *New England Journal of Medicine* 330(11):763-8.

Morin R. (2001). Misperceptions cloud whites' view of blacks. The Washington Post, July 11, 2001.

Morrison RS, Wallenstein S, Natale DK, Senzel RS, Huang 1. (2000). "We don't carry that"—Failure of pharmacies in predominantly nonwhite neighborhoods to stock opioid analgesics. New England Journal of Medicine 342(14):1023-1026.

Mukamel DB, Murthy AS, Weimer DL. (2000). Racial differences in access to high-quality cardiac surgeons. *American Journal of Public Health* 90:1774-1777.

Petersen LA, Wright SM, Peterson ED, Daley J. (2002). Impact of race on cardiac care and outcomes in veterans with acute myocardial infarction. *Medical Care* 40(ISuppl):I-86-96.

Peterson ED, Shaw LK, DeLong ER, Pryor DB, Califf RM, Mark DB. (1997). Racial variation in the use of coronary-vascularization procedures: Are the differences real? Do they matter? *New England Journal of Medicine 336:480-6*.

Phillips KA, Mayer ML, Aday LA. (2000). Barriers to care among racial! ethnic groups under managed care. *Health Affairs* 19:65-75.

Rice T. (this volume). The impact of cost-containment efforts on racial and ethnic disparities in health care: A conceptualization.

Robert Wood Johnson Foundation. (2001). New survey shows language barriers causing many Spanish-speaking Latinos to skip care. Fact sheet presented at press briefing, December 12, 2001. Washington, DC.

Rodney M, Clasen C, Goldman G, Markert R, Deane D. (1998). Three evaluation methods of a community health advocate program. *Journal of Community Health* 23(5):371-381.

Schneider EC, Leape LL, Weissman JS, Piana RN, Gatsonis C, Epstein AM. (2001). Racial differences in cardiac revascularization rates: Does "overuse" explain higher rates among white patients? *Annals of Internal Medicine* 135(5):328-37.

Schulman KA, Berlin JA, Harless W, et al. (1999). The effect of race and sex on physicians' recommendations for cardiac catheterization. *New England Journal of Medicine 340:618-626*.

Sedlis SP, Fisher VI, Tice D, Esposito R, Madmon L, Steinberg EH. (1997). Racial differences in performance of invasive cardiac procedures in a Department of Veterans Affairs Medical Center. *Journal of Clinical Epidemiology* 50(8):899-901.

Shapiro MF, Morton SC, McCaffrey DF, Senterfitt JW, Fleishman JA, Perlman JF, Athey LA, Keesey JW, Goldman DP, Berry SH, Bozette SA. (1999). Variations in the care of HIV-infected adults in the United States: Results from the HIV Yost and Services Utilization Study. *Journal of the American Medical Association* 281:2305-75.

Smith DB. (1999). Health Care Divided: Race and Healing a Nation. Ann Arbor: The University of Michigan Press.

Tai-Seale M, Freund D, LoSasso A. (2001). Racial disparities in service use among Medicaid beneficiaries after mandatory enrollment in managed care: A difference-in-differences approach. *Inquiry* 38(1):49-59.

U.s. Commission on Civil Rights. (2001). Funding Federal Civil Rights Enforcement: 2000 and Beyond. Washington, DC: U.s. Commission on Civil Rights.

van Ryn M. (2002). Research on the provider contribution to race/ ethnicity disparities in medical care. *Medical Care* 40(1): 1-140-1-151.

van Ryn M, Burke J. (2000). The effect of patient race and socio-economic status on physician's perceptions of patients. *Social Science and Medicine* 50:813-828.

Weisse CS, Sorum PC, Sanders KN, Syat BL. (2001). Do gender and race affect decisions about pain management? Journal of Ceneral Internal Medicine 16(4)211-217.

Weitzman S, Cooper L, Chambless L, Rosamond W, Clegg L, Marcucci G, Romm F, White A. (1997). Gender, racial, and geographic differences in the performance of cardiac diagnostic and therapeutic procedures for hospitalized acute myocardial infarction in four states. *The American Journal of Cardiology* 79:722-6.

Wenneker MB, Epstein AM. (1989) Racial inequalities in the use of procedures for patients with ischemic heart disease in Massachusetts. *Journal of the American Medical Association* 261:253-7.

Williams DR. (1999). Race, socioeconomic status, and health: The added effects of racism and discrimination. *Annals of the New York Academy of Sciences* 896:173-88.

Williams DR, Rucker TD (2000). Understanding and addressing racial disparities in health care. *Health Care Financing Review* 21:75-90.

Williams JF, Zimmerman JW, Wagner DP, Hawkins M, Knaus W A. (1995). African-American and white patients admitted to the intensive care unit: Is there a difference in therapy and outcome? *Critical Care Medicine* 23(4):626-636.

Witmer A, Seifer SD, Finocchio L, Leslie J, O'Neil EH. (1995). Community health workers:

Integral members of the health care work force. American Journal of Public Health 85(8): 1055-1058.

Woloshin S, Bickell NA, Schwartz LM, Gany F, Welch HG. (1995). Language barriers in medicine in the United States. *Journal of the American Medical Association* 273(9):724-728.

Wood AJJ. (2001). Racial differences in the response to drugs-pointers to genetic differences. *New England Journal of Medicine 344:1393-1395*.

Yergan J, Food AB, LoGerfo JP, Diher P. Relationship between patient race and the intensity of hospital services. *Medical Care* 25:592-603.

Copyright © National Academy of Sciences. Permission is granted for this material to be shared for noncommercial, educational purposes, provided that this notice appears on the reproduced materials, the Web address of the online, full authoritative version is retained, and copies are not altered. To disseminate otherwise or to republish requires written permission from the National Academies Press.