

LACTATE DEHYDROGENASE ISOZYMES IN TILAPIINE FISHES (CICHLIDAE): TISSUE EXPRESSION AND GENETIC VARIABILITY PATTERNS

SHERIF H. EL-ALFY*, MOHAMED B. ABDELMORDY*, MOHAMED S. SALAMA**

*Zoology Department, Faculty of Science, Ain Shams University, Cairo, Egypt

**Entomology Department, Faculty of Science, Ain Shams University, Cairo, Egypt.

Abstract

The tissue specific LDH isozyme pattern is examined in three species of the economically important tilapiine fishes, *Oreochromis niloticus*, *O. aureus* and *Tilapia zillii*, using horizontal starch gel electrophoresis. Like the majority of vertebrates, tilapiine fishes exhibited the usual relative electrophoretic mobility of the A₄ and B₄-that is, B₄ carries a greater net negative charge. Isozyme of A₄ homotetramer was predominant in the skeletal muscle, while B₄ one was greatly expressed in heart muscle. Heteropolymeric isozymes of A and B composition were recognized. They were expressed in some organs and absent in others. Therefore, the present study indicated that tilapiine species could be considered 3-5 isozyme fishes. The total number of LDH isozymes of A and B composition is under genetic control of two loci, *Ldh-A* and *Ldh-B*. The LDH patterns of eye and brain displayed an additional band which was detected anodal to B₄. This anodal band is designated as LDH-C₄ isozyme. The latter isozyme is encoded in a third *Ldh* locus designated as *Ldh-C*. While the brain has heteropolymers of CA, CB and CAB polypeptide composition, the eye was the only organ that expressed the C₄ homopolymer. The present investigation showed that C gene was able to function but at lower levels in non-neural tissues. The LDH isozyme patterns of muscle, heart and eye were also compared among the three tilapiine species. The isozymatic LDH patterns in the eye of the examined species were similar except at *Ldh-C* locus. The genetically controlled restriction of the heterotetramers assembly along with the homology of the three *Ldh* loci among species was discussed.

Keywords: Lactate dehydrogenase, isozymes, tilapiine fishes, *Oreochromis*, *Tilapia*, electrophoresis.

Corresponding author: (Sherif H. El-Alfy Tel: 002-0161705032, Fax: 00202-6842123, E-mail: s.helmy23@yahoo.com)

INTRODUCTION

Lactate dehydrogenase (LDH: L-lactate, NAD⁺ oxidoreductase, EC 1.1.1.27) in higher vertebrates has been shown to exist in the catalytically active form as a tetramer of four polypeptide chains. LDH enzyme is involved in the reversible reaction of lactate oxidation and pyruvate reduction (Zietara and Skorkowski, 1991, 1993). Electrophoresis of tissue extracts, followed by the appropriate visualization procedure revealed five isozyme bands corresponding to the five possible tetrameric

combinations of two kinds of subunits (Appella and Markert, 1961). The occurrence of the five possible HM or AB tetramers in fishes is not as common as in higher vertebrates. Frequently either only the homopolymers A_4 and B_4 isozymes or these two plus A_2B_2 heteropolymer are all observed. In fact, it would appear that there is such a range of LDH isozymes in fishes that few generalizations can be made. In Petromyzontiformes, LDH is encoded by a single *Ldh-A* gene locus. In Myxiniformes, however, two loci *A* and *B* are present forming the two homotetramers A_4 and B_4 , and these remained in the other vertebrates (Whitt, 1984; Baldwin and Lake, 1987; Stock and Whitt, 1992). Certain fish species, particularly flatfish, apparently contain only one isozyme of LDH in most their tissues (Markert, 1968). Electrophoresis of tissue extracts of longnose and blacknose dace (*Rhinichthys cataractae* and *R. atratulus*) and some phenotypes of the carp and the barb (Cypriniformes) revealed five isozymes corresponding to tetramers of heart and muscle type subunits in birds and mammals (Clayton and Gee, 1969; Engel *et al.*, 1973; Beck *et al.*, 1983; Frankel, 1987; Gronczewska *et al.*, 2003). In Salmoniformes, multiple genes have been described both for *Ldh-A* and *Ldh-B*. A gene duplication has produced two polypeptides, B^- from *B* and A^- from *A* (Bailey *et al.*, 1976; Kettler and Whitt, 1986). The relative electrophoretic mobility of the A_4 and B_4 isozymes was reversed from the usual in some fishes like the smallmouth and largemouth bass, and in some Serrasalmidae (Characiformes) (Whitt *et al.*, 1971; Almeida-Val *et al.*, 1991). The reverse mobility pattern does not imply on a reverse actuation of LDH enzyme on tissue glycolytic metabolism (Almeida-Val *et al.*, 1991).

An LDH isozyme unique to teleostean fishes is the LDH- C_4 isozyme. This isozyme is presumably encoded in a third *Ldh* locus (Markert and Faulhaber, 1965; Whitt *et al.*, 1971; Coppes *et al.*, 1987). The *Ldh-C* gene was expressed in many tissues in primitive teleosts while in advanced teleosts, LDH- C_4 was restricted to few tissues and it was much more specialized (Zietara and Skorkoski, 1991; Almeida-Val and Val, 1993; Farias *et al.*, 1997). The LDH-C polypeptides are synthesized almost solely in the region of the nervous system involved in vision (Goldberg, 1965; Markert and Faulhaber, 1965; Frankel, 1982). The highest level of C_4 isozyme activity is found in the neural retinal cells, particularly in the inner segment region of the photoreceptor cells (Whitt, 1970b; Whitt and Booth, 1970; Quattro *et al.*, 1993). In species belonging to the orders Gadiformes and Cypriniformes, C_4 isozyme was active in the liver, with cathodic migration (Shaklee *et al.*, 1973; Champion *et al.*, 1975; Kettler and Whitt, 1986; Rao *et al.*, 1989; Zietara and Skorkowski, 1991). Basaglia (1991) demonstrated that the hepatic electropherograms for some Sparidae species (Perciformes) unexpectedly showed evidence of the C_4 isozyme with a high anodic

rate. Rehse and Davidson (1986) suggested a close homology between C subunits of teleosts and mammals based on amino acid composition data. Markert (1994) used a chicken B-actin promoter to drive the coding sequence of a human *Ldh-C* transgene in order to examine the physiological effects of LDH isozymes containing C subunits when functioning in somatic tissues. In this study, the differential LDH patterns in different tissues of three tilapiine species are examined and the possible genetic controls of different tetramer expressions are discussed. In addition, the genetic heterogeneity among the candidate species, based on LDH patterns, is investigated.

MATERIALS AND METHODS

Chemicals and animals

Most chemicals were obtained from Sigma Chemical Co. (St Louis, MO). Fishes utilized in this study comprised *Oreochromis niloticus*, *Oreochromis aureus* and *Tilapia zillii*. These tilapiine fishes were collected from the Nile River in the region between Giza and Maser El-Kadima, Egypt. The total number of collected adult fishes was about 13 *O. niloticus*, 8 *O. aureus* and 12 *T. zillii*. The live specimens were transported to the laboratory and carefully dissected. The tissues examined were the left lobe of liver (hepatopancreas), heart muscle, ovary, testis, brain, stomach, eye, kidney and skeletal muscle. The dissected organs were stored at -20 °C until use.

Homogenization and sample preparation

Tissue samples were diluted in a ratio 1:1 with a precooled grinding solution (0.6% saline solution) except for stomach, kidney and liver where dilution was 3-folds. Homogenization was achieved using a motor-driven Teflon pestle. Homogenates were frozen and thawed twice, and then centrifuged at 14,000 rpm for 30 min in a cooling microfuge. The supernatants were isolated and recentrifuged for another 30 min under the same condition. Clear supernatant was divided into aliquots of 50 µL and kept at -20 °C.

Electrophoresis

Horizontal starch gel electrophoresis was carried out according to May (1992) to separate LDH isozymes. 11% starch in Tris (0.03M) citrate (0.005M) buffer, pH 8.5 (gel buffer) was used. The tray buffer (pH 8.1) was made up of lithium hydroxide (0.06M) and boric acid (0.3M). Run was carried out at a potential gradient of 10 V/cm for 6-7 h at 6 °C.

Staining

Visualization of LDH was carried out according to May (1992) with a slight modification. Gel slices were incubated in the dark at 37 °C for 1-2 h in a solution contained 45 ml gel buffer (pH 8.5), 5 ml substrate, 10 mg nicotinamide adenine

dinucleotide (NAD), 10 mg nitro blue tetrazolium (NBT) and 2 mg phenazine methosulphate (PMS). LDH substrate was prepared according to Shaw and Prasad (1970). After staining, the gels were fixed in a solution consisted of a mixture of methanol, H₂O and glacial acetic acid (5:4:1), and were then photographed.

RESULTS

I. Tissue distribution of the LDH isozymes of *Oreochromis niloticus*.

The tissue specific LDH isozyme pattern is displayed in Fig. 1 and Table 1. The subunit composition of LDH isozymes was designated on the basis of their tissue specificity and electrophoretic mobility according to Whitt (1970b). Like the majority of vertebrates, tilapiine fishes exhibited the usual (not reversal) relative electrophoretic mobility of the A₄ and B₄ - that is, the B₄ carries a greater net negative charge. Isozyme of B₄ homotetramer was well expressed in all tissues with their greatest activity in heart, while isozyme of A₄ homotetramer was found in varying degrees in all tissues but predominates in extracts of skeletal muscle. Heteropolymeric isozymes of mobility intermediate to the A₄ and B₄ homopolymers were recognized in this tilapiine fish. Among heteropolymers is A₂B₂ isozyme which gave the greatest stain density in ovary and brain. The remaining heteropolymers are A₃B₁ and B₃A₁. They were present in some organs and absent in others. A₃B₁ isozyme form was seen in the eye, ovary and stomach. The other heteropolymer B₃A₁ was detected in liver, eye, ovary, kidney and stomach. The LDH patterns of eye and brain are of special interest, because additional bands were detected anodal to LDH-B₄ (heart type LDH) in these organs. These anodal bands are designated as LDH-C isozymes. While the brain has heteropolymers of CA and CAB polypeptide composition, the eye was the only organ that expressed the C₄ homopolymer in addition to A₂C₂ heteropolymer. Moreover, a weak activity for A₃C₁ was scored in stomach and kidney.

II. Tissue distribution of the LDH isozymes of *Oreochromis aureus*.

The tissue specific LDH isozyme pattern is shown in Fig. 2 and Table 1. As in the case of *O. niloticus*, isozyme of B₄ homotetramer of *O. aureus* was well expressed in all tissues with their greatest activity in heart. Isozyme of A₄ homotetramer showed weak to strong stain density in all organs tested. Very strong activity for A₄ isozyme was noticed in muscle extract. Heteropolymeric isozymes that lie between A₄ and B₄ homopolymers were also recognized in this tilapiine fish. The symmetrical heteropolymers, A₂B₂ isozymes were well expressed in all organs investigated except liver which gave rather a trace of activity. The asymmetrical heteropolymer B₃A₁ isozyme gave weak staining activity in all organs except liver which has no activity for this isozyme. The other heteropolymer A₃B₁ showed a strong activity in muscle. Most

of the remaining organs showed weak to moderate activity for the latter isozyme. The anodal isozymes of C subunit composition were restricted in their expression to eye and brain. Extracts of brain contained heteropolymers of CA, CB and CAB polypeptide composition, while extracts of eye contained CAB composition in addition to C₄ homopolymer.

III. Tissue distribution of the LDH isozymes of *Tilapia zillii*.

The LDH isozyme patterns are portrayed in Fig. 3 and Table 1. The homopolymer A₄ was found in all tissues examined except liver. The most pronounced activity for A₄ isozyme was seen in muscle. The other homopolymer B₄ which occupies the anodal terminal was expressed in all tissues investigated. Expression of B₄ in heart comes on the top since a very strong activity for this isozyme was obtained. Heteropolymeric isozymes of electrophoretic mobility intermediate to the A₄ and B₄ homopolymers were also recognized in this species. The symmetrical heteropolymer A₂B₂ was expressed in varying degrees in all tissues. The asymmetrical A₃B₁ isozyme was found in all organs studied except liver and heart. The remaining heteropolymer B₃A₁ could not be detected in any of the organs analyzed in *T. zillii*. Brain extracts contained heteropolymers of CA and CAB polypeptide composition, while eye extracts contained CA and CB composition in addition to C₄ homopolymer. Heart extract contained heteropolymers of CA and CAB composition which indicated that the C subunit was synthesized in that tissue.

IV. LDH-species specificity (interspecific variation)

The LDH isozyme patterns of muscle, heart and eye were compared among the three tilapiine species (Figs. 4 and 5). Each species has its own specific LDH pattern. The electrophoretic mobilities of LDH of muscle and heart extracts from the three tilapiine species were quite similar. The A₄ (with average cathodic rate) and B₄ (with average anodic rate) isozymes of *O. niloticus* appeared to be identical in electrophoretic mobilities to those of *O. aureus* and *T. zillii*. Thus these bands were considered as monomorphic bands. The isozymatic LDH patterns in the eye of the three tilapiine species studied were also similar except for C₄ isozymes. The mobility of the latter homotetramer was different on the genus level. In the two species belonging to the genus *Oreochromis*, the mobility was the same while it differed from that of the species belonging to the other genus, *Tilapia*. The C₄ homotetramer of *O. niloticus* and *O. aureus* had a slightly faster mobility than that of *T. zillii*. Thus C₄ band was considered as polymorphic band. Concerning the heteropolymeric isozymes of CA and CB composition, they were considered in turn as polymorphic bands.

DISCUSSION

In most vertebrates, LDH is encoded by two gene loci, *Ldh-A* and *Ldh-B* which synthesize two subunits, A and B or M and H (Appella and Markert, 1961; Markert, 1968; Karlsson and Larsson, 1971; Champion *et al.*, 1975; Holt and Leibel, 1987; Haggblom *et al.*, 1988; Heinova and Blahovec, 1994; Tsoi and Li, 1994; Abdelmordy, 1999). These subunits are associated in the cytoplasm and produce five different tetramers: two homotetramers A_4 and B_4 , and three heterotetramers A_3B_1 , A_2B_2 and A_1B_3 . These tetramers have different distribution and different kinetic and physicochemical properties (Appella and Markert, 1961; Markert, 1968; Wu *et al.*, 1993; Schulte *et al.*, 2000; Chippari-Gomes *et al.*, 2005; Ishibashi *et al.*, 2007). Markert and Faulhaber (1965) found that out of 30 species of fishes, only three which produced all the five possible HM tetramers. The present results indicated that *T. zillii* is the only tilapiine fish species which did not show the five possible HM tetramers. The five possible HM tetramers were only detected in eye, ovary and stomach of *O. niloticus* and in most tissues of *O. aureus*. Markert and Faulhaber (1965) found that the electrophoretic mobility of the LDH isozymes of 10 out of 30 fish species with reversed heart-muscle LDH mobility. The same relative mobility of LDH isozymes was observed in some Poeciliidae, Percichthyidae and Serrasalminidae (Frankel, 1982; Almeida-Val *et al.*, 1991; Xia *et al.*, 1992). The present results indicated that the electrophoretic mobility of the LDH isozymes of tilapiine fishes is quite similar to what has been described for higher vertebrates and most teleosts (Battellino and Blanco, 1970; Champion *et al.*, 1975; Holt and Leibel, 1987; Rao *et al.*, 1989; Basaglia, 1991; Heinova and Blahovec, 1994; Abdelmordy, 1999). More specifically, the homotetramer A_4 has a lower negative charge and thus lower mobility (cathodic). B_4 has an intermediate negative charge and thus faster mobility (anodic). In the three tilapiine fishes, *Ldh-A* and *Ldh-B* were all expressed together in all tissues examined. A_4 homotetramer was predominant in the skeletal muscle (anaerobic tissue), while B_4 was expressed mainly in heart muscle (aerobic tissue). This distribution reflects functional characteristics of the isozymes that are well established since it involves regulatory genes and represents preferential metabolism of each tissue.

Sensabaugh and Kaplan (1972) have reported that LDH hybridization in fish is not random as it is for LDH of higher vertebrates, while Beck *et al.* (1983) have shown that the different LDH subunits associate at random to form the observed isozymes. It has been suggested that the A and B subunits of certain teleosts do not associate *in vivo* or *in vitro* producing 2 banded LDH pattern as a result of genetically controlled restriction on their assembly (Whitt, 1970a; Frankel, 1982). According to Markert (1984) and Zawadzki *et al.* (2001), restriction in copolymerization between paralogous

Ldh-A and *Ldh-B* gene products is common in fishes, amphibians and reptiles. It was assumed that the restricted (2-4) banding pattern may be explained by the evolutionary divergence between the two *Ldh* loci after genome duplication (Markert and Faulhaber, 1965; Whitt, 1970a, b). Furthermore, several reports have suggested that the occurrence of restricted LDH patterns in the majority of advanced teleosts may reflect a strong pressure for selecting subunits with low copolymerization abilities (Morizot and Siciliano, 1979; Philipp *et al.*, 1979). Almeida-Val *et al.* (1992) have shown that the isozyme number made up from A and B subunits within the family Serrasalmidae exhibits a non-phylogenetic variation, and the existence of different isozyme number can be explained by suitable differences in electric charges of subunits A and B. So in this case, the LDH isozyme number apparently is not determined by natural selection.

The present studies on tilapiine fishes indicated that these species could be considered 3-5 isozyme fishes. These restricted 3-5 banded patterns could be explained on the basis of either one of the two following alternatives: the first is the evolutionary divergence between the two *Ldh* loci after genome duplication. This divergence could provide molecular instability for co-polymerization of one homodimer (A_2 or B_2) with two different monomers (A and B subunits) leading to absence of the two or one of the asymmetrical heterotetramers. A_3B_1 and B_3A_1 were absent in liver and heart of *T. zillii*, and in brain, muscle and heart of *O. niloticus*. In the latter species, only A_3B_1 was absent in kidney and liver. On the other hand, B_3A_1 was absent in all organs of *T. zillii* and in liver of *O. aureus*. The result is the formation of LDH patterns of less than 5 isozymes in some organs of *O. niloticus* and *O. aureus*, and in all organs of *T. zillii*. The second explanation is the gene product of B_4 homotetramer is relatively not enough favoring the occurrence of the A_3B_1 heterotetramer in case of *T. zillii*. On the contrary, absence of A_3B_1 in some organs of *O. niloticus* reflects that gene product of A_4 in these organs is relatively not enough since it involves regulatory mechanisms favoring the absence of A_3B_1 . Ahmad and Hasnain (2005) reported that the contribution of heterotetramers depends on the levels of LDH homotetramers. Absence of the two asymmetrical heterotetramers A_3B_1 and B_3A_1 in heart, muscle and brain of *O. niloticus* although good expression of both A_4 and B_4 homotetramers may provide a supporting evidence to favor the first explanation in this case. The present results are not in common to what has been described by Rao *et al.* (1989). They have shown that *O. niloticus* contained only 1 or 2 LDH bands in their organs.

The members of tilapiine fishes showed a third homotetramer (LDH- C_4) that carries a strong negative charge and hence migrates rapidly to the anode during electrophoresis. The activity of the C type isozyme was found in eye and brain

extracts. This observation is consistent with most teleosts (Whitt, 1970b; Frankel, 1982; Holt and Leibel, 1987; Basaglia, 1991; Farias *et al.*, 1997). Isozyme C₄ is encoded in a third *Ldh* locus designated *Ldh-C* (Markert and Faulhaber, 1965; Markert, 1968; Whitt *et al.*, 1971; Coppes *et al.*, 1987; Farias *et al.*, 1997). The *Ldh-C* product is probably involved in the visual metabolism (Frankel, 1982). The present investigation showed that the C gene was able to function in non-neural tissues such as heart in *T. zillii*, kidney and stomach in *O. niloticus*. This observation is consistent with Whitt *et al.* (1971) and Basaglia (1991, 2002). However, the level of C polypeptide synthesis was lower in these tissues than in the eye and brain. According to Almeida-Val and Val (1993), the regulatory pattern of *Ldh-C* gene appears to be the result of a high selective pressure upon an ancient gene.

The heteropolymeric isozymes intermediate to the B₄ and C₄ isozymes in the eye of the bass fish appeared to be composed mainly of B and C subunits rather than A and C subunits (Whitt *et al.*, 1971; Philipp *et al.*, 1979). The eye of *Phallichthys amates* contains all heteropolymers of C subunit composition (CA, CB and CAB) (Frankel, 1982). The present investigation showed that eye and brain of tilapiine fishes run in common with *P. amates* in that respect. This proves that there is no isolation of C and A as well as B subunit synthesis. In contrast, in birds and mammals, during the time the C gene is turned on, the A and B genes are turned off. This coordinated behavior is not applicable to the LDH isozymes in many fish tissues.

Comparison of the LDH isozyme patterns of muscle, heart and eye between the three tilapiine species demonstrates that the lactate dehydrogenase isozymes are species-specific. Nonetheless, large similarities in terms of mobility and intensity exist in the patterns of these species, particularly those belonging to the same genus (*Oreochromis*). The electrophoretic mobilities of LDH-A₄ and LDH-B₄ from the three tilapiine species were quite similar indicating the homology of *Ldh-A* and *Ldh-B* genes among tilapiine fishes. The mobility of the C₄ in the eye of the three species was different on the genus level. In the two species belonging to genus *Oreochromis* the C₄ homotetramer migrated faster than the corresponding isozyme of the genus *Tilapia*. It must therefore carry a greater net negative charge than the C₄ isozyme of *T. zillii*. This in turn will be reflected on the mobility of heteropolymeric isozymes of CA, CB and CAB composition. It may be assumed that the *Ldh-C* locus was monomorphic at a time and the polymorphism was taken place prior to the divergence of the *Oreochromis* and *Tilapia*. Thus the evolution occurred more rapidly on an ancient gene (*Ldh-C* locus) than on the relatively more recent genes (*Ldh-A* and *Ldh-B* loci).

Table 1. Distribution and rate of activity of LDH isozymes in different organs of *Oreochromis niloticus*, *O. aureus* and *Tilapia zillii*.

Organs	Species	LDH isozymes											
		A ₄	A ₃ B ₁	A ₂ B ₂	B ₃ A ₁	B ₄	A ₃ C ₁	A ₂ B ₁ C ₁	A ₂ C ₂	A ₁ B ₁ C ₂	B ₂ C ₂	B ₁ C ₃	C ₄
Liver	<i>O. niloticus</i>	+	-	(+)	++	++	-	-	-	-	-	-	-
	<i>O. aureus</i>	+	+	(+)	-	++	-	-	-	-	-	-	-
	<i>T. zillii</i>	(+)	-	+	-	++	-	-	-	-	-	-	-
Heart	<i>O. niloticus</i>	++	-	++	-	++	-	-	-	-	-	-	-
	<i>O. aureus</i>	+	(+)	++	(+)	++	-	-	-	-	-	-	-
	<i>T. zillii</i>	+	-	++	-	++	++	+++	-	-	-	-	-
Brain	<i>O. niloticus</i>	++	-	++	-	++	-	-	++	++	-	-	-
	<i>O. aureus</i>	++	+	++	+	++	+	-	+	++	++	-	-
	<i>T. zillii</i>	++	++	++	-	++	++	-	++	++	-	-	-
Stomach	<i>O. niloticus</i>	+	+	++	+	++	+	-	-	-	-	-	-
	<i>O. aureus</i>	(+)	-	++	+	++	-	-	-	-	-	-	-
	<i>T. zillii</i>	++	++	++	-	++	-	-	-	-	-	-	-
Eye	<i>O. niloticus</i>	++	++	++	++	++	-	-	+	-	-	-	++
	<i>O. aureus</i>	++	++	++	+	++	-	-	-	+	-	-	++
	<i>T. zillii</i>	++	++	++	-	++	-	-	+	-	+	++	++
Kidney	<i>O. niloticus</i>	++	-	++	+	++	+	-	-	-	-	-	-
	<i>O. aureus</i>	++	-	++	+	++	-	-	-	-	-	-	-
	<i>T. zillii</i>	++	+	++	-	++	-	-	-	-	-	-	-
Muscle	<i>O. niloticus</i>	++	-	++	-	++	-	-	-	-	-	-	-
	<i>O. aureus</i>	++	++	++	+	++	-	-	-	-	-	-	-
	<i>T. zillii</i>	++	++	+	-	++	-	-	-	-	-	-	-
Ovary	<i>O. niloticus</i>	++	+	++	+	++	-	-	-	-	-	-	-
	<i>O. aureus</i>	?	?	?	?	?	?	?	?	?	?	?	?
	<i>T. zillii</i>	++	++	++	-	++	-	-	-	-	-	-	-
Testis	<i>O. niloticus</i>	?	?	?	?	?	?	?	?	?	?	?	?
	<i>O. aureus</i>	?	?	?	?	?	?	?	?	?	?	?	?
	<i>T. zillii</i>	++	+	+	-	++	-	-	-	-	-	-	-

(+) = trace of activity; + = weak activity; ++ = moderate activity; +++ = strong activity; ++++ = very strong activity; - = no activity; ? = not tested.




Fig. 1. Tissue distribution of LDH isozymes of *O. niloticus* (l, liver; e, eye; o, ovary; h, heart; m, muscle; b, brain; k, kidney; s, stomach).




Fig. 2. Tissue distribution of LDH isozymes of *O. aureus* (h, heart; b, brain; s, stomach; e, eye; k, kidney; m, muscle; l, liver).




Fig. 3. Tissue distribution of LDH isozymes of *T. zillii* (l, liver; h, heart; o, ovary; b, brain; s, stomach; e, eye; k, kidney; m, muscle; t, testis).




Fig. 4. LDH patterns in heart (a) and skeletal muscle (b) of tilapia fishes (1, *O. niloticus*; 2, *O. aureus*; 3, *T. zilli*).




Fig. 5. LDH patterns in eye of tilapia fishes (1, *O. niloticus*; 2, *O. aureus*; 3, *T. zilli*).

REFERENCES

1. Abdelmordy, M. 1999. Organ distribution of lactate and malate dehydrogenase isozymes in two species of the genus *Gerbillus* (Rodentia). *Biologia*, Bratislava, 54: 325-332.
2. Ahmad, R. and A. Hasnain. 2005. Ontogenetic changes and developmental adjustments in lactate dehydrogenase isozymes of an obligate air-breathing fish *Channa punctatus* during deprivation of air access. *Comp. Biochem. Physiol.*, 140B: 271-278.
3. Almeida-Val, V. M. F. and A. L. Val. 1993. Evolutionary trends of LDH isozymes in fishes. *Comp. Biochem. Physiol.*, 105B: 21-28.
4. Almeida-Val, V. M. F., M. L. B. Schwantes and A. L. Val. 1991. LDH isozymes in Amazon fish-II. Temperature and pH effects on LDH kinetic properties from *Mylossoma duriventris* and *Colossoma macropomum* (Serrasalminidae). *Comp. Biochem. Physiol.*, 98B: 79-86.
5. Almeida-Val, V. M. F., M. N. Paula-Silva, M. C. M. Caraciolo, L. S. B. Mesquita, I. P. Farias and A. L. Val. 1992. LDH isozymes in Amazon fishes-III. Distribution patterns and functional properties in Serrasalminidae (Teleostei: Ostariophysi). *Comp. Biochem. Physiol.*, 103B: 119-125.
6. Appella, E., and C. L. Markert. 1961. Dissociation of lactate dehydrogenase into subunits with guanidine hydrochloride. *Biochem. Biophys. Res. Commun.*, 6: 171-176.
7. Bailey, G. S., H. Tsuyki and A. C. Wilson. 1976. The number of genes for lactate dehydrogenase in salmonid fishes. *J. Fish. Res. Bd. Canada*, 33: 760-767.
8. Baldwin, J. and P. S. Lake. 1987. Lactate dehydrogenase homopolymer of hagfish heart and the single lactate dehydrogenase of lampreys display greater immunochemical similarity to LDHC₄ than to LDHB₄ of teleost fish. *J. Exp. Zool.*, 242: 99-102.
9. Basaglia, F. 1991. Lactate dehydrogenase isozymes and their genetic variation in fifteen Sparidae species (Perciformes, Teleostei). *Comp. Biochem. Physiol.*, 98B: 1-8.
10. Basaglia, F. 2002. Multilocus isozyme systems in African lungfish, *Protopterus annectens*: distribution, differential expression and variation in dipnoans. *Comp. Biochem. Physiol.*, 131B: 89-102.
11. Battellino, L. J. and A. Blanco. 1970. Catalytic properties of the lactate dehydrogenase isozyme "X" from mouse testis. *J. Exp. Zool.*, 174: 173-186.

12. Beck, M. L., C. J. Biggers and H. K. Dupree. 1983. Electrophoretic analysis of protein systems of *Ctenopharyngodon idella* (Val.), *Hypophthalmichthys nobilis* (Rich.) and their F1 triploid hybrid. J. Fish. Biol., 22: 603-611.
13. Champion, M. J., J. B. Shaklee and G. S. Whitt. 1975. Developmental genetics of teleost isozymes. In: Markert, C. L., (Ed.), Developmental Genetics. Academic Press, New York. pp. 417-437.
14. Chippari-Gomes, A. R., L. C. Gomes, N. P. Lopes, A. L. Val and V. M. F. Almeida-Val. 2005. Metabolic adjustments in two Amazonian cichlids exposed to hypoxia and anoxia. Comp. Biochem. Physiol., 141B: 347-355.
15. Clayton, J. W. and J. H. Gee. 1969. Lactate dehydrogenase isozymes in longnose and blacknose dace (*Rhinichthys cataractae* and *R. atratulus*) and their hybrid. J. Fish. Res. Bd. Canada, 26: 3049-3053.
16. Coppes, Z. L., M. L. Schwantes and A. R. Schwantes. 1987. Adaptive features of enzymes from family Sciaenidae-III. Studies on lactate dehydrogenase (LDH) of fishes from the south coast of Uruguay. Comp. Biochem. Physiol., 88B: 1005-1012.
17. Engel, W., J. Schmidtke, W. Vogel and U. Wolf. 1973. Genetic polymorphism of lactate dehydrogenase isozymes in the carp (*Cyprinus carpio*) apparently due to a "null allele". Biochem. Genet. 8: 281-289.
18. Farias, I. P., M. N. Paula-Silva and V. M. F. Almeida-Val. 1997. No co-expression of LDH-C in amazon cichlids. Comp. Biochem. Physiol., 117B: 315-319.
19. Frankel, J. S. 1982. Lactate dehydrogenase specificity and subunit assembly in neural tissues of the teleost *Phallichthys amates*. Experientia. 38: 673-674.
20. Frankel, J. S. 1987. Lactate dehydrogenase isozymes of the island barb, *Barbus oligolepis* (Cypriniformes, Teleostei): their characterization and ontogeny. Comp. Biochem. Physiol., 87B: 581-585.
21. Goldberg, E. 1965. Lactate dehydrogenase in trout: evidence for a third subunit. Ibid., 148: 391-392.
22. Gronczewska, J., M. S. Zietara, A. Biegñiewska and E. F. Skorkowski. 2003. Enzyme activities in fish spermatozoa with focus on lactate dehydrogenase isoenzymes from herring *Clupea harengus*. Comp. Biochem. Physiol., 134B: 399-406.
23. Haggblom, L., R. C. Terwilliger and N. B. Terwilliger. 1988. Changes in myoglobin and lactate dehydrogenase in muscle tissues of a diving bird, the pigeon guillemot, during maturation. Comp. Biochem. Physiol., 91B: 273-277.
24. Heinova, D. and J. Blahovec. 1994. Lactate dehydrogenase isozymes in mammalian and chicken serum. Vet. Med. Praha, 39: 75-84.

25. Holt, R. W., and W. S. Leibel. 1987. Coexpression of distinct eye- and liver-specific LDH isozymes in cichlid fish. *J. Exp. Zool.*, 244: 337-343.
26. Ishibashi, Y., T. Kotaki, Y. Yamada and H. Ohta. 2007. Ontogenic changes in tolerance to hypoxia and energy metabolism of larval and juvenile Japanese flounder *Paralichthys olivaceus*. *J. Experimental Marine Biology and Ecology*, 352: 42-49.
27. Karlsson, B. W., and G. B. Larsson. 1971. Lactic and malic dehydrogenases and their multiple molecular forms in the Mongolian gerbil as compared with the rat, mouse and rabbit. *Comp. Biochem. Physiol.*, 40B: 93-96.
28. Kettler, M. K. and G. S. Whitt. 1986. An apparent progressive and recurrent evolutionary restriction in tissue expression of a gene, the lactate dehydrogenase-C gene, within a family of bony fish (Salmoniformes: Umbridae). *J. Mol. Evol.*, 23: 95-107.
29. Markert, C. L. 1968. The molecular basis for isozymes. *Ann. N. Y. Acad. Sci.*, 151: 14-40.
30. Markert, C. L. 1984. Lactate dehydrogenase-biochemistry and function of lactate dehydrogenase. *Cell. Biochem. Funct.*, 2: 131-134.
31. Markert, C. L. 1994. Transgenic creation of novel isozyme systems for challenging and studying the physiology and development of organisms. In: Markert, C. L., J. G. Scandalios, H. A. Lim and O. L. Serov, (Eds.), *Isozymes: Organization and Roles in Evolution, Genetics and Physiology*. The Seventh International Congress on Isozymes. World Scientific, Singapore. pp. 3-12.
32. Markert, C. L. and I. Faulhaber. 1965. Lactate dehydrogenase isozyme patterns of fish. *J. Exp. Zool.*, 159: 319-332.
33. May, B. 1992. Starch gel electrophoresis of allozymes. In: Hoelzel, A. R., (Ed.), *Molecular Genetic Analysis of Populations, a Practical Approach*. IRL Press, Oxford. pp. 1-27.
34. Morizot, D. C. and M. J. Siciliano. 1979. Polymorphisms, linkage and mapping of four enzyme loci in the fish genus *Xiphophorus* (Poeciliidae). *Genetics*, 93: 947-960.
35. Philipp, D. P., W. F. Childers and G. S. Whitt. 1979. Evolution of patterns of differential gene expression: a comparison of the temporal and spatial patterns of isozyme locus expression in two closely related fish species (northern largemouth bass, *Micropterus salmoides* and smallmouth bass, *micropterus dolomieu*). *J. Exp. Zool.*, 210: 473-488.

36. Quattro, J. M., H. A. Woods and D. A. Powers. 1993. Sequence analysis of teleost retina-specific lactate dehydrogenase C: evolutionary implications for the vertebrate lactate dehydrogenase gene family. *Proc. Natl. Acad. Sci.*, 90: 242-246.
37. Rao, M. R., B. K. Padhi and A. R. Khuda-Bukhsh. 1989. Lactate dehydrogenase isozymes in fifty-two species of teleostean fishes: taxonomic significance of *Ldh-C* gene expression. *Biochem. Syst. Ecol.*, 17: 69-76.
38. Rehse, P. H. and W. S. Davidson. 1986. Evolutionary relationship of a fish C type lactate dehydrogenase to other vertebrate lactate dehydrogenase isozymes. *Can. J. Fish. Aquat. Sci.*, 43: 1045-1051.
39. Schulte, P. M., H. C. Glemet, A. A. Fiebig and D. A. Powers. 2000. Adaptive variation in lactate dehydrogenase-B gene expression: Role of a stress-responsive regulatory element. *Proc. Natl. Acad. Sci.*, 97: 6597-6602.
40. Sensabaugh, G. F. Jr., and N. O. Kaplan. 1972. A lactate dehydrogenase specific to the liver of gadoid fish. *J. Biol. Chem.*, 247: 585-593.
41. Shaklee, J. B., K. L. Kepes and G. S. Whitt. 1973. Specialized lactate dehydrogenase isozymes: the molecular and genetic basis for the unique eye and liver LDHs of teleost fishes. *J. Exp. Zool.*, 185: 217-240.
42. Shaw, C. R. and R. Prasad. 1970. Starch gel electrophoresis of enzymes- a compilation of recipes. *Biochem. Genet.*, 4: 297-320.
43. Stock, D. W. and G. S. Whitt. 1992. Evolutionary implications of the cDNA sequence of the single lactate dehydrogenase of a lamprey. *Proc. Natl. Acad. Sci.*, 89: 1799-1803.
44. Tsoi, S. C. M. and S. S. L. Li. 1994. The nucleotide and deduced amino-acid sequences of a cDNA encoding lactate dehydrogenase from *Caenorhabditis elegans*: the evolutionary relationships of lactate dehydrogenases from mammals, birds, amphibian, fish, nematode, plants, bacteria, mycoplasma and plasmodium. *Biochem. Biophys. Res. Commun.*, 205: 558-564.
45. Whitt, G. S. 1970a. Directed assembly of polypeptides of the isozymes of lactate dehydrogenase. *Arch. Biochem. Biophys.*, 138: 352-354.
46. Whitt, G. S. 1970b. Developmental genetics of the lactate dehydrogenase isozymes of fish. *J. Exp. Zool.*, 175: 1-36.
47. Whitt, G. S. 1984. Genetic, developmental and evolutionary aspects of the lactate dehydrogenase isozyme system. *Cell Biochem. Funct.*, 2: 134-139.
48. Whitt, G. S., and G. M. Booth. 1970. Localization of lactate dehydrogenase activity in the cells of the fish (*Xiphophorus helleri*) eye. *J. Exp. Zool.*, 174: 215-224.

49. Whitt, G. S., W. F. Childers and T. E. Wheat. 1971. The inheritance of tissue-specific lactate dehydrogenase isozymes in interspecific bass (*Micropterus*) hybrids. *Biochem. Genet.*, 5: 257-273.
50. Wu, T., D. Xia and H. Wang. 1993. Purification and immunochemical analysis of lactate dehydrogenase (LDH) isozymes of grass carp (*Ctenopharyngodon idella*). *Aquaculture*, 110: 41-50.
51. Xia, D., T. Wu and H. Wang. 1992. Differential gene expression for lactate dehydrogenase of mandarin fish (*Siniperca chuatsi*). *Aquaculture*, 108: 207-214.
52. Zawadzki, C. H., M. F. P. Machado and E. Renesto. 2001. Differential expression for tissue-specific isozymes in three species of *Hypostomus* Lacepede, 1803 (Teleostei: Loricariidae). *Biochem. Syst. Ecol.*, 29: 911-922.
53. Zietara, M. S. and E. F. Skorkowski. 1991. Unusual expression of the threespine stickleback (*Gasterosteus aculeatus*) lactate dehydrogenase isoenzymes and partial characterization of purified LDH-A₄. *Comp. Biochem. Physiol.*, 99B: 51-56.
54. Zietara, M. S. and E. F. Skorkowski. 1993. Purification and properties of the heart type lactate dehydrogenase of the cod (*Gadus morhua*) from the Baltic sea: comparison with LDH-A₄ and LDH-C₄. *Comp. Biochem. Physiol.*, 105B: 349-356.

