

INDEPENDENCE IN ABSTRACT ELEMENTARY CLASSES

SEBASTIEN VASEY

ABSTRACT. We study general methods to build forking-like notions in the framework of tame abstract elementary classes (AECs) with amalgamation. We show that whenever such classes are categorical in a high-enough cardinal, they admit a good frame: a forking-like notion for types of singleton elements.

Theorem 0.1 (Superstability from categoricity). Let K be a $(< \kappa)$ -tame AEC with amalgamation. If $\kappa = \beth_\kappa > \text{LS}(K)$ and K is categorical in a $\lambda > \kappa$, then:

- K is stable in all cardinals $\geq \kappa$.
- K is categorical in κ .
- There is a type-full good λ -frame with underlying class K_λ .

Under more locality conditions, we prove that the frame extends to a global independence notion (for types of arbitrary length).

Theorem 0.2 (A global independence notion from categoricity). Let K be a densely type-local, fully tame and type short AEC with amalgamation. If K is categorical in unboundedly many cardinals, then there exists $\lambda \geq \text{LS}(K)$ such that $K_{\geq \lambda}$ admits a global independence relation with the properties of forking in a superstable first-order theory.

Modulo an unproven claim of Shelah, we deduce that Shelah's categoricity conjecture (without assuming categoricity in a successor cardinal) follows from the weak generalized continuum hypothesis and the existence of unboundedly many strongly compact cardinals.

Date: April 1, 2015

AMS 2010 Subject Classification: Primary 03C48. Secondary: 03C45, 03C52, 03C55, 03C75, 03E55.

Key words and phrases. Abstract elementary classes; Forking; Independence; Classification theory; Stability; Good frames; Categoricity; Superstability; Tameness.

The author is supported by the Swiss National Science Foundation.

CONTENTS

1. Introduction	2
2. Preliminaries	7
3. Independence relations	13
4. Some independence calculus	23
5. Skeletons	29
6. Generating an independence relation	33
7. Weakly good independence relations	37
8. Good independence relations	44
9. Canonicity	48
10. Superstability	51
11. Domination	59
12. A fully good long frame	67
13. Extending the base and right hand side	71
14. Extending the left hand side	74
15. The main theorems	81
References	84

1. INTRODUCTION

Independence (or forking) is one of the central notions of model theory. In the first-order setup, it was introduced by Shelah [She78] and is one of the main devices of his book. One can ask whether there is such a notion in the nonelementary context. In homogeneous model theory, this was investigated in [HL02] for the superstable case and [BL03] for the simple and stable cases. Some of their results were later generalized by Hyttinen and Kesälä [HK06] to tame and \aleph_0 -stable finitary abstract elementary classes¹. What about general² abstract elementary classes? There we believe the answer is still a work in progress.

¹Note that by [HK06, Theorem 4.11], such classes are actually fully $(< \aleph_0)$ -tame and short.

²For a discussion of how the framework of tame AECs compare to other non first-order frameworks, see the introduction of [Vasb].

Already in [She99, Remark 4.9.1] it was asked by Shelah whether there is such a notion as forking in AECs. In his book on AECs [She09], the central concept is that of a good λ -frame (a local independence notion for types of singletons) and some conditions are given for their existence. Shelah’s main construction (see [She09, Theorem II.3.7]) uses categoricity in two successive cardinals and non-ZFC principles like the weak diamond³. It has been suggested⁴ that replacing Shelah’s strong local model-theoretic hypotheses by the global hypotheses of amalgamation and tameness (a locality property for types introduced by Grossberg and VanDieren [GV06b]) should lead to ZFC theorems and simpler proofs. Furthermore, one can argue that any “reasonable” AEC should be tame and have amalgamation, see for example the discussion in Section 5 of [BG], and the introductions of [Bon14b] or [GV06b]. Examples of the use of tameness and amalgamation include [BKV06] (an upward stability transfer), [Lie11] (showing that tameness is equivalent to a natural topology on Galois types being Hausdorff), [GV06c] (an upward categoricity transfer theorem which Boney [Bon14b] used to prove that Shelah’s categoricity conjecture for a successor follows from unboundedly many strongly compact) and [Bon14a, BVb, Jarb], showing that good frames behave well in tame classes.

In [Vas], we constructed good frames in ZFC using global model-theoretic hypotheses: tameness, amalgamation, and categoricity in a cardinal of high-enough cofinality. However we were unable to remove the assumption on the cofinality of the cardinal or to show that the frame was ω -successful, a key technical property of frames. Both in Shelah’s book and in [Vas], the question of whether there exists a *global* independence notion (for longer types) was also left open. In this paper, we continue working in ZFC with tameness and amalgamation, and make progress toward these problems. Regarding the cofinality of the categoricity cardinal, we show (Theorem 10.16):

Theorem 1.1. Let K be a $(< \kappa)$ -tame AEC with amalgamation. If $\kappa = \beth_\kappa > \text{LS}(K)$ and K is categorical in a $\lambda > \kappa$, then there is a type-full good λ -frame with underlying class K_λ .

³Shelah claims to construct a good frame in ZFC in [She09, Theorem IV.4.10] but he has to change the class and still uses the weak diamond to show his frame is ω -successful.

⁴The program of using tameness and amalgamation to prove Shelah’s results in ZFC is due to Rami Grossberg and dates back to at least [GV06b], see the introduction there.

As a consequence, the class K above has several superstable-like properties: for all $\mu \geq \lambda$, K is stable⁵ in μ (this is also part of Theorem 10.16) and has a unique limit model of cardinality μ (by e.g. [BVb, Theorem 1.1] and Remark 2.26). The proof of Theorem 1.1 also yields a downward categoricity transfer⁶:

Theorem 1.2. Let K be a $(< \kappa)$ -tame AEC with amalgamation. If $\kappa = \beth_\kappa > \text{LS}(K)$ and K is categorical in a $\lambda > \kappa$, then K is categorical in κ .

The construction of the good frame in the proof of Theorem 1.1 is similar to that in [Vasa] but uses local character of coheir (or $(< \kappa)$ -satisfiability) rather than splitting. A milestone study of coheir in the nonelementary context is [MS90], working in classes of models of an $L_{\kappa, \omega}$ -sentence, κ a strongly compact cardinal. Makkai and Shelah's work was generalized to fully tame and short AECs in [BG], and some results were improved in [Vasb]. Building on these works, we are able to show that under the assumptions above, coheir has enough superstability-like properties to apply the methods of [Vasa], and obtain that coheir restricted to types of length one in fact induces a good frame.

Note that coheir is a candidate for a global independence relation. In fact, one of the main result of [BGKV] is that it is canonical: if there is a global forking-like notion, it must be coheir. Unfortunately, the paper assumes that coheir has the extension property, and it is not clear that it is a reasonable assumption. Here, we prove that coheir is canonical without this assumption (Theorem 9.3). We also obtain results on the canonicity of good frames. For example, any two type-full good λ -frames with the same *categorical* underlying AEC must be the same (Theorem 9.7). This answers several questions asked in [BGKV].

Using that coheir is global and (under categoricity) induces a good frame, we can use more locality assumptions to get that the good frame is ω -successful (Theorem 15.6):

⁵The downward stability transfer from categoricity is an early result of Shelah [She99, Claim 1.7], but the upward transfer is new and improves on [Vasa, Theorem 7.5]. In fact, the proof here is new even when K is the class of models of a first-order theory.

⁶[MS90, Conclusion 5.1] proved a similar conclusion under stronger assumptions (namely that K is the class of models of an $L_{\kappa, \omega}$ sentence, κ a strongly compact cardinal).

Theorem 1.3. Let K be a fully $(< \kappa)$ -tame and short AEC. If $\text{LS}(K) < \kappa = \beth_\kappa < \lambda = \beth_\lambda$, $\text{cf}(\lambda) \geq \kappa$, and K is categorical in a $\mu \geq \lambda$, then there exists an ω -successful type-full good λ -frame with underlying class K_λ .

We believe that the locality hypotheses in Theorem 1.3 are reasonable: they follow from large cardinals [Bon14b] and slightly weaker assumptions can be derived from the existence of a global forking-like notion, see the discussion in Section 15.

Theorem 1.3 can be used to build a global independence notion (Theorem 15.1 formalizes Theorem 0.2 from the abstract). Unfortunately we assume one more locality hypothesis (dense type-locality) there, but we suspect it can be removed, see the discussion in Section 15. Without dense type-locality, one still obtains an independence relation for types of length less than or equal to λ (see Theorem 15.6).

These results bring us closer to solving one of the main test questions in the classification theory of abstract elementary classes⁷:

Conjecture 1.4 (Shelah’s categoricity conjecture). If K is an AEC that is categorical in unboundedly⁸ many cardinals, then K is categorical on a tail of cardinals.

The power of ω -successful frames comes from Shelah’s analysis in Chapter III of his book. Unfortunately, Shelah could not quite prove the stronger results he had hoped for. Still, in [She09, Discussion III.12.40], he claims the following (a proof should appear in [She]):

Claim 1.5. Assume the weak generalized continuum hypothesis⁹ (WGCH). Let K be an AEC such that there is an ω -successful good λ -frame with underlying class K_λ . Then K is categorical in some $\mu > \lambda^{+\omega}$ if and only if K is categorical in all $\mu > \lambda^{+\omega}$.

Modulo this claim, we obtain the consistency of Shelah’s categoricity conjecture from large cardinals. This partially answers [She00, Question 6.14]:

Theorem 1.6. Assume Claim 1.5 and WGCH.

⁷A version of Shelah’s categoricity conjecture already appears as [She90, Open problem D.3(a)] and the statement here appears in [She09, Conjecture N.4.2], see [Gro02] or the introduction to [She09] for history and motivation.

⁸Since (e.g. by the presentation theorem) there are only set many AECs with a given Löwenheim-Skolem number, this is equivalent to requiring categoricity in a single high-enough cardinal.

⁹Namely, $2^\lambda < 2^{\lambda^+}$ for all cardinals λ .

- (1) Shelah's categoricity conjecture holds in fully tame and short AECs with amalgamation.
- (2) If there exists unboundedly many strongly compact cardinals, then Shelah's categoricity conjecture holds.

Proof. Let K be an AEC which is categorical in unboundedly many cardinals.

- (1) Assume K is fully $\text{LS}(K)$ -tame and short and has amalgamation. Pick κ and λ such that $\text{LS}(K) < \kappa = \beth_\kappa < \lambda = \beth_\lambda$ and $\text{cf}(\lambda) \geq \kappa$. By Theorem 1.3, there is an ω -successful good λ -frame on K_λ . By Claim 1.5, K is categorical in all $\mu > \lambda^{+\omega}$.
- (2) Let $\kappa > \text{LS}(K)$ be strongly compact. By [Bon14b], K is fully ($< \kappa$)-tame and short. By the methods of [MS90, Proposition 1.13], $K_{\geq \kappa}$ has amalgamation. Now apply the previous part to $K_{\geq \kappa}$.

□

Remark 1.7. Previous works (e.g. [MS90, She99, GV06c, Bon14b]) all assume categoricity in a successor cardinal, and this was thought to be hard to remove. Here, we do *not* need to assume categoricity in a successor.

Note that [She09, Theorem IV.7.12] is stronger than Theorem 1.6 (since Shelah assumes only Claim 1.5, WGCH, and amalgamation) but we were unable to verify Shelah's proof. Also, the statement contains an error as it contradicts Morley's categoricity theorem.

This paper is organized as follows. In Section 2 we review some of the background. In Sections 3-4, we introduce the framework with which we will study independence. In Sections 5-8, we introduce the definition of a *generator* for an independence relation and show how to use it to build good frames. In Section 9, we use the theory of generators to prove results on the canonicity of coheir and good frames. In Section 10, we use generators to give a general definition of superstability (closely related to those implicit in [GVV, Vasa]). We derive superstability from categoricity and use it to construct good frames. In Section 11, we show how to prove a good frame is ω -successful provided it is induced by coheir. In Sections 12-14, we show how to extend such a frame to a global independence relation. In Section 15, some of the main theorems are established.

This paper was written while working on a Ph.D. thesis under the direction of Rami Grossberg at Carnegie Mellon University and I would

like to thank Professor Grossberg for his guidance and assistance in my research in general and in this work specifically.

I also thank Andrés Villaveces for sending his thoughts on my results and Will Boney for thoroughly reading this paper and giving invaluable feedback.

2. PRELIMINARIES

We review some of the basics of abstract elementary classes and fix some notation. The reader is advised to skim through this section quickly and go back to it as needed. We refer the reader to the preliminaries of [Vasb] for more motivation on some of the definitions below.

2.1. Set theoretic terminology.

Notation 2.1. When we say that \mathcal{F} is an interval of cardinals, we mean that $\mathcal{F} = [\lambda, \theta)$ is the set of cardinals μ such that $\lambda \leq \mu < \theta$. Here, $\lambda \leq \theta$ are (possibly finite) cardinals except we also allow $\theta = \infty$.

We will often use the following function:

Definition 2.2 (Hanf function). For λ an infinite cardinal, define $h(\lambda) := \beth_{(2^\lambda)^+}$.

Note that for λ infinite, $\lambda = \beth_\lambda$ if and only if for all $\mu < \lambda$, $h(\mu) < \lambda$.

Definition 2.3. For κ an infinite cardinal, let κ_r be the least regular cardinal $\geq \kappa$. That is, κ_r is κ^+ if κ is singular and κ otherwise.

2.2. Abstract classes. Recall¹⁰ [Vasb, Definition 2.7] that an abstract class (AC for short) is a pair (K, \leq) , where K is a class of structures of the same (possibly infinitary) language and \leq is an ordering on K extending substructure and respecting isomorphisms. We will use the same notation as in [Vasb]; for example $M < N$ means $M \leq N$ and $M \neq N$.

Definition 2.4. Let K be an abstract class and let R be a binary relation on K . A sequence $\langle M_i : i < \delta \rangle$ of elements of K is *R-increasing* if for all $i < j < \delta$, $M_i R M_j$. When $R = \leq$, we omit it. *Strictly increasing* means $<$ -increasing. $\langle M_i : i < \delta \rangle$ is *continuous* if for all limit $i < \delta$, $M_i = \bigcup_{j < i} M_j$.

¹⁰The definition is due to Rami Grossberg and appears in [Gro].

Notation 2.5. For K an abstract class, \mathcal{F} an interval of cardinals, we write $K_{\mathcal{F}} := \{M \in K \mid \|M\| \in \mathcal{F}\}$. When $\mathcal{F} = \{\lambda\}$, we write K_{λ} for $K_{\{\lambda\}}$. We also use notation like $K_{\geq \lambda}$, $K_{< \lambda}$, etc.

Definition 2.6. An abstract class K is *in* \mathcal{F} if $K_{\mathcal{F}} = K$.

We now recall the definition of an abstract elementary class (AEC) in \mathcal{F} , for \mathcal{F} an interval of cardinal. Localizing to an interval is convenient when dealing with good frames and appears already (for $\mathcal{F} = \{\lambda\}$) in [JS13, Definition 1.0.3.2]. Confusingly, Shelah earlier on called an AEC in λ a λ -AEC (in [She09, Definition II.1.18]).

Definition 2.7. For $\mathcal{F} = [\lambda, \theta)$ an interval of cardinals, we say an abstract class K in \mathcal{F} is an *abstract elementary class* (AEC for short) *in* \mathcal{F} if it satisfies:

- (1) Coherence: If M_0, M_1, M_2 are in K , $M_0 \leq M_2$, $M_1 \leq M_2$, and $|M_0| \subseteq |M_1|$, then $M_0 \leq M_1$.
- (2) $L(K)$ is finitary.
- (3) Tarski-Vaught axioms: If $\langle M_i : i < \delta \rangle$ is an increasing chain in K and $\delta < \theta$, then $M_{\delta} := \bigcup_{i < \delta} M_i$ is such that:
 - (a) $M_{\delta} \in K$.
 - (b) $M_0 \leq M_{\delta}$.
 - (c) If $M_i \leq N$ for all $i < \delta$, then $M_{\delta} \leq N$.
- (4) Löwenheim-Skolem axiom: There exists a cardinal $\lambda \geq |L(K)| + \aleph_0$ such that for any $M \in K$ and any $A \subseteq |M|$, there exists $M_0 \leq M$ containing A with $\|M_0\| \leq |A| + \lambda$. We write $\text{LS}(K)$ (the *Löwenheim-Skolem number of* K) for the least such cardinal.

When $\mathcal{F} = [0, \infty)$, we omit it. We say K is an *AEC in* λ if it is an AEC in $\{\lambda\}$.

Recall that an AEC in \mathcal{F} can be made into an AEC:

Fact 2.8 (Lemma II.1.23 in [She09]). If K is an AEC in $\lambda := \text{LS}(K)$, then there exists a unique AEC K' such that $(K')_{\lambda} = K$ and $\text{LS}(K') = \lambda$. The same holds if K is an AEC in \mathcal{F} , $\mathcal{F} = [\lambda, \theta)$ (apply the previous sentence to K_{λ}).

Notation 2.9. Let K be an AEC in \mathcal{F} with $\mathcal{F} = [\lambda, \theta)$, $\lambda = \text{LS}(K)$. Write K^{up} for the unique AEC K' described by Fact 2.8.

When studying independence, the following definition will be useful:

Definition 2.10. A *coherent abstract class* in \mathcal{F} is an abstract class in \mathcal{F} satisfying the coherence property (see Definition 2.7).

We also define the following weakening of the existence of a Löwenheim-Skolem number:

Definition 2.11. An abstract class K is $(< \lambda)$ -closed if for any $M \in K$ and $A \subseteq |M|$ with $|A| < \lambda$, there exists $M_0 \leq M$ which contains A and has size less than λ . λ -closed means $(< \lambda^+)$ -closed.

Remark 2.12. An AEC K is $(< \lambda)$ -closed in every $\lambda > \text{LS}(K)$.

We will sometimes use the following consequence of Shelah's presentation theorem:

Fact 2.13 (Conclusion I.1.11 in [She09]). Let K be an AEC. If $K_{\geq \lambda} \neq \emptyset$ for every $\lambda < h(\text{LS}(K))$, then K has arbitrarily large models.

As in the preliminaries of [Vasb], we can define a notion of embedding for abstract classes and go on to define amalgamation, joint embedding, no maximal models, Galois types, tameness, and type-shortness (that we will just call shortness).

The following fact tells us that an AEC with amalgamation is a union of AECs with amalgamation and joint embedding (see also the notion of the *diagram* of an AEC [She09, Definition I.2.2]):

Fact 2.14 (Lemma 16.14 in [Bal09]). Let K be an AEC with amalgamation. Then we can write $K = \bigcup_{i \in I} K^i$ where the K^i 's are disjoint AECs with $\text{LS}(K^i) = \text{LS}(K)$ and each K^i has joint embedding and amalgamation.

Using Galois types, a natural notion of saturation can be defined (see [Vasb, Definition 2.22] for more explanation on the definition):

Definition 2.15. Let K be an abstract class and μ be an infinite cardinal.

- (1) A model $M \in K$ is μ -saturated if for all $N \geq M$ and all $A_0 \subseteq |M|$ of size less than μ , any $p \in \text{gS}^{<\mu}(A_0; N)$ is realized inside M . When $\mu = \|M\|$, we omit it.
- (2) We write $K^{\mu\text{-sat}}$ for the class of μ -saturated models of $K_{\geq \mu}$ (ordered by the ordering of K).

Remark 2.16. By [She09, Lemma II.1.14], if K is an AEC with amalgamation and $\mu > \text{LS}(K)$, $M \in K$ is μ -saturated if and only if for all $N \geq M$ and all $A_0 \subseteq |M|$ with $|A_0| < \mu$, any $p \in \text{gS}(A_0; N)$ is realized in M . That is, it is enough to consider types of length 1 in the definition. We will use this fact freely.

Finally, we recall there is a natural notion of stability in this context. This paper's definition follows [Vasb, Definition 2.20].

Definition 2.17 (Stability). Let α be a cardinal, μ be a cardinal. A model $N \in K$ is $(< \alpha)$ -stable in μ if for all $A \subseteq |N|$ of size $\leq \mu$, $|\text{gS}^{<\alpha}(A; N)| \leq \mu$. Here and below, α -stable means $(< (\alpha^+))$ -stable. We say “stable” instead of “1-stable”.

K is $(< \alpha)$ -stable in μ if every $N \in K$ is $(< \alpha)$ -stable in μ . K is $(< \alpha)$ -stable if it is $(< \alpha)$ -stable in unboundedly¹¹ many cardinals.

A corresponding definition of the order property in AECs appears in [She99, Definition 4.3]. For simplicity, we have removed one parameter from the definition.

Definition 2.18. Let α and μ be cardinals and let K be an abstract class. A model $M \in K$ has the α -order property of length μ if there exists $\langle \bar{a}_i : i < \mu \rangle$ inside M with $\ell(\bar{a}_i) = \alpha$ for all $i < \mu$, such that for any $i_0 < j_0 < \mu$ and $i_1 < j_1 < \mu$, $\text{gtp}(\bar{a}_{i_0} \bar{a}_{j_0} / \emptyset; N) \neq \text{gtp}(\bar{a}_{j_1} \bar{a}_{i_1} / \emptyset; N)$.

M has the $(< \alpha)$ -order property of length μ if it has the β -order property of length μ for some $\beta < \alpha$. M has the order property of length μ if it has the α -order property of length μ for some α .

K has the α -order of length μ if some $M \in K$ has it. K has the order property if it has the order property for every length.

For completeness, we also recall the definition of the following variation on the $(< \kappa)$ -order property of length κ that appears in [BG, Definition 4.2] (but is adapted from a previous definition of Shelah, see there for more background):

Definition 2.19. Let K be an AEC. For $\kappa > \text{LS}(K)$, K has the weak κ -order property if there are $\alpha, \beta < \kappa$, $M \in K_{<\kappa}$, $N \geq M$, types $p \neq q \in \text{gS}^{\alpha+\beta}(M)$, and sequences $\langle \bar{a}_i : i < \kappa \rangle$, $\langle \bar{b}_i : i < \kappa \rangle$ from N so that for all $i, j < \kappa$:

- (1) $i \leq j$ implies $\text{gtp}(\bar{a}_i \bar{b}_j / M; N) = p$.
- (2) $i > j$ implies $\text{gtp}(\bar{a}_i \bar{b}_j / M; N) = q$.

The following sums up all the results we will use about stability and the order property:

Fact 2.20. Let K be an AEC.

¹¹Note ([GV06b, Corollary 6.4]) that in a $\text{LS}(K)$ -tame AEC with amalgamation, this is equivalent to stability in *some* cardinal.

- (1) [Vasb, Lemma 4.8] Let $\kappa = \beth_\kappa > \text{LS}(K)$. The following are equivalent:
- (a) K has the weak κ -order property.
 - (b) K has the $(< \kappa)$ -order property of length κ .
 - (c) K has the $(< \kappa)$ -order property.
- (2) [Vasb, Theorem 4.13] Assume K is $(< \kappa)$ -tame and has amalgamation. The following are equivalent:
- (a) K is stable in some $\lambda \geq \kappa + \text{LS}(K)$.
 - (b) There exists $\mu \leq \lambda_0 < h(\kappa + \text{LS}(K))$ such that K is stable in any $\lambda \geq \lambda_0$ with $\lambda = \lambda^{<\mu}$.
 - (c) K does not have the order property.
 - (d) K does not have the $(< \kappa)$ -order property.
- (3) [BKV06, Theorem 4.5] If K is $\text{LS}(K)$ -tame, has amalgamation, and is stable in $\text{LS}(K)$, then it is stable in $\text{LS}(K)^+$.

2.3. Universal and limit extensions.

Definition 2.21. Let K be an abstract class, λ be a cardinal.

- (1) For $M, N \in K$, say $M <_{\text{univ}} N$ (N is *universal over* M) if and only if $M < N$ and whenever we have $M' \geq M$ such that $\|M'\| \leq \|N\|$, then there exists $f : M' \xrightarrow{M} N$. Say $M \leq_{\text{univ}} N$ if and only if $M = N$ or $M <_{\text{univ}} N$.
- (2) For $M, N \in K$, λ a cardinal and $\delta \leq \lambda^+$, say $M <_{\lambda, \delta} N$ (N is (λ, δ) -*limit over* M) if and only if $M \in K_\lambda$, $N \in K_{\lambda+|\delta|}$, $M < N$, and there exists $\langle M_i : i \leq \delta \rangle$ increasing continuous such that $M_0 = M$, $M_i <_{\text{univ}} M_{i+1}$ for all $i < \delta$, and $M_\delta = N$ if $\delta > 0$. Say $M \leq_{\lambda, \delta}$ if $M = N$ or $M <_{\lambda, \delta} N$. We say $N \in K$ is a (λ, δ) -*limit model* if $M <_{\lambda, \delta} N$ for some M . We say N is λ -*limit* if it is (λ, δ) -limit for some limit $\delta < \lambda^+$. When λ is clear from context, we omit it.

Remark 2.22. So for $M, N \in K_\lambda$, $M <_{\lambda, 0} N$ if and only if $M < N$, while $M <_{\lambda, 1}$ if and only if $M <_{\text{univ}} N$.

Remark 2.23. Variations on $<_{\lambda, \delta}$ already appear as [She99, Definition 2.1]. This paper's definition of being universal is different from the usual one (see e.g. [Van06, Definition I.2.1.2]) because we ask only for $\|M'\| \leq \|N\|$ rather than $\|M'\| = \|M\|$.

Fact 2.24. Let K be an AC with amalgamation, λ be an infinite cardinal, and $\delta \leq \lambda^+$. Then:

- (1) $M_0 <_{\text{univ}} M_1 \leq M_2$ implies $M_0 <_{\text{univ}} M_2$.
- (2) $M_0 \leq M_1 <_{\text{univ}} M_2$ implies $M_0 <_{\text{univ}} M_2$.

- (3) If $M_0 \in K_\lambda$, then $M_0 \leq M_1 <_{\lambda, \delta} M_2$ implies $M_0 <_{\lambda, \delta} M_2$.
- (4) If $\delta < \lambda^+$, K is an AEC in $\lambda = \text{LS}(K)$ with no maximal models and stability in λ , then for any $M_0 \in K$ there exists M'_0 such that $M_0 <_{\lambda, \delta} M'_0$.

Proof. All are straightforward, except perhaps the last which is due to Shelah. For proofs and references see [Vasa, Proposition 2.12]. \square

By a routine back and forth argument, we have:

Fact 2.25 (Fact 1.3.6 in [SV99]). Let K be an AEC in $\lambda := \text{LS}(K)$ with amalgamation. Let $\delta \leq \lambda^+$ be a limit ordinal and for $\ell = 1, 2$, let $\langle M_i^\ell : i \leq \delta \rangle$ be increasing continuous with $M_0 := M_0^1 = M_0^2$ and $M_i^\ell <_{\text{univ}} M_{i+1}^\ell$ for all $i < \delta$ (so they witness $M_0^\ell <_{\lambda, \delta} M_\delta^\ell$).

Then there exists $f : M_\delta^1 \cong_{M_0} M_\delta^2$ such that for all $i < \delta$, there exists $j < \delta$ such that $f[M_i^1] \leq M_j^2$ and $f^{-1}[M_j^2] \leq M_i^1$.

Remark 2.26. Uniqueness of limit models that are *not* of the same length (i.e. the statement $M_0 <_{\lambda, \delta} M_1$, $M_0 <_{\lambda, \delta'} M_2$ implies $M_1 \cong_{M_0} M_2$ for any limit $\delta, \delta' < \lambda^+$) has been argued to be an important dividing line, akin to superstability in the first-order theory. See for example [SV99, Van06, Van13, GVV]. It is known to follow from the existence of a good λ -frame (see [She09, Lemma II.4.8], or [Bon14a, Theorem 9.2] for a detailed proof).

We couldn't find a proof of the next result in the literature, so we included one here.

Lemma 2.27. Let K be an AEC with amalgamation. Let δ be a (not necessarily limit) ordinal and assume $(M_i)_{i \leq \delta}$ is increasing continuous with $M_i <_{\text{univ}} M_{i+1}$ for all $i < \delta$. Then $M_i <_{\text{univ}} M_\delta$ for all $i < \delta$.

Proof. By induction on δ . If $\delta = 0$, there is nothing to do. If $\delta = \alpha + 1$ is a successor, let $i < \delta$. We know $M_i \leq M_\alpha$. By hypothesis, $M_\alpha <_{\text{univ}} M_\delta$. By Fact 2.24, $M_i <_{\text{univ}} M_\delta$. Assume now δ is limit. In that case it is enough to show $M_0 <_{\text{univ}} M_\delta$. By the induction hypothesis, we can further assume that $\delta = \text{cf}(\delta)$. Let $N \geq M_0$ be given such that $\mu := \|N\| \leq \|M_\delta\|$, and N, M_δ are inside a common model \widehat{N} . If $\mu < \|M_\delta\|$, then there exists $i < \delta$ such that $\mu \leq \|M_i\|$, and we can use the induction hypothesis, so assume $\mu = \|M_\delta\|$. We can further assume $\mu > \|M_0\|$, for otherwise N directly embeds into M_1 over M_0 . The M_i s show that $\gamma := \text{cf}(\mu) \leq \delta$. Let $\langle N_i : i \leq \gamma \rangle$ be increasing continuous such that for all $i < \gamma$.

- (1) $N_0 = M_0$.
- (2) $N_\gamma = N$.
- (3) $\|N_i\| < \mu$.

This exists since $\gamma = \text{cf}(\mu)$.

Build $\langle f_i : i \leq \gamma \rangle$, increasing continuous such that for all $i < \gamma$, $f_i : N_i \xrightarrow{M_0} M_{k_i}$ for some $k_i < \delta$. This is enough, since then f_γ will be the desired embedding. This is possible: For $i = 0$, take $f_0 := \text{id}_{M_0}$. At limits, take unions: since δ is regular and $\gamma \leq \delta$, $k_j < \delta$ for all $j < i < \gamma$ implies $k_i := \sup_{j < i} k_j < \delta$.

Now given $i = j + 1$, first pick $k = k_j < \delta$ such that $f_j[N_j] \leq M_k$. Such a k exists by the induction hypothesis. Find $k' > k$ such that $\|N_i\| \leq \|M_{k'}\|$. This exists since $\|N_i\| < \mu = \|M_\delta\|$. Now by the induction hypothesis, $M_k <_{\text{univ}} M_{k'}$, so by Fact 2.24, $f_j[N_j] <_{\text{univ}} M_{k'}$. Hence by some renaming, we can extend f_i as desired. \square

Remark 2.28. (K, \leq_{univ}) is in general not an AEC as it may fail the Löwenheim-Skolem axiom, the coherence axiom, and (3c) in the Tarski-Vaught axioms of Definition 2.7.

3. INDEPENDENCE RELATIONS

Since this section mostly lists definitions, the reader already familiar with independence (e.g. in the first-order context) may want to skip it and refer to it as needed. We would like a general framework in which to study independence. One such framework is Shelah's good λ -frames [She09, Section II.6]. Another is given by the definition of independence relation in [BGKV, Definition 3.1] (itself adapted from [BG, Definition 3.3]). Both definitions describe a relation “ p does not fork over M ” for p a Galois type over N and $M \leq N$.

In [BGKV], it is also shown how to “close” such a relation to obtain a relation “ p does not fork over M ” when p is a type over an arbitrary set. We find that starting with such a relation makes the statement of symmetry transparent, and hence makes many proofs easier. Perhaps even more importantly, we can be very precise¹² when dealing with

¹²Assume for example that \mathfrak{s} is a good-frame on a class of saturated models of an AEC K . Let $\langle M_i : i < \delta \rangle$ be an increasing chain of saturated models. Let $M_\delta := \bigcup_{i < \delta} M_i$ and let $p \in \text{gS}(M_\delta)$. We would like to say that there is $i < \delta$ such that p does not fork over M_i but we may not know that M_δ is saturated, so maybe forking is not even defined for types over M_δ . However if the forking relation were defined for types over sets, there would be no problem.

chain local character properties (see Definition 3.14). We also give a more general definition than [BGKV], as we do not assume that everything happens in a big homogeneous monster model, and we allow working inside coherent abstract classes (recall Definition 2.10) rather than only abstract elementary classes. The later feature is convenient when working with classes of saturated models.

Because we quote extensively from [She09], which deals with frames, and also because it is sometimes convenient to “forget” the extension of the relation to arbitrary sets, we will still define frames and recall their relationship with independence relations over sets.

3.1. Frames. Shelah’s definition of a pre-frame appears in [She09, Definition III.0.2.1] and is meant to axiomatize the bare minimum of properties a relation must satisfy in order to be a meaningful independence notions.

We make several changes: we do not mention basic types (we have no use for them), so in Shelah’s terminology our pre-frames will be *type-full*. In fact, it is notationally convenient for us to define our frame on every type, not just the nonalgebraic ones. The *disjointness* property (see Definition 3.10) tells us that the frame behaves trivially on the algebraic types. We do not require it (as it is not required in [BGKV, Definition 3.1]) but it will hold of all frames we consider.

We require that the class on which the independence relation operates has amalgamation¹³, and we do not require that the base monotonicity property holds (this is to preserve the symmetry between right and left properties in the definition. All the frames we consider will have base monotonicity). Finally, we allow the size of the models to lie in an interval rather than just be restricted to a single cardinal as Shelah does. We also parametrize on the length of the types. This allows more flexibility and was already the approach favored in [Vas, BVb].

Definition 3.1. Let $\mathcal{F} = [\lambda, \theta)$ be an interval of cardinals with $\aleph_0 \leq \lambda < \theta$, $\alpha \leq \theta$ be a cardinal or ∞ .

A *type-full pre- $(< \alpha, \mathcal{F})$ -frame* is a pair $\mathfrak{s} = (K, \perp)$, where:

- (1) K is a coherent abstract class in \mathcal{F} (see Definition 2.10) with amalgamation.
- (2) \perp is a relation on quadruples of the form (M_0, A, M, N) , where $M_0 \leq M \leq N$ are all in K , $A \subseteq |N|$ is such that $|A \setminus M_0| < \alpha$.

¹³This is required in Shelah’s definition of good frames, but not in his definition of pre-frames.

We write $\perp(M_0, A, M, N)$ or $A \underset{M_0}{\overset{N}{\perp}} M$ instead of $(M_0, A, M, N) \in \perp$.

(3) The following properties hold:

(a) Invariance: If $f : N \cong N'$ and $A \underset{M_0}{\overset{N}{\perp}} M$, then $f[A] \underset{f[M_0]}{\overset{N'}{\perp}} f[M]$.

(b) Monotonicity: Assume $A \underset{M_0}{\overset{N}{\perp}} M$. Then:

(i) Ambient monotonicity: If $N' \geq N$, then $A \underset{M_0}{\overset{N'}{\perp}} M$. If

$M \leq N_0 \leq N$ and $A \subseteq |N_0|$, then $A \underset{M_0}{\overset{N_0}{\perp}} M$.

(ii) Left and right monotonicity: If $A_0 \subseteq A$, $M_0 \leq M' \leq M$, then $A_0 \underset{M_0}{\overset{N}{\perp}} M'$.

(c) Left normality: If $A \underset{M_0}{\overset{N}{\perp}} M$, then¹⁴ $AM_0 \underset{M_0}{\overset{N}{\perp}} M$.

When α or \mathcal{F} are clear from context or irrelevant, we omit them and just say that \mathfrak{s} is a pre-frame (or just a frame). We may omit the “type-full”. A $(\leq \alpha)$ -frame is just a $(< (\alpha + 1))$ -frame. We might omit α when $\alpha = 2$ (i.e. \mathfrak{s} is a (≤ 1) -frame) and we might talk of a λ -frame or a $(\geq \lambda)$ -frame instead of a $\{\lambda\}$ -frame or a $[\lambda, \infty)$ -frame.

Notation 3.2. For $\mathfrak{s} = (K, \perp)$ a pre- $(< \alpha, \mathcal{F})$ -frame with $\mathcal{F} = [\lambda, \theta]$, write¹⁵ $K_{\mathfrak{s}} := K$, $\underset{\mathfrak{s}}{\perp} := \perp$, $\alpha_{\mathfrak{s}} := \alpha$, $\mathcal{F}_{\mathfrak{s}} = \mathcal{F}$, $\lambda_{\mathfrak{s}} := \lambda$, $\theta_{\mathfrak{s}} := \theta$.

Notation 3.3. For $\mathfrak{s} = (K, \perp)$ a pre-frame, we write $\perp(M_0, \bar{a}, M, N)$ or $\bar{a} \underset{M_0}{\overset{N}{\perp}} M$ for $\text{ran}(\bar{a}) \underset{M_0}{\overset{N}{\perp}} M$ (similarly when other parameters are sequences). When $p \in \text{gS}^{<\infty}(M)$, we say p *does not \mathfrak{s} -fork over M_0* (or just *does not fork over M_0* if \mathfrak{s} is clear from context) if $\bar{a} \underset{M_0}{\overset{N}{\perp}} M$ whenever $p = \text{gtp}(\bar{a}/M; N)$ (using monotonicity and invariance, it is easy to check that this does not depend on the choice of representatives).

Remark 3.4. In the definition of a pre-frame given in [BVb, Definition 3.1], the left hand side of the relation \perp is a sequence, not just a set.

¹⁴For sets A and B , we sometimes write AB instead of $A \cup B$.

¹⁵Pedantically, α , \mathcal{F} , and θ should be part of the data of the frame but we ignore this detail.

Here, we simply assume outright that the relation is defined so that order does not matter.

Remark 3.5. We can go back and forth from this paper's definition of pre-frame to Shelah's. We sketch how. From a pre-frame \mathfrak{s} in our sense (with $K_{\mathfrak{s}}$ an AEC), we can let $S^{\text{bs}}(M)$ be the set of nonalgebraic $p \in \text{gS}(M)$ that do not \mathfrak{s} -fork over M . Then restricting \perp to the basic types we obtain (assuming that \mathfrak{s} has base monotonicity, see Definition 3.10) a pre-frame in Shelah's sense. From a pre-frame $(K, \perp, S^{\text{bs}})$ in Shelah's sense (where K has amalgamation), we can extend \perp by specifying that algebraic and basic types do not fork over their domains. We then get a pre-frame \mathfrak{s} in our sense with base monotonicity and disjointness.

3.2. Independence relations. We now give a definition for an independence notion that also takes sets on the right hand side.

Definition 3.6 (Independence relation). Let $\mathcal{F} = [\lambda, \theta)$ be an interval of cardinals with $\aleph_0 \leq \lambda < \theta$, $\alpha, \beta \leq \theta$ be cardinals or ∞ . A $(< \alpha, \mathcal{F}, < \beta)$ -independence relation is a pair $\mathfrak{i} = (K, \perp)$, where:

- (1) K is a coherent abstract class in \mathcal{F} with amalgamation.
- (2) \perp is a relation on quadruples of the form (M, A, B, N) , where $M \leq N$ are all in K , $A \subseteq |N|$ is such that $|A \setminus M| < \alpha$ and $B \subseteq |N|$ is such that $|B \setminus M| < \beta$. We write $\perp(M, A, B, N)$ or $A \underset{M}{\overset{N}{\perp}} B$ instead of $(M, A, B, N) \in \perp$.
- (3) The following properties hold:

(a) Invariance: If $f : N \cong N'$ and $A \underset{M}{\overset{N}{\perp}} B$, then $f[A] \underset{f[M]}{\overset{N'}{\perp}} f[B]$.

(b) Monotonicity: Assume $A \underset{M}{\overset{N}{\perp}} B$. Then:

(i) Ambient monotonicity: If $N' \geq N$, then $A \underset{M}{\overset{N'}{\perp}} B$. If

$M \leq N_0 \leq N$ and $A \cup B \subseteq |N_0|$, then $A \underset{M}{\overset{N_0}{\perp}} B$.

(ii) Left and right monotonicity: If $A_0 \subseteq A$, $B_0 \subseteq B$, then $A_0 \underset{M}{\overset{N}{\perp}} B_0$.

(c) Left and right normality: If $A \underset{M}{\overset{N}{\perp}} B$, then $AM \underset{M}{\overset{N}{\perp}} BM$.

When $\beta = \theta$, we omit it. More generally, when α, β are clear from context or irrelevant, we omit them and just say that \mathbf{i} is an independence relation.

We adopt the same notational conventions as for pre-frames.

Remark 3.7. It seems that in every case of interest $\beta = \theta$. We did not make it part of the definition to avoid breaking the symmetry between α and β . Note also that the case $\alpha > \lambda$ is of particular interest in Section 14.

Before listing the properties independence relations and frames could have, we discuss how to go from one to the other. The cl operation is called the *minimal closure* in [BGKV, Definition 3.4].

Definition 3.8.

- (1) Given a pre-frame $\mathfrak{s} := (K, \perp)$, let $\text{cl}(\mathfrak{s}) := (K, \perp^{\text{cl}})$, where $\perp^{\text{cl}}(M, A, B, N)$ if and only if $M \leq N$, $|B| < \theta_{\mathfrak{s}}$, and there exists $N' \geq N$, $M' \geq M$ containing B such that $\perp(M, A, M', N')$.
- (2) Given a $(< \alpha, \mathcal{F})$ -independence relation $\mathbf{i} = (K, \perp)$ let $\text{pre}(\mathbf{i}) := (K, \perp^{\text{pre}})$, where $\perp^{\text{pre}}(M, A, M', N)$ if and only if $M \leq M' \leq N$ and $\perp(M, A, M', N)$.

Remark 3.9.

- (1) If \mathbf{i} is a $(< \alpha, \mathcal{F})$ -independence relation, then $\text{pre}(\mathbf{i})$ is a pre- $(< \alpha, \mathcal{F})$ -frame.
- (2) If \mathfrak{s} is a pre- $(< \alpha, \mathcal{F})$ -frame, then $\text{cl}(\mathfrak{s})$ is a $(< \alpha, \mathcal{F})$ -independence relation and $\text{pre}(\text{cl}(\mathfrak{s})) = \mathfrak{s}$.

Other properties of cl and pre are given by Proposition 4.1.

Next, we give a long list of properties that an independence relation may or may not have. Most are classical and already appear for example in [BGKV]. We give them here again both for the convenience of the reader and because their definition is sometimes slightly modified compared to [BGKV] (for example, symmetry there is called right full symmetry here, and some properties like uniqueness and extensions are complicated by the fact we do not work in a monster model). They will be used throughout this paper (for example, Section 4 discusses implications between the properties).

Definition 3.10 (Properties of independence relations). Let $\mathbf{i} := (K, \perp)$ be a $(< \alpha, \mathcal{F}, < \beta)$ -independence relation.

- (1) \mathfrak{i} has *disjointness* if $A \underset{M}{\downarrow}^N B$ implies $A \cap B \subseteq |M|$.
- (2) \mathfrak{i} has *symmetry* if $A \underset{M}{\downarrow}^N B$ implies that for all¹⁶ $B_0 \subseteq B$ of size less than α and all $A_0 \subseteq A$ of size less than β , $B_0 \underset{M}{\downarrow}^N A_0$.
- (3) \mathfrak{i} has *right full symmetry* if $A \underset{M}{\downarrow}^N B$ implies that for all $B_0 \subseteq B$ of size less than α and all $A_0 \subseteq A$ of size less than β , there exists $N' \geq N$, $M' \geq M$ containing A_0 such that $B_0 \underset{M}{\downarrow}^{N'} M'$.
- (4) \mathfrak{i} has *right base monotonicity* if $A \underset{M}{\downarrow}^N B$ and $M \leq M' \leq N$, $|M'| \subseteq B \cup |M|$ implies $A \underset{M'}{\downarrow}^N B$.
- (5) \mathfrak{i} has *right existence* if $A \underset{M}{\downarrow}^N M$ for any $A \subseteq |N|$ with $|A| < \alpha$.
- (6) \mathfrak{i} has *right uniqueness* if whenever $M_0 \leq M \leq N_\ell$, $\ell = 1, 2$, $|M_0| \subseteq B \subseteq |M|$, $q_\ell \in \text{gS}^{<\alpha}(B; N_\ell)$, $q_1 \restriction M_0 = q_2 \restriction M_0$, and q_ℓ does not fork over M_0 , then $q_1 = q_2$.
- (7) \mathfrak{i} has *right extension* if whenever $p \in \text{gS}^{<\alpha}(MB; N)$ does not fork over M and $B \subseteq C \subseteq |N|$ with $|C| < \beta$, there exists $N' \geq N$ and $q \in \text{gS}^{<\alpha}(MC; N')$ extending p such that q does not fork over M .
- (8) \mathfrak{i} has *right independent amalgamation* if $\alpha > \lambda$, $\beta = \theta$, and¹⁷ whenever $M_0 \leq M_\ell$ are in K , $\ell = 1, 2$, there exists $N \in K$ and $f_\ell : M_\ell \xrightarrow{M_0} N$ such that $f_1[M_1] \underset{M_0}{\downarrow}^N f_2[M_2]$.
- (9) \mathfrak{i} has the *right ($< \kappa$)-model-witness property* if whenever $M \leq M' \leq N$, $||M'| \setminus |M|| < \beta$, $A \subseteq |N|$, and $A \underset{M}{\downarrow}^N B_0$ for all $B_0 \subseteq |M'|$ of size less than κ , then $A \underset{M}{\downarrow}^N M'$. \mathfrak{i} has the *right ($< \kappa$)-witness property* if this is true when M' is allowed to be an arbitrary set. The λ -[model-]witness property is the $(< \lambda^+)$ -[model-]witness property.

¹⁶Why not just take $B_0 = B$? Because the definition of \downarrow requires that the left hand side has size less than α . Similarly for right full symmetry.

¹⁷Note that even though the next condition is symmetric, the condition on α and β make the left version of the property different from the right.

- (10) \mathbf{i} has *right transitivity* if whenever $M_0 \leq M_1 \leq N$, $A \underset{M_0}{\overset{N}{\downarrow}} M_1$ and $A \underset{M_1}{\overset{N}{\downarrow}} B$ implies $A \underset{M_0}{\overset{N}{\downarrow}} B$. *Strong right transitivity* is the same property when we do not require $M_0 \leq M_1$.
- (11) \mathbf{i} has *right full model-continuity* if K is an AEC in \mathcal{F} , $\alpha > \lambda$, $\beta = \theta$, and whenever $\langle M_i^\ell : i \leq \delta \rangle$ is increasing continuous with δ limit, $\ell \leq 3$, for all $i < \delta$, $M_i^0 \leq M_i^\ell \leq M_i^3$, $\ell = 1, 2$, $\|M_\delta^1\| < \alpha$, and $M_i^1 \underset{M_i^0}{\overset{M_i^3}{\downarrow}} M_i^2$ for all $i < \delta$, then $M_\delta^1 \underset{M_\delta^0}{\overset{M_\delta^3}{\downarrow}} M_\delta^2$.
- (12) *Weak chain local character* is a technical property used to generate weakly good independence relations, see Definition 6.6.

Whenever this makes sense, we similarly define the same properties for pre-frames.

Note that we have defined the right version of the asymmetric properties. One can define a left version by looking at the *dual independence relation*.

Definition 3.11. Let $\mathbf{i} := (K, \downarrow)$ be a $(< \alpha, \mathcal{F}, < \beta)$ -independence relation. Define the *dual independence relation* $\mathbf{i}^d := (K, \underset{d}{\downarrow})$ by $\underset{d}{\downarrow}(M, A, B, N)$ if and only if $\downarrow(M, B, A, N)$.

Remark 3.12.

- (1) If \mathbf{i} is a $(< \alpha, \mathcal{F}, < \beta)$ -independence relation, then \mathbf{i}^d is a $(< \beta, \mathcal{F}, < \alpha)$ -independence relation and $(\mathbf{i}^d)^d = \mathbf{i}$.
- (2) Let \mathbf{i} be a $(< \alpha, \mathcal{F}, < \alpha)$ -independence relation. Then \mathbf{i} has symmetry if and only if $\mathbf{i} = \mathbf{i}^d$.

Definition 3.13. For P a property, we will say \mathbf{i} has *left P* if \mathbf{i}^d has right P . When we omit left or right, we mean the right version of the property.

Definition 3.14 (Locality cardinals). Let $\mathbf{i} = (K, \downarrow)$ be a $(< \alpha, \mathcal{F})$ -independence relation, $\mathcal{F} = [\lambda, \theta)$. Let $\alpha_0 < \alpha$.

- (1) Let $\bar{\kappa}_{\alpha_0}(\mathbf{i})$ be the minimal cardinal $\mu \geq |\alpha_0|^+ + \lambda^+$ such that for any $M \leq N$ in K , any $A \subseteq |N|$ with $|A| \leq \alpha_0$, there exists $M_0 \leq M$ in $K_{<\mu}$ with $A \underset{M_0}{\overset{N}{\downarrow}} M$. When μ does not exist, we set $\bar{\kappa}_{\alpha_0}(\mathbf{i}) = \infty$.

- (2) For R a binary relation on K , Let $\kappa_{\alpha_0}(\mathbf{i}, R)$ be the minimal cardinal $\mu \geq |\alpha_0|^+ + \aleph_0$ such that for any regular $\delta \geq \mu$, any R -increasing (recall Definition 2.4) $\langle M_i : i < \delta \rangle$ in K , any N extending all the M_i 's, and any $A \subseteq |N|$ of size $\leq \alpha_0$, there exists $i < \delta$ such that $A \overset{N}{\underset{M_i}{\downarrow}} M_\delta$. Here, we have set¹⁸ $M_\delta := \bigcup_{i < \delta} M_i$. When $R = \leq$, we omit it. When μ does not exist or $\mu_r \geq \theta$, we set $\kappa_{\alpha_0}(\mathbf{i}) = \infty$.

When K is clear from context, we may write $\bar{\kappa}_{\alpha_0}(\downarrow)$. For $\alpha_0 \leq \alpha$, we also let $\bar{\kappa}_{<\alpha_0}(\mathbf{i}) := \sup_{\alpha'_0 < \alpha_0} \bar{\kappa}_{\alpha'_0}(\mathbf{i})$. Similarly define $\kappa_{<\alpha_0}$.

We similarly define $\bar{\kappa}_{\alpha_0}(\mathfrak{s})$ and $\kappa_{\alpha_0}(\mathfrak{s})$ for \mathfrak{s} a pre-frame (in the definition of $\kappa_{\alpha_0}(\mathfrak{s})$, we require in addition that M_δ be a member of K).

We will use the following notation to restrict independence relations to smaller domains:

Notation 3.15. Let \mathbf{i} be a $(< \alpha, \mathcal{F}, < \beta)$ -independence relation.

- (1) For $\alpha_0 \leq \alpha$, $\beta_0 \leq \beta$, let $\mathbf{i}^{<\alpha_0, <\beta_0}$ denotes the $(< \alpha_0, \mathcal{F}, < \beta_0)$ -independence relation obtained by restricting the types to have length less than α_0 and the right hand side to have size less than β_0 (in the natural way). When $\beta_0 = \beta$, we omit it.
- (2) For K' a coherent sub-AC of $K_{\mathbf{i}}$, let $\mathbf{i} \upharpoonright K'$ be the $(< \alpha, \mathcal{F}, < \beta)$ -independence relation obtained by restricting the underlying class to K' . When \mathbf{i} is a $(< \alpha, \mathcal{F})$ -independence relation and $\mathcal{F}_0 \subseteq \mathcal{F}$ is an interval of cardinals, $\mathcal{F}_0 = [\lambda_{\mathbf{i}}, \theta_0)$, we let $\mathbf{i}_{\mathcal{F}_0} := \mathbf{i}^{<\min(\alpha, \theta_0)} \upharpoonright (K_{\mathbf{i}})_{\mathcal{F}_0}$ be the restriction of \mathbf{i} to models of size in \mathcal{F}_0 and types of appropriate length.

We end this section with two examples of independence relations. The first is coheir. In first-order logic, coheir was first defined in [LP79]¹⁹. A definition of coheir for classes of models of an $L_{\kappa, \omega}$ sentence appears in [MS90] and was later adapted to general AECs in [BG]. In [Vasb], we gave a more conceptual (but equivalent) definition and improved some of the results of Boney and Grossberg. Here, we use Boney and Grossberg's definition but rely on [Vasb].

¹⁸Recall that K is only a coherent abstract class, so may not be closed under unions of chains of length δ .

¹⁹The equivalence of nonforking with coheir (for stable theories) was already established by Shelah in the early seventies and appears in Section III.4 of [She78], see also [She90, Corollary III.4.10].

Definition 3.16. Let K be an AEC with amalgamation and let $\kappa > \text{LS}(K)$.

Define $\mathbf{i}_{\kappa\text{-ch}}(K) := (K^{\kappa\text{-sat}}, \perp)$ by $\perp(M, A, B, N)$ if and only if $M \leq N$ are in $K^{\kappa\text{-sat}}$, $A \cup B \subseteq |N|$, and for any $\bar{a} \in {}^{<\kappa}A$ and $B_0 \subseteq |M| \cup B$ of size less than κ , there exists $\bar{a}' \in {}^{<\kappa}|M|$ such that $\text{gtp}(\bar{a}/B_0; N) = \text{gtp}(\bar{a}'/B_0; M)$.

Fact 3.17 (Theorem 5.13 in [Vasb]). Let K be an AEC with amalgamation and let $\kappa > \text{LS}(K)$. Let $\mathbf{i} := \mathbf{i}_{\kappa\text{-ch}}(K)$.

- (1) \mathbf{i} is a $(< \infty, [\kappa, \infty))$ -independence relation with disjointness, base monotonicity, left and right existence, left and right $(< \kappa)$ -witness property, and strong left transitivity.
- (2) If K does not have the $(< \kappa)$ -order property of length κ , then:
 - (a) \mathbf{i} has symmetry and strong right transitivity.
 - (b) For all α , $\bar{\kappa}_\alpha(\mathbf{i}) \leq ((\alpha + 2)^{<\kappa_r})^+$.
 - (c) If $M_0 \leq M \leq N_\ell$ for $\ell = 1, 2$, $|M_0| \subseteq B \subseteq |M|$, $q_\ell \in \text{gS}^{<\infty}(B; N_\ell)$, $q_1 \upharpoonright M_0 = q_2 \upharpoonright M_0$, q_ℓ does not \mathbf{i} -fork over M_0 for $\ell = 1, 2$, and K is $(< \kappa)$ -tame and short for $\{q_1, q_2\}$, then $q_1 = q_2$.
 - (d) If K is $(< \kappa)$ -tame and short for types of length less than α , then $\text{pre}(\mathbf{i}^{<\alpha})$ has uniqueness. Moreover²⁰ $\mathbf{i}_{[\kappa, \alpha]}^{<\alpha}$ has uniqueness.

Remark 3.18. The extension property²¹ seems to be more problematic. In [BG], Boney and Grossberg simply assumed it (they also showed that it followed from κ being strongly compact [BG, Theorem 8.2.1]). From superstability-like hypotheses, we will obtain more results on it (see Theorem 10.16, Theorem 15.1, and Theorem 15.6).

We now consider another independence notion: splitting. This was first defined for AECs in [She99, Definition 3.2]. Here we define the negative property (nonsplitting), as it is the one we use the most.

Definition 3.19 (λ -nonsplitting). Let K be a coherent abstract class with amalgamation.

- (1) For λ an infinite cardinal, define $\mathfrak{s}_{\lambda\text{-ns}}(K) := (K, \perp)$ by $\bar{a} \perp M$ if and only if $M_0 \leq M \leq N$, $A \subseteq |N|$, and whenever $M_0 \leq$

²⁰Of course, this is only interesting if $\alpha \leq \kappa$.

²¹A word of caution: In [HL02, Section 4], the authors give Shelah's example of an ω -stable class that does not have extension. However, the extension property they consider is *over all sets*, not only over models.

$N_\ell \leq M$, $N_\ell \in K_{\leq \lambda}$, $\ell = 1, 2$, and $f : N_1 \cong_{M_0} N_2$, then $f(\text{gtp}(\bar{a}/N_1; N)) = \text{gtp}(\bar{a}/N_2; N)$.

- (2) Define $\mathfrak{s}_{\text{ns}}(K)$ to have underlying AEC K and forking relation defined such that $p \in \text{gS}^{<\infty}(M)$ does not $\mathfrak{s}_{\text{ns}}(K)$ -fork over $M_0 \leq M$ if and only if p does not $\mathfrak{s}_{\lambda\text{-ns}}(K)$ -fork over M_0 for all infinite λ .
- (3) Let $\mathfrak{i}_{\lambda\text{-ns}}(K) := \text{cl}(\mathfrak{s}_{\lambda\text{-ns}}(K))$, $\mathfrak{i}_{\text{ns}}(K) := \text{cl}(\mathfrak{s}_{\text{ns}}(K))$.

Fact 3.20. Assume K is a coherent AC in $\mathcal{F} = [\lambda, \theta]$ with amalgamation. Let $\mathfrak{s} := \mathfrak{s}_{\text{ns}}(K)$, $\mathfrak{s}' := \mathfrak{s}_{\lambda\text{-ns}}(K)$.

- (1) \mathfrak{s} and \mathfrak{s}' are pre- $(< \infty, \mathcal{F})$ -frame with base monotonicity, left and right existence. If K is λ -closed, \mathfrak{s}' has the right λ -model-witness property.
- (2) If K is an AEC in \mathcal{F} and is stable in λ , then $\bar{\kappa}_{<\omega}(\mathfrak{s}') = \lambda^+$.
- (3) If \mathfrak{t} is a pre- $(< \infty, \mathcal{F})$ -frame with uniqueness and $K_{\mathfrak{t}} = K$, then $\downarrow_{\mathfrak{t}} \subseteq \downarrow_{\mathfrak{s}}$.
- (4) Always, $\downarrow_{\mathfrak{s}} \subseteq \downarrow_{\mathfrak{s}'}$. Moreover if K is λ -tame for types of length less than α , then $\mathfrak{s}^{<\alpha} = (\mathfrak{s}')^{<\alpha}$.
- (5) Let $M_0 <_{\text{univ}} M \leq N$ with $\|M\| = \|N\|$.
 - (a) Weak uniqueness: If $p_\ell \in \text{gS}^\alpha(N)$, $\ell = 1, 2$, do not \mathfrak{s} -fork over M_0 and $p_1 \upharpoonright M = p_2 \upharpoonright M$, then $p_1 = p_2$.
 - (b) Weak extension: If $p \in \text{gS}^{<\infty}(M)$ does not \mathfrak{s} -fork over M_0 and $f : N \xrightarrow{M_0} M$, then $q := f^{-1}(p)$ is an extension of p to $\text{gS}^{<\infty}(N)$ that does not \mathfrak{s} -fork over M_0 . Moreover q is algebraic if and only if p is algebraic.

Proof.

- (1) Easy.
- (2) By [She99, Claim 3.3.1] (see also [GV06b, Fact 4.6]).
- (3) By [BGKV, Lemma 4.2].
- (4) By [BGKV, Proposition 3.12].
- (5) By [Van06, Theorem I.4.10, Theorem I.4.12] (the moreover part is easy to see from the definition of q).

□

Remark 3.21. Fact 3.20.(3) tells us that any reasonable independence relation will be extended by nonsplitting. In this sense, nonsplitting is a *maximal* candidate for an independence relation²².

²²Moreover, $(< \kappa)$ -coheir is a *minimal* candidate in the following sense: Let us say an independence relation $\mathfrak{i} = (K, \downarrow)$ has the *strong* $(< \kappa)$ -witness property if

4. SOME INDEPENDENCE CALCULUS

We investigate relationships between properties and how to go from a frame to an independence relation. Most of it appears already in [BGKV] and has a much longer history, described there. The following are new: Lemma 4.5 gives a way to get the witness properties from tameness, partially answering [BGKV, question 5.5]. Lemmas 4.8 and 4.7 are technical results used in the last sections.

The following proposition investigates what properties are preserved by the operations cl and pre (recall Definition 3.8). This was done already in [BGKV, Section 5.1], so we cite from there.

Proposition 4.1. Let \mathfrak{s} be a $\text{pre-}(< \alpha, \mathcal{F})$ -frame and let \mathfrak{i} be a $(< \alpha, \mathcal{F})$ -independence relation.

- (1) For P in the list of properties of Definition 3.10, if \mathfrak{i} has P , then $\text{pre}(\mathfrak{i})$ has P .
- (2) For P a property in the following list, \mathfrak{i} has P if (and only if) $\text{pre}(\mathfrak{i})$ has P : existence, independent amalgamation, full model-continuity.
- (3) For P a property in the following list, $\text{cl}(\mathfrak{s})$ has P if (and only if) \mathfrak{s} has P : disjointness, full symmetry, base monotonicity, extension, transitivity.
- (4) If $\text{pre}(\mathfrak{i})$ has extension, then $\text{cl}(\text{pre}(\mathfrak{i})) = \mathfrak{i}$ if and only if \mathfrak{i} has extension.
- (5) The following are equivalent:
 - (a) \mathfrak{s} has full symmetry.
 - (b) $\text{cl}(\mathfrak{s})$ has symmetry.
 - (c) $\text{cl}(\mathfrak{s})$ has full symmetry.
- (6) If $\text{pre}(\mathfrak{i})$ has uniqueness and \mathfrak{i} has extension, then \mathfrak{i} has uniqueness.
- (7) If $\text{pre}(\mathfrak{i})$ has extension and \mathfrak{i} has uniqueness, then \mathfrak{i} has extension.

whenever $A \not\underset{M}{\overset{N}{\forking}} B$, there exists $\bar{a}_0 \in {}^{<\kappa}A$ and $B_0 \subseteq |M| \cup B$ of size less than κ

such that $\text{gtp}(\bar{a}'_0/B_0; N) = \text{gtp}(\bar{a}_0/B_0; N)$ implies $\bar{a}_0 \not\underset{M}{\overset{N}{\forking}} B$. Intuitively, this says that forking is witnessed by a formula (and this could be made precise using the methods of [Vasb]). It is easy to check that $(< \kappa)$ -coheir has this property, and any independence relation with strong $(< \kappa)$ -witness and left existence must extend $(< \kappa)$ -coheir.

- (8) ²³ $\bar{\kappa}_{<\alpha}(\mathbf{i}) = \bar{\kappa}_{<\alpha}(\text{pre}(\mathbf{i}))$.
- (9) $\kappa_{<\alpha}(\text{pre}(\mathbf{i})) \leq \kappa_{<\alpha}(\mathbf{i})$. If $K_{\mathbf{i}}$ is an AEC, then this is an equality.

Proof. All are straightforward. See [BGKV, Lemma 5.3, Lemma 5.4]. \square

To what extent is an independence relation determined by its corresponding frame? There is an easy answer:

Lemma 4.2. Let \mathbf{i} and \mathbf{i}' be independence relations with $\text{pre}(\mathbf{i}) = \text{pre}(\mathbf{i}')$. If \mathbf{i} and \mathbf{i}' both have extension, then $\mathbf{i} = \mathbf{i}'$.

Proof. By Proposition 4.1.(4), $\mathbf{i} = \text{cl}(\text{pre}(\mathbf{i}))$ and $\mathbf{i}' = \text{cl}(\text{pre}(\mathbf{i}')) = \text{cl}(\text{pre}(\mathbf{i})) = \mathbf{i}$. \square

The next proposition gives relationships between the properties. We state most results for frames, but they usually have an analog for independence relations that can be obtained using Proposition 4.1.

Proposition 4.3. Let \mathbf{i} be a $(< \alpha, \mathcal{F})$ -independence relation with base monotonicity. Let \mathfrak{s} be a $\text{pre-}(< \alpha, \mathcal{F})$ -frame with base monotonicity.

- (1) If \mathbf{i} has full symmetry, then it has symmetry. If \mathbf{i} has the $(< \kappa)$ -witness property, then it has the $(< \kappa)$ -model-witness property. If $\mathbf{i}[\mathfrak{s}]$ has strong transitivity, then it has transitivity.
- (2) If \mathfrak{s} has uniqueness and extension, then it has transitivity.
- (3) For $\alpha > \lambda$, if \mathfrak{s} has extension and existence, then \mathfrak{s} has independent amalgamation. Conversely, if \mathfrak{s} has transitivity and independent amalgamation, then \mathfrak{s} has extension and existence. Moreover if \mathfrak{s} has uniqueness and independent amalgamation, then it has transitivity.
- (4) If $\min(\kappa_{<\alpha}(\mathfrak{s}), \bar{\kappa}_{<\alpha}(\mathfrak{s})) < \infty$, then \mathfrak{s} has existence.
- (5) $\kappa_{<\alpha}(\mathfrak{s}) \leq \bar{\kappa}_{<\alpha}(\mathfrak{s})$.
- (6) If $K_{\mathfrak{s}}$ is $\lambda_{\mathfrak{s}}$ -closed, $\bar{\kappa}_{<\alpha}(\mathfrak{s}) = \lambda_{\mathfrak{s}}^+$ and \mathfrak{s} has transitivity, then \mathfrak{s} has the right λ -model-witness property.
- (7) If $K_{\mathfrak{s}}$ does not have the order property (Definition 2.18), any chain in $K_{\mathfrak{s}}$ has an upper bound, $\theta = \infty$, and \mathfrak{s} has uniqueness, existence, and extension, then \mathfrak{s} has full symmetry.

Proof.

- (1) Easy.

²³Note that maybe $\alpha = \infty$. However we can always apply the proposition to $\mathfrak{s}^{<\alpha_0}$ for an appropriate $\alpha_0 \leq \alpha$.

- (2) As in the proof of [She09, Claim II.2.18].
- (3) The first sentence is easy, since independent amalgamation is a particular case of extension and existence. Moreover to show existence it is enough by monotonicity to show it for types of models. The proof of transitivity from uniqueness and independent amalgamation is as in (2).
- (4) By definition of the local character cardinals.
- (5) Let $\delta = \text{cf}(\delta) \geq \bar{\kappa}_{<\alpha}(\mathfrak{s})$ and $\langle M_i : i < \delta \rangle$ be increasing in K , $N \geq M_i$ for all $i < \delta$ and $A \subseteq |N|$ with $|A| < \alpha$. Assume $M_\delta := \bigcup_{i < \delta} M_i$ is in K . By definition of $\bar{\kappa}_{<\alpha}$ there exists $N \leq M_\delta$ of size less than $\bar{\kappa}_{\alpha_0}(\mathfrak{i})$ such that p does not fork over N . Now use regularity of δ to find $i < \delta$ with $N \leq M_i$.
- (6) Let $\lambda := \lambda_{\mathfrak{s}}$, say $\mathfrak{s} = (K, \perp)$. Let $M_0 \leq M \leq N$ and assume $A \underset{M_0}{\overset{N}{\perp}} B$ for all $B \subseteq |M|$ with $|B| \leq \lambda$. By definition of $\bar{\kappa}_{<\alpha}(\mathfrak{s})$, there exists $M'_0 \leq M$ of size λ such that $A \underset{M'_0}{\overset{N}{\perp}} M$. By λ -closure and base monotonicity, we can assume without loss of generality that $M_0 \leq M'_0$. By assumption, $A \underset{M_0}{\overset{N}{\perp}} M'_0$, so by transitivity, $A \underset{M_0}{\overset{N}{\perp}} M$.
- (7) As in [BGKV, Corollary 5.16].

□

Remark 4.4. The precise statement of [BGKV, Corollary 5.16] shows that Proposition 4.3.(7) is local in the sense that to prove symmetry over the base model M , it is enough to require uniqueness and extension over this base model (i.e. any two types that do not fork over M , have the same domain, and are equal over M are equal over their domain, and any type over M can be extended to an arbitrary domain so that it does not fork over M).

Lemma 4.5. Let $\mathfrak{i} = (K, \perp)$ be a $(< \alpha, \mathcal{F})$ -independence relation. If \mathfrak{i} has extension and uniqueness, then:

- (1) If K is $(< \kappa)$ -tame for types of length less than α , then K has the right $(< \kappa)$ -model-witness property.
- (2) If K is $(< \kappa)$ -tame and short for types of length less than $\theta_{\mathfrak{i}}$, then K has the right $(< \kappa)$ -witness property.
- (3) If K is $(< \kappa)$ -tame and short for types of length less than $\kappa + \alpha$ and \mathfrak{i} has symmetry, then K has the left $(< \kappa)$ -witness property.

Proof.

- (1) Let $M \leq M' \leq N$ be in K , $A \subseteq |N|$ have size less than α . Assume $A \underset{M}{\overset{N}{\perp}} B_0$ for all $B_0 \subseteq |M'|$ of size less than κ . We want to show that $A \underset{M}{\overset{N}{\perp}} M'$. Let \bar{a} be an enumeration of A , $p := \text{gtp}(\bar{a}/M; N)$. Note that (taking $B_0 = \emptyset$ above) normality implies p does not fork over M . By extension, let $q \in \text{gS}^{<\alpha}(M')$ be an extension of p that does not fork over M . Using amalgamation and some renaming, we can assume without loss of generality that q is realized in N . Let $p' := \text{gtp}(\bar{a}/M'; N)$. We claim that $p' = q$, which is enough by invariance. By the tameness assumption, it is enough to check that $p' \upharpoonright B_0 = q \upharpoonright B_0$ for all $B_0 \subseteq |M'|$ of size less than κ . Fix such a B_0 . By assumption, $p' \upharpoonright B_0$ does not fork over M . By monotonicity, $q \upharpoonright B_0$ does not fork over M . By uniqueness, $p' \upharpoonright B_0 = q \upharpoonright B_0$, as desired.
- (2) Similar to before, noting that for $M \leq N$, $\text{gtp}(\bar{a}/M\bar{b}; N) = \text{gtp}(\bar{a}'/M\bar{b}; N)$ if and only if $\text{gtp}(\bar{a}\bar{b}/M; N) = \text{gtp}(\bar{a}'\bar{b}/M; N)$.
- (3) Observe that in the proof of the previous part, if the set on the right hand side has size less than κ , it is enough to require $(< \kappa)$ -tameness and shortness for types of length less than $(\alpha + \kappa)$. Now use symmetry.

□

Having a nice independence relation makes the class nice. The results below are folklore:

Proposition 4.6. Let $\mathbf{i} = (K, \perp)$ be a $(< \alpha, \mathcal{F})$ -independence relation with base monotonicity. Assume K is an AEC with $\text{LS}(K) = \lambda_{\mathbf{i}}$.

- (1) If \mathbf{i} has uniqueness, and $\kappa := \bar{\kappa}_{<\alpha}(\mathbf{i}) < \infty$, then K is $(< \kappa)$ -tame for types of length less than α .
- (2) If \mathbf{i} has uniqueness and $\kappa := \bar{\kappa}_{<\alpha}(\mathbf{i}) < \infty$, then K is $(< \alpha)$ -stable in any infinite μ such that $\mu = \mu^{<\kappa}$.
- (3) If \mathbf{i} has uniqueness, $\mu > \text{LS}(K)$, K is $(< \alpha)$ -stable in unboundedly many $\mu_0 < \mu$, and $\text{cf}(\mu) \geq \kappa_{<\alpha}(\mathbf{i})$, then K is $(< \alpha)$ -stable in μ .

Proof.

- (1) As in the proof of [Bon14a, Theorem 3.2].

- (2) Let $\mu = \mu^{<\kappa}$ be infinite. Let $M \in K_{\leq \mu}$, $\langle p_i : i < \mu^+ \rangle$ be elements in $\text{gS}^{<\alpha}(M)$. It is enough to show that for some $i < j$, $p_i = p_j$. For each $i < \lambda^+$, there exists $M_i \leq M$ in $K_{<\kappa}$ such that p_i does not fork over M_i . Since $\mu = \mu^{<\kappa}$, we can assume without loss of generality that $M_i = M_0$ for all $i < \mu^+$. Also, $|\text{gS}^{<\alpha}(M_0)| \leq 2^{<\kappa} \leq \mu^{<\kappa} = \mu$ so there exists $i < j < \lambda^+$ such that $p_i \upharpoonright M_0 = p_j \upharpoonright M_0$. By uniqueness, $p_i = p_j$, as needed.
- (3) As in the proof of [Vasa, Lemma 5.5].

□

The following technical result is also used in the last sections. Roughly, it gives conditions under which we can take the base model given by local character to be contained in both the left and right hand side.

Lemma 4.7. Let $\mathbf{i} = (K, \perp)$ be a $(< \alpha, \mathcal{F})$ -independence relation, $\mathcal{F} = [\lambda, \theta)$, with $\alpha > \lambda$. Assume:

- (1) K is an AEC with $\text{LS}(K) = \lambda$.
- (2) \mathbf{i} has base monotonicity and transitivity.
- (3) μ is a cardinal, $\lambda \leq \mu < \theta$.
- (4) \mathbf{i} has the left $(< \kappa)$ -model-witness property for some regular $\kappa \leq \mu$.
- (5) $\bar{\kappa}_\mu(\mathbf{i}) = \mu^+$.

Let $M^0 \leq M^\ell \leq N$ be in K , $\ell = 1, 2$ and assume $M^1 \overset{N}{\underset{M^0}{\perp}} M^2$. Let $A \subseteq |M^1|$, be such that $|A| \leq \mu$. Then there exists $N^1 \leq M^1$ and $N^0 \leq M^0$ such that:

- (1) $A \subseteq |N^1|$, $A \cap |M^0| \subseteq |N^0|$.
- (2) $N^0 \leq N^1$ are in $K_{\leq \mu}$.
- (3) $N^1 \overset{N}{\underset{N^0}{\perp}} M^2$.

Proof. For $\ell = 0, 1$, we build $\langle N_i^\ell : i \leq \kappa \rangle$ increasing continuous in $K_{\leq \mu}$ such that for all $i < \kappa$ and $\ell = 0, 1$:

- (1) $A \subseteq |N_0^1|$, $A \cap |M^0| \subseteq |N_0^0|$.
- (2) $N_i^\ell \leq M^\ell$.
- (3) $N_i^0 \leq N_i^1$.
- (4) $N_i^1 \overset{N}{\underset{N_{i+1}^0}{\perp}} M^2$.

This is possible. Pick any $N_0^0 \leq M^0$ in $K_{\leq \mu}$ containing $A \cap |M^0|$. Now fix $i < \kappa$ and assume inductively that $\langle N_j^0 : j \leq i \rangle, \langle N_j^1 : j < i \rangle$ have been built. If i is a limit, we take unions. Otherwise, pick any $N_i^1 \leq M^1$ in $K_{\leq \mu}$ that contains A , N_j^1 for all $j < i$ and N_i^0 . Now use right transitivity and $\bar{\kappa}_\mu(\mathbf{i}) = \mu^+$ to find $N_{i+1}^0 \leq M^0$ such that $N_i^1 \underset{N_{i+1}^0}{\downarrow}^N M^2$.

By base monotonicity, we can assume without loss of generality that $N_i^0 \leq N_{i+1}^0$.

This is enough. We claim that $N^\ell := N_\kappa^\ell$ are as required. By coherence, $N^0 \leq N^1$ and since $\kappa \leq \mu$ they are in $K_{\leq \mu}$. Since $A \subseteq |N_0^1|$, $A \subseteq |N^1|$. It remains to see $N^1 \underset{N^0}{\downarrow}^N M^2$. By the left witness property²⁴, it is enough to check it for every $B \subseteq |N^1|$ of size less than κ . Fix such a B . Since κ is regular, there exists $i < \kappa$ such that $B \subseteq |N_i^1|$. By assumption and monotonicity, $B \underset{N_{i+1}^0}{\downarrow}^N M^2$. By base monotonicity,

$B \underset{N_\kappa^0}{\downarrow}^N M^2$, as needed. \square

With a similar proof, we can clarify the relationship between full model continuity and local character. Essentially, the next lemma says that local character for types up to a certain length plus full model-continuity implies local character for all lengths. It will be used in Section 14.

Lemma 4.8. Let $\mathbf{i} = (K, \downarrow)$ be a $(< \theta, \mathcal{F})$ -independence relation, $\mathcal{F} = [\lambda, \theta)$. Assume:

- (1) K is an AEC with $\text{LS}(K) = \lambda$.
- (2) \mathbf{i} has base monotonicity, transitivity, and full model continuity.
- (3) \mathbf{i} has the left $(< \kappa)$ -model-witness property for some regular $\kappa \leq \lambda$.
- (4) For all cardinals $\mu \leq \lambda$, $\bar{\kappa}_\mu(\mathbf{i}) = \lambda^+$.

Then for all cardinals $\mu < \theta$, $\bar{\kappa}_\mu(\mathbf{i}) = \lambda^+ + \mu^+$.

Proof. By induction on μ . If $\mu \leq \lambda$, this holds by hypothesis, so assume $\mu > \lambda$. Let $\delta := \text{cf}(\mu)$.

²⁴Note that we do *not* need to use full model continuity, as we only care about chains of cofinality $\geq \kappa$.

Let $M^0 \leq M^1$ be in K and let $A \subseteq |M^1|$ have size μ . We want to find $M \leq M^0$ such that $A \underset{M}{\overset{N}{\downarrow}} M^0$ and $\|M\| \leq \mu$. Let $\langle A_i : i \leq \delta \rangle$ be increasing continuous such that $A = A_\delta$ and $|A_i| < \mu$ for all $i < \delta$.

For $\ell = 0, 1$, we build $\langle N_i^\ell : i \leq \delta \rangle$ increasing continuous such that for all $i < \delta$ and $\ell = 0, 1$:

- (1) $N_i \in K_{<\mu}$.
- (2) $A_i \subseteq |N_i^1|$, $A_i \cap |M^0| \subseteq |N_i^0|$.
- (3) $N_i^\ell \leq M^\ell$.
- (4) $N_i^0 \leq N_i^1$.
- (5) $N_i^1 \underset{N_{i+1}^0}{\overset{M^1}{\downarrow}} M^0$.

This is possible. By (3) and (4), we have $M^1 \underset{M^0}{\overset{M^1}{\downarrow}} M^0$. Now proceed as in the proof of Lemma 4.7.

This is enough. As in the proof of Lemma 4.7, for any $i < \delta$ of cofinality at least κ we have $N_i^1 \underset{N_i^0}{\overset{M^1}{\downarrow}} M^0$. Thus by full model continuity (applied to the sequences $\langle N_i^\ell : i < \delta, \text{cf}(i) \geq \kappa \rangle$), $N_\delta^1 \underset{N_\delta^0}{\overset{M^1}{\downarrow}} M^0$. Since $A = A_\delta \subseteq |N_\delta^1|$, $M := N_\delta^0$ is as needed. \square

5. SKELETONS

We define what it means for an abstract class K' to be a *skeleton* of an abstract class K . The main examples are classes of saturated models with the usual ordering (or even universal or limit extension). Except perhaps for Lemma 5.7, the results of this section are either easy or well known, we simply put them in the general language of this paper.

We will use skeletons to generalize various statements of chain local character (for example in [GVV, Vasa]) that only ask that if $\langle M_i : i < \delta \rangle$ is an increasing chain with respect to *some restriction of the* ordering of K (usually being universal over) and the M_i s are inside some subclass of K (usually some class of saturated models), then any $p \in \text{gS}(\bigcup_{i < \delta} M_i)$ does not fork over some M_i . Lemma 6.8, is they key upward transfer of that property. Note that Lemma 6.7 shows that one can actually assume that skeletons have a particular form. However the generality is still useful when one wants to prove the local character statement.

Also, it seems that many key subclasses appearing in the theory of AECs are skeletons, see the examples below.

Definition 5.1. For (K, \leq) an abstract class, we say (K', \trianglelefteq) is a sub-AC of K if $K' \subseteq K$, (K', \trianglelefteq) is an AC, and $M \trianglelefteq N$ implies $M \leq N$. We similarly define sub-AEC, etc. When $\trianglelefteq = \leq \upharpoonright K'$, we omit it (or may abuse notation and write (K', \leq)).

Definition 5.2. For (K, \leq) an abstract class, we say a set $S \subseteq K$ is *dense* in (K, \leq) if for any $M \in K$ there exists $M' \in S$ with $M \leq M'$.

Definition 5.3. An abstract class (K', \trianglelefteq) is a *skeleton* of (K, \leq) if:

- (1) (K', \trianglelefteq) is a sub-AC of (K, \leq) .
- (2) K' is dense in (K, \leq) .
- (3) If $\langle M_i : i < \alpha \rangle$ is a \trianglelefteq -increasing chain in K' (α not necessarily limit) and there exists $N \in K'$ such that $M_i < N$ for all $i < \alpha$, then we can choose such an N with $M_i \triangleleft N$ for all $i < \alpha$.

Remark 5.4. The term “skeleton” is inspired from the term “skeletal” in [Vas], although there “skeletal” is applied to frames. The intended philosophical meaning is the same: K' has enough information about K so that for many purposes we can work with K' rather than K .

Remark 5.5. Let (K, \leq) be an abstract class. Assume (K', \trianglelefteq) is a dense sub-AC of (K, \leq) with no maximal models satisfying in addition: If $M_0 \leq M_1 \triangleleft M_2$ are in K' , then $M_0 \triangleleft M_2$. Then (K', \trianglelefteq) is a skeleton of (K, \leq) . This property of the ordering already appears in the definition of an abstract universal ordering in [Vas, Definition 2.13]. In the terminology there, if (K, \leq) is an AEC and \triangleleft is an (invariant) universal ordering on K_λ , then $(K_\lambda, \trianglelefteq)$ is a skeleton of (K_λ, \leq) .

Example 5.6. Let K be an AEC.

- (1) Assume K has no maximal models. Let K' be the class of Ehrenfeucht-Mostowski models of K (for some fixed diagram, see for example [Bal09, Theorem 8.18]). Then K' is dense in K (because for a fixed $M \in K$, the AEC K_M of models in K containing a copy of M has EM models), so (K', \leq) is a skeleton of (K, \leq) .
- (2) Let \mathfrak{s} be a weakly successful good λ -frame (see e.g. Definition 11.4 or [She09, Chapter II]) with $K_{\mathfrak{s}} = K_\lambda$. Then $(K_{\lambda^+}^{\lambda^+ \text{-sat}}, \leq_{\lambda^+}^{\text{NF}})$ (see [JS13, Definition 6.1.4] or [She09, Definition II.7.2], where $\leq_{\lambda^+}^{\text{NF}}$ is called $\leq_{\lambda^+}^*$) is a skeleton of (K_{λ^+}, \leq) (use [She09, Claim II.7.4.1, II.7.7.3]).

- (3) Let $\lambda \geq \text{LS}(K)$. Assume K_λ has amalgamation, no maximal models and is stable in λ . Let K' be dense in K_λ and let $\delta < \lambda^+$. Then $(K', \leq_{\lambda, \delta})$ (recall Definition 2.21) is a skeleton of (K_λ, \leq) (use Fact 2.24 and Remark 5.5).

We will only use Example 5.6.(3). However the above suggests that other types of skeletons are also relevant. For example *solvability*, Shelah's definition of superstability [She09, Definition IV.1.4], uses EM models (see the discussion at the beginning of section 10).

The next lemma is a useful tool to find extensions in the skeleton of an AEC with amalgamation:

Lemma 5.7. Let (K', \trianglelefteq) be a skeleton of (K, \leq) . Assume K is an AEC in $\mathcal{F} := [\lambda, \theta)$ with amalgamation. If $M \leq N$ are in K' , then there exists $N' \in K'$ such that $M \trianglelefteq N'$ and $N \trianglelefteq N'$.

Proof. If N is not maximal (with respect to either of the orderings, it does not matter by definition of a skeleton), then by definition of a skeleton we can find $N' \in K'$ such that $N \triangleleft N'$ and $M \triangleleft N'$, as needed.

Now assume N is maximal. We claim that $M \trianglelefteq N$, so $N' := N$ is as desired. Suppose not. Let $\mu := \|N\|$.

We build $\langle M_i : i < \mu^+ \rangle$ and $\langle f_i : M_i \xrightarrow[M]{\rightarrow} N : i < \mu^+ \rangle$ such that:

- (1) $\langle M_i : i < \mu^+ \rangle$ is a strictly increasing chain in (K', \trianglelefteq) with $M_0 = M$.
- (2) $\langle f_i : i < \mu^+ \rangle$ is a strictly increasing chain of K -embeddings.

This is enough. Let $B_{\mu^+} := \bigcup_{i < \mu^+} |M_i|$ and $f_{\mu^+} := \bigcup_{i < \mu^+} f_i$ (Note that it could be that $\mu^+ = \theta$, so B_{μ^+} is just a set and we do not claim that f_{μ^+} is a K -embedding). Then f_{μ^+} is an injection from B_{μ^+} into $|N|$. This is impossible because $|B_{\mu^+}| \geq \mu^+ > \mu = \|N\|$.

This is possible. Set $M_0 := M$, $f_0 := \text{id}_M$. If $i < \mu^+$ is limit, let $M'_i := \bigcup_{j < i} M_j \in K$. By density, find $M''_i \in K'$ such that $M'_i \leq M''_i$. We have that $M_j < M''_i$ for all $j < i$. By definition of a skeleton, this means we can find $M_i \in K'$ with $M_j \triangleleft M_i$ for all $j < i$. Let $f'_i := \bigcup_{j < i} f_j$. Using amalgamation and the fact that N is maximal, we can extend it to $f_i : M_i \xrightarrow[M]{\rightarrow} N$. If $i = j + 1$ is successor, we consider two cases:

- If M_j is not maximal, let $M_i \in K'$ be a \triangleleft -extension of M_j . Using amalgamation and the fact N is maximal, pick $f_i : M_i \xrightarrow[M]{M} N$ an extension of f_j .
- If M_j is maximal, then by amalgamation and the fact both N and M_j are maximal, we must have $N \cong_M M_j$. However by assumption $M_0 \trianglelefteq M_j$ so $M = M_0 \trianglelefteq N$, a contradiction.

□

Thus we get that many properties of a class transfer to its skeletons.

Proposition 5.8. Let (K, \leq) be an AEC in \mathcal{F} and let (K', \trianglelefteq) be a skeleton of K .

- (1) (K, \leq) has no maximal models if and only if (K', \trianglelefteq) has no maximal models.
- (2) If (K, \leq) has amalgamation, then:
 - (a) (K', \trianglelefteq) has amalgamation.
 - (b) (K, \leq) has joint embedding if and only if (K', \trianglelefteq) has joint embedding.
 - (c) Galois types are the same in (K, \leq) and (K', \trianglelefteq) : For any $N \in K'$, $A \subseteq |N|$, $\bar{b}, \bar{c} \in {}^\alpha |N|$, $\text{gtp}_K(\bar{b}/A; N) = \text{gtp}_K(\bar{c}/A; N)$ if and only if $\text{gtp}_{K'}(\bar{b}/A; N) = \text{gtp}_{K'}(\bar{c}/A; N)$. Here, by gtp_K we denote the Galois type computed in (K, \leq) and by $\text{gtp}_{K'}$ the Galois type computed in (K', \trianglelefteq) .
 - (d) (K, \leq) is α -stable in λ if and only if (K', \trianglelefteq) is α -stable in λ .

Proof.

- (1) Directly from the definition.
- (2) (a) Let $M_0 \trianglelefteq M_\ell$ be in K' , $\ell = 1, 2$. By density, find $N \in K'$ and $f_\ell : M_\ell \xrightarrow[M_0]{M} N$ K -embeddings. By Lemma 5.7, there exists $N_1 \in K'$ such that $N \trianglelefteq N_1$, $f_1[M_1] \trianglelefteq N_1$. By Lemma 5.7 again, there exists $N_2 \in K'$ such that $N_1 \trianglelefteq N_2$, $f_2[M_2] \trianglelefteq N_2$. Thus we also have $f_1[M_1] \trianglelefteq N_2$. It follows that $f_\ell : M_\ell \xrightarrow[M]{M} N_2$ is a \trianglelefteq -embedding.
- (b) If (K', \trianglelefteq) has joint embedding, then by density (K, \leq) has joint embedding. The converse is similar to the proof of amalgamation above.
- (c) Note that by density any Galois type (in K) is realized in an element of K' . Since (K', \trianglelefteq) is a sub-AC of (K, \leq) , equality of the types in K' implies equality in K (this

- doesn't use amalgamation). Conversely, assume $\text{gtp}_K(\bar{b}/A; N) = \text{gtp}_K(\bar{c}/A; N)$. Fix $N' \geq N$ in K and a K -embedding $f : N \xrightarrow{A} N'$ such that $f(\bar{b}) = \bar{c}$. By density, we can assume without loss of generality that $N' \in K'$. By Lemma 5.7, find $N'' \in K'$ such that $N \trianglelefteq N''$, $N' \trianglelefteq N''$. By Lemma 5.7 again, find $N''' \in K'$ such that $f[N] \trianglelefteq N'''$, $N'' \trianglelefteq N'''$. By transitivity, $N \trianglelefteq N'''$ and $f : N \xrightarrow{A} N'''$ witnesses equality of the Galois types in (K', \trianglelefteq) .
- (d) Because Galois types are the same in K and K' .

□

We end with an observation concerning universal extensions that will be used in the proof of Lemma 6.7.

Lemma 5.9. Let K be an AEC in $\lambda := \text{LS}(K)$. Assume K has amalgamation, no maximal models, and is stable in λ . Let (K', \trianglelefteq) be a skeleton of K . For any $M \in K'$, there exists $N \in K'$ such that both $M \triangleleft N$ and $M <_{\text{univ}} N$. Thus $(K', \trianglelefteq \cap \leq_{\text{univ}})$ is a skeleton of K .

Proof. For the last sentence, let $\trianglelefteq' := \trianglelefteq \cap \leq_{\text{univ}}$. Note that if $\langle M_i : i < \alpha \rangle$ is a \trianglelefteq' -increasing chain in K' and $M \in K'$ is such that $M_i < M$ for all $i < \alpha$, then by definition of a skeleton we can take M so that $M_i \triangleleft M$ for all $i < \alpha$. If we know that there exists $N \in K'$ with $M \triangleleft N$ and $M <_{\text{univ}} N$, then for all $i < \alpha$, $M_i \triangleleft N$ by transitivity, and $M_i <_{\text{univ}} N$ by Lemma 2.27.

Now let $M \in K'$. By Fact 2.24, there exists $N \in K$ with $M <_{\text{univ}} N$. By density (note that if $N' \geq N$ is in K , then $M <_{\text{univ}} N'$) we can take $N \in K'$. By Lemma 5.7, there exists $N' \in K'$ such that $M \trianglelefteq N'$ and $N \trianglelefteq N'$. Thus (using Fact 2.24 again) $M <_{\text{univ}} N'$, as desired. □

6. GENERATING AN INDEPENDENCE RELATION

In [She09, Section II.2], Shelah showed how to extend a good λ -frame to a pre- $(\geq \lambda)$ -frame. Later, [Bon14a] (with improvements in [BVb]) gave conditions under which all the properties transferred. Similar ideas are used in [Vasa] to directly build a good frame. In this section we adapt Shelah's definition to this paper's more general setup. It is useful to think of the initial λ -frame as a *generator*²⁵ for a $(\geq \lambda)$ -frame, since in case the frame is not good we usually can only get a nice independence

²⁵In [Vasa], we called a generator a skeletal frame (and in earlier version a poor man's frame) but never defined it precisely.

relation on λ^+ -saturated models. Moreover, it is often useful to work with the independence relation being only defined on a dense sub-AC of the original AECs.

Definition 6.1. (K, \mathbf{i}) is a λ -generator for a $(< \alpha)$ -independence relation if:

- (1) α is a cardinal with $2 \leq \alpha \leq \lambda^+$. λ is an infinite cardinal.
- (2) K is an AEC in $\lambda = \text{LS}(K)$
- (3) $\mathbf{i} = (K', \perp)$, where (K', \leq) is a dense sub-AC (recall Definitions 5.1, 5.2) of²⁶ (K, \leq) .
- (4) \mathbf{i} is a $(< \alpha, \lambda)$ -independence relation.
- (5) K^{up} (recall Definition 2.9) has amalgamation.

Remark 6.2. We could similarly define a λ -generator for a $(< \alpha)$ -independence relation below θ , where we require $\theta \geq \lambda^{++}$ and only $K_{\mathcal{F}}^{\text{up}}$ has amalgamation (so when $\theta = \infty$ we recover the above definition). We will not adopt this approach as we have no use for the extra generality and do not want to complicate the notation further. We could also have required less than “ K is an AEC in λ ” but again we have no use for it.

Definition 6.3. Let (K, \mathbf{i}) be a λ -generator for a $(< \alpha)$ -independence relation. Define $(K, \mathbf{i})^{\text{up}} := (K^{\text{up}}, \perp^{\text{up}})$ by $\perp^{\text{up}}(M, A, B, N)$ if and only if $M \leq N$ are in K^{up} and there exists $M_0 \leq M$ in K' such that for all $B_0 \subseteq B$ with $|B_0| \leq \lambda$ and all $N_0 \leq N$ in K' with $A \cup B_0 \subseteq |N_0|$, $M_0 \leq N_0$, we have $\perp_{\mathbf{i}}(M_0, A, B_0, N_0)$.

When $K = K_{\mathbf{i}}$, we write \mathbf{i}^{up} for $(K, \mathbf{i})^{\text{up}}$.

Remark 6.4. In general, we do not claim that $(K, \mathbf{i})^{\text{up}}$ is even an independence relation (the problem is that given $A \subseteq |N|$ with $N \in K^{\text{up}}$ and $|A| \leq \lambda$, there might not be any $M \in K'$ with $M \leq N$ and $A \subseteq |M|$ so the monotonicity properties can fail). Nevertheless, we will abuse notation and use the restriction operations on it.

Lemma 6.5. Let (K, \mathbf{i}) be a λ -generator for a $(< \alpha)$ -independence relation. Then:

- (1) If $K = K_{\mathbf{i}}$, then $\mathbf{i}^{\text{up}} := (K, \mathbf{i})^{\text{up}}$ is an independence relation.
- (2) $(K, \mathbf{i})^{\text{up}} \upharpoonright (K^{\text{up}})^{\lambda^+ \text{-sat}}$ is an independence relation.

²⁶Why not be more general and require only (K', \leq) to be a skeleton of K ? Because some examples of skeletons do not satisfy the coherence axiom which is required by the definition of an independence relation.

Proof. As in [She09, Claim II.2.11], using density and homogeneity in the second case. \square

The case (1) of Lemma 6.5 has been well studied (at least when $\alpha = 2$): see [She09, Section II.2] and [Bon14a, BVb]. We will further look at it in the last sections. We will focus on case (2) for now. It has been studied (implicitly) in [Vasa] when \mathbf{i} is nonsplitting and satisfies some superstability-like assumptions. We will use the same methods to obtain more general results. The generality will be used, since for example we also care about what happens when \mathbf{i} is coheir.

The following property of a generator will be very useful in the next section. The point is that $\bigcup_{i < \lambda^+} M_i$ below is usually not a member of $K_{\mathbf{i}}$ so forking is not defined on it.

Definition 6.6. Let (K, \mathbf{i}) be a λ -generator for a $(< \alpha)$ -independence relation.

(K, \mathbf{i}) has *weak chain local character* if there exists \trianglelefteq such that $(K_{\mathbf{i}}, \trianglelefteq)$ is a skeleton of K and whenever $\langle M_i : i < \lambda^+ \rangle$ is \trianglelefteq -increasing in $K_{\mathbf{i}}$ and $p \in \text{gS}^{<\alpha}(\bigcup_{i < \lambda^+} M_i)$, there exists $i < \lambda^+$ such that $p \upharpoonright M_{i+1}$ does not fork over M_i .

The following technical lemma shows that local character in a skeleton implies local character in a bigger class with the universal ordering:

Lemma 6.7. Let (K, \mathbf{i}) be a λ -generator for a $(< \alpha)$ -independence relation.

Assume that K has amalgamation, no maximal models, and is stable in λ . Assume \mathbf{i} has base monotonicity and let $K' := K_{\mathbf{i}}$. Let (K'', \trianglelefteq) be a skeleton of (K', \leq) and let $\mathbf{i}' := \mathbf{i} \upharpoonright (K'', \trianglelefteq)$. Then:

- (1) $\kappa_{<\alpha}(\mathbf{i}, \leq_{\text{univ}}) \leq \kappa_{<\alpha}(\mathbf{i}', \trianglelefteq)$.
- (2) If (K, \mathbf{i}') has weak chain local character, then (K, \mathbf{i}) has it and it is witnessed by $<_{\text{univ}}$.

Proof.

- (1) By Lemma 5.9, we can (replacing \trianglelefteq by $\trianglelefteq \cap \leq_{\text{univ}}$) assume without loss of generality that \trianglelefteq is extended by \leq_{univ} . Let $\langle M_i : i < \delta \rangle$ be \leq_{univ} -increasing in K' , $\delta = \text{cf}(\delta) \geq \kappa_{<\alpha}(\mathbf{i}', \trianglelefteq)$, $\delta < \lambda^+$. Without loss of generality, $\langle M_i : i < \delta \rangle$ is $<_{\text{univ}}$ -increasing. Let $M_\delta := \bigcup_{i < \delta} M_i$ and let $p \in \text{gS}^{<\alpha}(M_\delta)$.

By density, pick $M'_0 \in K''$ such that $M_0 <_{\text{univ}} M'_0$. Now build $\langle M'_i : i < \delta \rangle \triangleleft$ -increasing in K'' . Let $M'_\delta := \bigcup_{i < \delta} M'_i$. By Fact

2.25, there exists $f : M'_\delta \cong_{M_0} M_\delta$ such that for every $i < \delta$ there exists $j < \delta$ with $f[M'_i] \leq M_j$, $f^{-1}[M_i] \leq M'_j$. By definition of $\kappa_{<\alpha}(\mathbf{i}', \trianglelefteq)$, there exists $i < \delta$ such that $f^{-1}(p)$ does not \mathbf{i}' -fork over M'_i . Let $j < \delta$ be such that $f[M'_i] \leq M_j$. By invariance, p does not \mathbf{i}' -fork over $f[M_i]$, so does not \mathbf{i} -fork over $f[M_i]$. By base monotonicity, p does not \mathbf{i} -fork over M_j , as desired.

(2) Similar.

□

The last lemma of this section investigates what properties directly transfer up.

Lemma 6.8. Let (K, \mathbf{i}) be a λ -generator for a $(< \alpha)$ -independence relation. Let $\mathbf{i}' := (K, \mathbf{i})^{\text{up}} \upharpoonright (K^{\text{up}})^{\lambda^+-\text{sat}}$.

- (1) If \mathbf{i} has base monotonicity, then \mathbf{i}' has base monotonicity.
- (2) Assume \mathbf{i} has base monotonicity and (K, \mathbf{i}) has weak chain local character. Then:
 - (a) $\bar{\kappa}_{<\alpha}(\mathbf{i}') = \lambda^{++}$.
 - (b) If \trianglelefteq is an ordering such that $(K_{\mathbf{i}}, \trianglelefteq)$ is a skeleton of K , then for any $\alpha_0 < \alpha$, $\kappa_{\alpha_0}(\mathbf{i}') \leq \kappa_{\alpha_0}(\mathbf{i}, \trianglelefteq)$.

Proof.

- (1) As in [She09, Claim II.2.11]
- (2) This is a generalization of the proof of [Vasa, Lemma 4.8] but we have to say slightly more so we give the details. Let \trianglelefteq^0 be an ordering witnessing weak chain local character. We first prove (2b). Fix $\alpha_0 < \alpha$, and assume $\kappa_{\alpha_0}(\mathbf{i}, \trianglelefteq) < \infty$. Then by definition $\kappa_{\alpha_0}(\mathbf{i}, \trianglelefteq) \leq \lambda$. Let $\delta = \text{cf}(\delta) \geq \kappa_{\alpha_0}(\mathbf{i}, \trianglelefteq)$.

Let $\langle M_i : i < \delta \rangle$ be increasing in $K^{\lambda^+-\text{sat}}$ and write $M_\delta := \bigcup_{i < \delta} M_i$ (note that we do not claim $M_\delta \in K^{\lambda^+-\text{sat}}$. However, $M_\delta \in K_{\geq \lambda}$). Let $p \in \text{gS}^{\alpha_0}(M_\delta)$. We want to find $i < \delta$ such that p does not fork over M_i . There are two cases:

- Case 1: $\delta < \lambda^+$:

We imitate the proof of [She09, Claim II.2.11.5]. Assume the conclusion fails. Build $\langle N_i : i < \delta \rangle$ \trianglelefteq -increasing in K' , $\langle N'_i : i < \delta \rangle$ \leq -increasing in K' such that for all $i < \delta$:

- (a) $N_i \leq M_i$.
- (b) $N_i \leq N'_i \leq M_\delta$.
- (c) $p \upharpoonright N'_i$ \mathbf{i} -forks over N_i .
- (d) $\bigcup_{j < i} (|N'_j| \cap |M_j|) \subseteq |N_i|$.

This is possible. Assume N_j and N'_j have been constructed for $j < i$. Choose $N_i \leq M_i$ satisfying (2d) so that $N_j \leq N_i$ for all $j < i$ (This is possible: use that M_i is λ^+ -saturated and that in skeletons of AECs, chains have upper bounds). By assumption, p i' -forks over M_i , and so by definition of forking there exists $N'_i \leq M_\delta$ in K' such that $p \restriction N'_i$ forks over N_i . By monotonicity, we can of course assume $N'_i \geq N_i$, $N'_i \geq N'_j$ for all $j < i$.

This is enough. Let $N_\delta := \bigcup_{i < \delta} N_i$, $N'_\delta := \bigcup_{i < \delta} N'_i$. By local character for i , there is $i < \delta$ such that $p \restriction N_\delta$ does not fork over N_i . By (2b) and (2d), $N'_\delta \leq N_\delta$. Thus by monotonicity $p \restriction N'_i$ does not i -fork over N_i , contradicting (2c).

- Case 2: $\delta \geq \lambda^+$: Assume the conclusion fails. As in the previous case (in fact it is easier), we can build $\langle N_i : i < \lambda^+ \rangle \leq^0$ -increasing in K' such that $N_i \leq M_\delta$ and $p \restriction N_{i+1}$ i -forks over N_i . Since i has weak chain local character, there exists $i < \lambda^+$ such that $p \restriction N_{i+1}$ does not i -fork over N_i , contradiction.

For (2a), assume not: then there exists $M \in K^{\lambda^+ \text{-sat}}$ and $p \in \text{gS}^{<\alpha}(M)$ such that for all $M_0 \leq M$ in $K^{\lambda^+ \text{-sat}}$, p i' -forks over M_0 . By stability, for any $A \subseteq |M|$ with $|A| \leq \lambda$, there exists $M_0 \leq M$ containing A which is λ^+ -saturated of size λ^+ . As in case 2 above, we build $\langle N_i : i < \lambda^+ \rangle \leq^0$ -increasing in K' such that $N_i \leq M$ and $p \restriction N_{i+1}$ i -fork over N_i . This is possible (for the successor step, given N_i , take any $M_0 \leq M$ saturated of size λ^+ containing N_i . By definition of i' and the fact p i' -forks over M_0 , there exists $N'_{i+1} \leq M$ in K' witnessing the forking. This can further extended to N_{i+1} which is as desired). This is enough: we get a contradiction to weak chain local character.

□

7. WEAKLY GOOD INDEPENDENCE RELATIONS

Interestingly, nonsplitting and $(< \kappa)$ -coheir (for a suitable choice of κ) are already well-behaved if the AEC is stable. This raises the question of whether there is an object playing the role of a good frame (see the next section) in strictly stable classes. Note that [BGKV] proves the canonicity of independence relations that satisfy much less than the full properties of good frames, so it is reasonable to expect existence of such an object. The next definition comes from extracting all the

properties we are able to prove from the construction of a good frame in [Vas] assuming only stability.

Definition 7.1. Let $\mathbf{i} = (K, \perp)$ be a $(< \alpha, \mathcal{F})$ -independence relation, $\mathcal{F} = [\lambda, \theta]$. \mathbf{i} is *weakly good*²⁷ if:

- (1) K is nonempty, is λ -closed (Recall Definition 2.11), and every chain in K of ordinal length less than θ has an upper bound.
- (2) K is stable in λ .
- (3) \mathbf{i} has base monotonicity, disjointness, existence, and transitivity.
- (4) $\text{pre}(\mathbf{i})$ has uniqueness.
- (5) \mathbf{i} has the left λ -witness property and the right λ -model-witness property.
- (6) Local character: For all $\alpha_0 < \min(\lambda^+, \alpha)$, $\bar{\kappa}_{\alpha_0}(\mathbf{i}) = \lambda^+$.
- (7) Local extension and uniqueness: $\mathbf{i}_\lambda^{<\lambda^+}$ has extension and uniqueness.

We say a pre- $(< \alpha, \mathcal{F})$ -frame \mathfrak{s} is *weakly good* if $\text{cl}(\mathfrak{s})$ is weakly good. \mathbf{i} is *pre-weakly good* if $\text{pre}(\mathbf{i})$ is weakly good.

Remark 7.2. By Propositions 4.3.(4), 4.3.(6), existence and the right λ -witness property follow from the others.

Our main tool to build weakly good independence relations will be to start from a λ -generator (see Definition 6.1) which satisfies some additional properties:

Definition 7.3. (K, \mathbf{i}) is a λ -generator for a weakly good $(< \alpha)$ -independence relation if:

- (1) (K, \mathbf{i}) is a λ -generator for a $(< \alpha)$ -independence relation.
- (2) K is nonempty, has no maximal models, and is stable in λ .
- (3) $(K^{\text{up}})^{\lambda^+ \text{-sat}}$ is λ -tame for types of length less than α .
- (4) \mathbf{i} has base monotonicity, existence, and is extended by λ -nonsplitting: whenever $p \in \text{gS}^{<\alpha}(M)$ does not \mathbf{i} -fork over $M_0 \leq M$, then p does not $\mathfrak{s}_{\lambda\text{-ns}}(K_{\mathbf{i}})$ -fork over M_0 .
- (5) (K, \mathbf{i}) has weak chain local character.

Both coheir and λ -nonsplitting induce a generator for a weakly good independence relation:

²⁷The name “weakly good” is admittedly not very inspired. A better choice may be to rename good independence relations to superstable independence relations and weakly independence relations to stable independence relations. We did not want to change Shelah’s terminology here and wanted to make the relationship between “weakly good” and “good” clear.

Proposition 7.4. Let K be an AEC with amalgamation and let $\lambda \geq \text{LS}(K)$ be such that K_λ is nonempty, has no maximal models, and K is stable in λ . Let $2 \leq \alpha \leq \lambda^+$.

- (1) Let $\text{LS}(K) < \kappa \leq \lambda$. Assume that K is $(< \kappa)$ -tame and short for types of length less than α . Let $\mathbf{i} := (\mathbf{i}_{\kappa\text{-ch}}(K))^{<\alpha}$.
 - (a) If K does not have the $(< \kappa)$ -order property of length κ , $\kappa_{<\alpha}(\mathbf{i}) \leq \lambda^+$, and $K_\lambda^{\kappa\text{-sat}}$ is dense in K_λ , then $(K_\lambda, \mathbf{i}_\lambda)$ is a λ -generator for a weakly good $(< \alpha)$ -independence relation.
 - (b) If $\kappa = \beth_{\kappa}$, $(\alpha_0 + 2)^{<\kappa_r} \leq \lambda$ for all $\alpha_0 < \alpha$, then $(K_\lambda, \mathbf{i}_\lambda)$ is a λ -generator for a weakly good $(< \alpha)$ -independence relation.
- (2) Assume $\alpha \leq \omega$ and $K^{\lambda^+\text{-sat}}$ is λ -tame for types of length less than α . Then $(K_\lambda, (\mathbf{i}_{\lambda\text{-ns}}(K_\lambda))^{<\alpha})$ is a λ -generator for a weakly good $(< \alpha)$ -independence relation.
- (3) Let K' be a dense sub-AC of K such that $K^{\lambda^+\text{-sat}} \subseteq K'$ and let \mathbf{i} be a $(< \alpha, \geq \lambda)$ -independence relation with $K_{\mathbf{i}} = K'$, such that $\text{pre}(\mathbf{i})$ has uniqueness, \mathbf{i} has base monotonicity, and $\bar{\kappa}_{<\alpha}(\mathbf{i}) = \lambda^+$. If K'_λ is dense in K_λ , then $(K_\lambda, \mathbf{i}_\lambda)$ is a λ -generator for a weakly good $(< \alpha)$ -independence relation.

Proof.

- (1) (a) By Fact 3.17, \mathbf{i} has base monotonicity, existence, and uniqueness. By Fact 3.20.(3), coheir is extended by λ -nonsplitting. The other properties are easy. For example, weak chain local character follows from $\kappa_{<\alpha}(\mathbf{i}) \leq \lambda^+$ and monotonicity.
- (b) We check that K and \mathbf{i} satisfy all the conditions of the previous part. By Fact 2.20, K does not have the $(< \kappa)$ -order property of length κ . By (the proof of) Proposition 4.3.(5) and Fact 3.17:

$$\kappa_{<\alpha}(\mathbf{i}) \leq \bar{\kappa}_{<\alpha}(\mathbf{i}) \leq \sup_{\alpha_0 < \alpha} ((\alpha_0 + 2)^{<\kappa_r})^+ \leq \lambda^+$$

Since K is stable in λ , if $\kappa < \lambda$ then $K_\lambda^{\kappa\text{-sat}}$ is dense in K_λ . If $\kappa = \lambda$, then $\kappa = 2^{<\kappa_r}$ so is regular, hence strongly inaccessible, so $\kappa = \kappa^{<\kappa}$ so again it is easy to check that $K_\lambda^{\kappa\text{-sat}}$ is dense in K_λ .

- (2) Let $\mathbf{i} := (\mathbf{i}_{\lambda\text{-ns}}(K))^{<\alpha}$. By Fact 3.20.(2) and Proposition 4.3.(5), $\kappa_{<\alpha}(\mathbf{i}) = \lambda^+$. By monotonicity, weak chain local character follows. The other properties are easy to check.

- (3) By Fact 3.20.(3), \mathbf{i} is extended by λ -nonsplitting. Weak chain character follows from $\kappa_{<\alpha}(\mathbf{i}) = \lambda^+$. By (the proof of) Proposition 4.6, $K^{\lambda^+-\text{sat}}$ is λ -tame for types of length less than α . The other properties are easy to check.

□

The next result is that a generator for a weakly good independence relation indeed induces a weakly good independence relation.

Theorem 7.5. Let (K, \mathbf{i}) be a λ -generator for a weakly good $(< \alpha)$ -independence relation. Then $(K, \mathbf{i})^{\text{up}} \upharpoonright (K^{\text{up}})^{\lambda^+-\text{sat}}$ is a pre-weakly good $(< \alpha, \geq \lambda^+)$ -independence relation.

Proof. This follows from the methods of [Vas], but we give some details. Let $\mathbf{i}' := (K, \mathbf{i})^{\text{up}} \upharpoonright (K^{\text{up}})^{\lambda^+-\text{sat}}$. Let $\perp := \perp_{\mathbf{i}'}$, $K' := K_{\mathbf{i}'}$, $\mathbf{s}' := \text{pre}(\mathbf{i}')$. We check the conditions in the definition of a weakly good independence relation. Note that by Remark 7.2 we do not need to check existence or the right λ^+ -witness property.

- \mathbf{i}' is a $(< \alpha, \geq \lambda^+)$ -independence relation: By Lemma 6.5.
- $K_{\mathbf{i}'}$ is stable in λ^+ : By Fact 2.20, K^{up} is stable in λ^+ . By stability, $K_{\mathbf{i}'}$ is dense in K so by Proposition 5.8, $K_{\mathbf{i}'}$ is stable in λ^+ .
- $K_{\mathbf{i}'} \neq \emptyset$ since it is dense in $K_{\lambda^+}^{\text{up}}$ and $K_{\lambda^+}^{\text{up}} = K$ is nonempty and has no maximal models. Every chain $\langle M_i : i < \delta \rangle$ in $K_{\mathbf{i}'}$ has an upper bound: we have $M_\delta := \bigcup_{i < \delta} M_i \in K$, and by density there exists $M \geq M_\delta$ in $K_{\mathbf{i}'}$. $K_{\mathbf{i}'}$ is λ^+ -closed by an easy increasing chain argument, using stability in λ^+ .
- Local character: $\bar{\kappa}_{<\alpha}(\mathbf{i}') = \lambda^{++}$ by Lemma 6.8.
- \mathbf{s}' has:
 - Base monotonicity: By Lemma 6.8.
 - Uniqueness: First observe that using local character, base monotonicity, λ^+ -closure, and the fact that $K_{\mathbf{i}'}$ is λ^+ -tame for types of length less than α , it is enough to show uniqueness for $(\mathbf{s}')_{\lambda^+}$. For this imitate the proof of [Vas, Lemma 5.3] (the key is weak uniqueness: Fact 3.20.(5)).
 - Local extension: Let $p \in \text{gS}^{<\alpha}(M)$, $M_0 \leq M \leq N$ be in $(K_{\mathbf{i}'})_{\lambda^+}$ such that p does not fork over M_0 . Let $M'_0 \leq M_0$ be in K' and witness it. By homogeneity, $M'_0 <_{\text{univ}} M$ so there exists $f : N \xrightarrow{M'_0} M$. Let $q := f^{-1}(p) \upharpoonright N$. By invariance, q does not fork over M_0 (as witnessed by M'_0).

Since λ -nonsplitting extends nonforking, $p \upharpoonright M'$ does not $\mathfrak{s}_{\lambda\text{-ns}}(K')$ -fork over M'_0 whenever $M'_0 \leq M' \leq M$ is such that $M' \in K'$. Let $K'' := K' \cup K^{\lambda^+-\text{sat}}$. By (the proof of) Fact 3.20.(4), p does not $\mathfrak{s}_{\text{ns}}(K'')$ -fork over M'_0 . By weak extension (Fact 3.20.(5)), q extends p and is algebraic if and only if q is.

- Transitivity: Imitate the proof of [Vasa, Lemma 4.7].
- Disjointness: It is enough to prove it for types of length 1 so assume $\alpha = 2$. Assume $a \underset{M_0}{\downarrow}^N M$ (with $M_0 \leq M \leq N$ in $K^{\lambda^+-\text{sat}}$) and $a \in M$. We show $a \in M_0$. Using local character, we can assume without loss of generality that $\|M_0\| = \lambda^+$ and (by taking a submodel of M containing a of size λ^+) that also $\|M\| = \lambda^+$. Find $M'_0 \leq M_0$ in K' witnessing the nonforking. By the proof of local extension, we can find $p \in \text{gS}(M)$ extending $p_0 := \text{gtp}(a/M_0; N)$ such that p_0 is algebraic if and only if p is. Since $a \in N$, we must have by uniqueness that p is algebraic so p_0 is algebraic, i.e. $a \in M_0$.

Now by Proposition 4.1, $\text{cl}(\mathfrak{s}')$ has the above properties.

- $\text{cl}(\mathfrak{s}')$ has the left λ -witness property: Because $\alpha \leq \lambda^+$.

□

Interestingly, the generator can always be taken to have a particular form:

Lemma 7.6. Let (K, \mathfrak{i}) be a λ -generator for a weakly good ($< \alpha$)-independence relation. Let $\mathfrak{i}' := \mathfrak{i}_{\lambda\text{-ns}}(K)^{<\alpha}$. Then:

- (1) (K, \mathfrak{i}') is a λ -generator for a weakly good ($< \alpha$)-independence relation and $<_{\text{univ}}$ is the ordering witnessing weak chain local character.
- (2) $\text{pre}((K, \mathfrak{i})^{\text{up}}) \upharpoonright (K^{\text{up}})^{\lambda^+-\text{sat}} = \text{pre}((K, \mathfrak{i}')^{\text{up}}) \upharpoonright (K^{\text{up}})^{\lambda^+-\text{sat}}$.

Proof.

- (1) By Lemma 6.7 (with K, \mathfrak{i}', K_i here standing for K, \mathfrak{i}, K'' there), (K, \mathfrak{i}') has weak chain local character (witnessed by $<_{\text{univ}}$) and the other properties are easy to check.
- (2) Let $\mathfrak{s} := \text{pre}((K, \mathfrak{i})^{\text{up}}) \upharpoonright (K^{\text{up}})^{\lambda^+-\text{sat}}$, $\mathfrak{s}' := \text{pre}((K, \mathfrak{i}')^{\text{up}}) \upharpoonright (K^{\text{up}})^{\lambda^+-\text{sat}}$. We want to see that $\underset{\mathfrak{s}}{\downarrow} = \underset{\mathfrak{s}'}{\downarrow}$. Since $\text{pre}(\mathfrak{i})$ is extended by λ -nonsplitting, it is easy to check that $\underset{\mathfrak{s}}{\downarrow} \subseteq \underset{\mathfrak{s}'}{\downarrow}$. By the proof of

[BGKV, Lemma 4.1], $\downarrow_{\mathfrak{s}_{\lambda^+}} = \downarrow_{\mathfrak{s}'_{\lambda^+}}$. By the right λ -model-witness property, $\downarrow_{\mathfrak{s}} = \downarrow_{\mathfrak{s}'}$.

□

In Theorem 7.5, $\mathfrak{i}' := (K, \mathfrak{i})^{\text{up}} \upharpoonright (K^{\text{up}})^{\lambda^+ \text{-sat}}$ is only pre-weakly good, not necessarily weakly good: in general, only $\mathfrak{i}'' := \text{cl}(\text{pre}(\mathfrak{i}'))$ will be weakly good. The following technical lemma shows that \mathfrak{i}' and \mathfrak{i}'' agree on slightly more than $\text{pre}(\mathfrak{i}')$.

Lemma 7.7. Let (K, \mathfrak{i}) be a λ -generator for a weakly good $(< \alpha)$ -independence relation. Let $\mathfrak{i}' := (K, \mathfrak{i})^{\text{up}} \upharpoonright (K^{\text{up}})^{\lambda^+ \text{-sat}}$ and let $\mathfrak{i}'' := \text{cl}(\text{pre}(\mathfrak{i}'))$. Let $M \leq N$ be in $K^{\text{up}}_{\geq \lambda^+}$ with $M \in K^{\lambda^+ \text{-sat}}$ (but maybe $N \notin K^{\lambda^+ \text{-sat}}$). Assume K^{up} is λ -tame²⁸ for types of length less than α . Let $p \in \text{gS}^{< \alpha}(N)$.

If $\|N\| = \lambda^+$ or \mathfrak{i}'' has extension, then p does not \mathfrak{i}' -fork over M if and only if p does not \mathfrak{i}'' -fork over M .

Proof. Assume p does not \mathfrak{i}'' -fork over M . Then by definition there exists an extension of p to a model in $K^{\lambda^+ \text{-sat}}$ that does not \mathfrak{i}' -fork over M so by monotonicity p does not \mathfrak{i}' -fork over M . Assume now that p does not \mathfrak{i}' -fork over M . Note that the proof of Theorem 7.5 (more precisely [Vasa, Lemma 5.3]) implies that p is the unique type over N that does not \mathfrak{i}' -fork over M .

Pick $N' \geq N$ in $K^{\lambda^+ \text{-sat}}$ with $\|N'\| = \|N\|$. We imitate the proof of [BGKV, Lemma 4.1]. By extension (or local extension if $\|N\| = \lambda^+$, recall that \mathfrak{i}'' is weakly good, see Theorem 7.5), there exists $q \in \text{gS}^{< \alpha}(N')$ that does not \mathfrak{i}'' -fork over M and extends $p \upharpoonright M$. By the above, q does not \mathfrak{i}' -fork over M . By uniqueness, q extends p , so $q \upharpoonright N = p$ does not \mathfrak{i}'' -fork over M . □

Note that if the independence relation of the generator is coheir, then the weakly good independence relation obtained from it is also coheir. We first prove a slightly more abstract lemma:

Lemma 7.8. Let K be an AEC, $\lambda \geq \text{LS}(K)$. Let K' be a dense sub-AC of K such that $K^{\lambda^+ \text{-sat}} \subseteq K'$ and K'_λ is dense in K_λ . Let \mathfrak{i} be a $(< \alpha, \geq \lambda)$ -independence relation with base monotonicity and

²⁸Note that the definition of a generator for a weakly good independence relation only requires that $(K^{\text{up}})^{\lambda^+ \text{-sat}}$ be λ -tame for types of length less than α .

$K_i = K'$, $2 \leq \alpha \leq \lambda^+$. Assume that \mathbf{i} has base monotonicity and the right λ -model-witness property.

Assume $\bar{\kappa}_{<\alpha}(\mathbf{i}) = \lambda^+$ and $(K_\lambda, \mathbf{i}_\lambda)$ is a λ -generator for a weakly good $(< \alpha)$ -independence relation. Let $\mathbf{i}' := (K_\lambda, \mathbf{i}_\lambda)^{\text{up}} \upharpoonright K^{\lambda^+-\text{sat}}$. Then $\text{pre}(\mathbf{i}') = \text{pre}(\mathbf{i}) \upharpoonright K^{\lambda^+-\text{sat}}$.

Moreover if \mathbf{i} has the right λ -witness property, then $\mathbf{i}' = \mathbf{i} \upharpoonright K^{\lambda^+-\text{sat}}$.

Proof. We prove the moreover part and it will be clear how to change the proof to prove the weaker statement (just replace the use of the witness property by the model-witness property).

Let $M \leq N$ be in $K^{\lambda^+-\text{sat}}$, $p \in \text{gS}^{<\alpha}(B; N)$. We want to show that p does not \mathbf{i} -fork over M if and only if there exists $M_0 \leq M$ in K'_λ so that for all $B_0 \subseteq B$ of size $\leq \lambda$, $p \upharpoonright B_0$ does not \mathbf{i} -fork over M_0 . Assume first that p does not \mathbf{i} -fork over M . Since $\bar{\kappa}_{<\alpha}(\mathbf{i}) = \lambda^+$, there exists $M_0 \leq M$ in K_λ such that p does not \mathbf{i} -fork over M_0 . By base monotonicity and homogeneity, we can assume that $M_0 \in K'_\lambda$. In particular $p \upharpoonright B_0$ does not \mathbf{i} -fork over B_0 for all $B \subseteq B$ of size $\leq \lambda$.

Conversely, assume p does not \mathbf{i}' -fork over M , and let $M_0 \leq M$ in K'_λ witness it. Then by the right λ -witness property, p does not \mathbf{i} -fork over M_0 , so over M , as desired.

□

Lemma 7.9. Let K be an AEC, $\text{LS}(K) < \kappa \leq \kappa' \leq \lambda$. Let $2 \leq \alpha \leq \lambda^+$. Let $\mathbf{i} := (\mathbf{i}_{\kappa\text{-ch}}(K))_{\geq \lambda}^{<\alpha} \upharpoonright K^{\kappa'\text{-sat}}$.

Assume $\bar{\kappa}_{<\alpha}(\mathbf{i}) = \lambda^+$ and $(K_\lambda, \mathbf{i}_\lambda)$ is a λ -generator for a weakly good $(< \alpha)$ -independence relation. Let $\mathbf{i}' := (K_\lambda, \mathbf{i}_\lambda)^{\text{up}} \upharpoonright K^{\lambda^+-\text{sat}}$. Then $\mathbf{i}' = \mathbf{i} \upharpoonright K^{\lambda^+-\text{sat}}$.

Proof. By Lemma 7.8 applied with $K' = K^{\kappa'\text{-sat}}$.

□

We end this section by showing how to build a weakly good independence relation in any stable fully tame and short AEC (with amalgamation and no maximal models).

Theorem 7.10. Let K be a $\text{LS}(K)$ -tame AEC with amalgamation and no maximal models. Let $\kappa = \beth_\kappa > \text{LS}(K)$. Assume K is stable and $(< \kappa)$ -tame and short for types of length less than α , $\alpha \geq 2$.

If $K_\kappa \neq \emptyset$, then $\mathbf{i}_{\kappa\text{-ch}}(K)^{<\alpha} \upharpoonright K^{(2^\kappa)^+-\text{sat}}$ is a pre-weakly good $(< \alpha, \geq (2^\kappa)^+)$ -independence relation. Moreover if $\alpha = \infty$, then it is weakly good.

Proof. Let $\lambda := 2^\kappa$. By Fact 3.17, $\mathbf{i}_{\kappa\text{-ch}}(K)^{<\alpha} \restriction K^{\lambda^+ \text{-sat}}$ already has many properties of a weakly good independence relation, and in particular has the left λ -witness property so it is enough to check that $\mathbf{i} := \mathbf{i}_{\kappa\text{-ch}}(K)^{<(\min(\alpha, \lambda^+))} \restriction K^{\lambda^+ \text{-sat}}$ is weakly good, so assume now without loss of generality that $\alpha \leq \lambda^+$. Note that by Fact 3.17, $\bar{\kappa}_{<\alpha}(\mathbf{i}) \leq (\lambda^\kappa)^+ = \lambda^+$. By Lemma 7.9 it is enough to check that $(K_\lambda, \mathbf{i}_\lambda)$ is a λ -generator for a weakly good ($< \alpha$)-independence relation. From Fact 2.20, we get that K is stable in λ . Finally, note that $K_\lambda \neq \emptyset$. Now apply Proposition 7.4.

If $\alpha = \infty$, then by Fact 3.17, \mathbf{i} has uniqueness. Since \mathbf{i} is pre-weakly good, $\text{pre}(\mathbf{i}_\lambda)$ has extension, so by Proposition 4.1.(7), \mathbf{i}_λ also has extension. The other properties of a weakly good independence relation follow from Fact 3.17. \square

8. GOOD INDEPENDENCE RELATIONS

Good frames were introduced by Shelah [She09, Definition II.2.1] as a “bare bone” definition of superstability in AECs. Here we adapt Shelah’s definition to independence relations. We also define a variation, being *fully* good. This is only relevant when the types are allowed to have length $\geq \lambda$, and asks for more continuity (like in [BVb], but the continuity property asked for is different). This is used to enlarge a good frame in the last sections.

Definition 8.1.

- (1) A *good* ($< \alpha, \mathcal{F}$)-independence relation $\mathbf{i} = (K, \perp)$ is a ($< \alpha, \mathcal{F}$)-independence relation satisfying:
 - (a) K is a nonempty AEC in \mathcal{F} , $\text{LS}(K) = \lambda_i$, K has no maximal models and joint embedding, K is stable in all cardinals in \mathcal{F} .
 - (b) \mathbf{i} has base monotonicity, disjointness, symmetry, uniqueness, extension, the left λ_i -witness property, and for all $\alpha_0 < \alpha$, $\kappa_{\alpha_0}(\mathbf{i}) = |\alpha_0|^+ + \aleph_0$ and $\bar{\kappa}_{\alpha_0}(\mathbf{i}) = |\alpha_0|^+ + \lambda_i^+$.
- (2) A *type-full good* ($< \alpha, \mathcal{F}$)-frame \mathfrak{s} is a pre- $(< \alpha, \mathcal{F})$ -frame so that $\text{cl}(\mathfrak{s})$ is good.
- (3) \mathbf{i} is *pre-good* if $\text{pre}(\mathbf{i})$ is good.

When we add “fully”, we require in addition that the frame/independence relation satisfies full model-continuity.

Remark 8.2. This paper’s definition is equivalent to that of Shelah [She09, Definition II.2.1] if we remove the requirement there on the

existence of a superlimit (as was done in almost all subsequent papers, for example in [JS13]) and assume the frame is type-full (i.e. the basic types are all the nonalgebraic types). For example, the continuity property that Shelah requires follows from $\kappa_1(\mathfrak{s}) = \aleph_0$ ([She09, Claim II.2.17.3]).

Remark 8.3. If \mathfrak{i} is a good $(< \alpha, \mathcal{F})$ -independence relation (except perhaps for the symmetry axiom) then \mathfrak{i} is weakly good.

Definition 8.4. An AEC K is *[fully] $(< \alpha, \mathcal{F})$ -good* if there exists a [fully] $(< \alpha, \mathcal{F})$ -good independence relation \mathfrak{i} with $K_{\mathfrak{i}} = K$. When $\alpha = \infty$ and $\mathcal{F} = [\text{LS}(K), \infty)$, we omit them.

As in the previous section, we give conditions for a generator to induce a good independence relation:

Definition 8.5. (K, \mathfrak{i}) is a λ -generator for a good $(< \alpha)$ -independence relation if:

- (1) (K, \mathfrak{i}) is a λ -generator for a weakly good $(< \alpha)$ -independence relation.
- (2) K^{up} is λ -tame.
- (3) There exists $\mu \geq \lambda$ such that K_{μ}^{up} has joint embedding.
- (4) Local character: For all $\alpha_0 < \min(\alpha, \lambda)$, there exists an ordering \trianglelefteq such that $(K_{\mathfrak{i}}, \trianglelefteq)$ is a skeleton of K and $\kappa_{\alpha_0}(\mathfrak{i}, \trianglelefteq) = |\alpha_0|^+ + \aleph_0$.

Remark 8.6. If (K, \mathfrak{i}) is a λ -generator for a good $(< \alpha)$ -independence relation, then it is a λ -generator for a weakly good $(< \alpha)$ -independence relation. Moreover if $\alpha < \lambda^+$, the weak chain local character axiom follows from the local character axiom.

As before, the generator can always be taken to be of a particular form:

Lemma 8.7. Let (K, \mathfrak{i}) be a λ -generator for a good $(< \alpha)$ -independence relation. Let $\mathfrak{i}' := \mathfrak{i}_{\lambda\text{-ns}}(K)^{<\alpha}$. Then:

- (1) (K, \mathfrak{i}') is a λ -generator for a good $(< \alpha)$ -independence relation and $<_{\text{univ}}$ is the ordering witnessing local character.
- (2) $\text{pre}((K, \mathfrak{i})^{\text{up}}) \upharpoonright (K^{\text{up}})^{\lambda^+ \text{-sat}} = \text{pre}((K, \mathfrak{i}')^{\text{up}}) \upharpoonright (K^{\text{up}})^{\lambda^+ \text{-sat}}$.

Proof.

- (1) By Lemma 6.7 (with $K, \mathfrak{i}', K_{\mathfrak{i}}$ here standing for K, \mathfrak{i}, K'' there), (K, \mathfrak{i}') has the local character properties, witnessed by $<_{\text{univ}}$, and the other properties are easy to check.
- (2) By Lemma 7.6.

□

Unfortunately it is not strictly true that a generator for a good $(< \alpha)$ -independence relation induces a good independence relation. For one thing, the extension property is problematic when $\alpha > \omega$ and this in turn creates trouble in the proof of symmetry. Also, we are unable to prove $K^{\lambda^+ \text{-sat}}$ is an AEC (although we suspect it should be true, see also Fact 10.18). For the purpose of stating a clean result, we introduce the following definition:

Definition 8.8. \mathbf{i} is an *almost pre-good* $(< \alpha, \mathcal{F})$ -independence relation if:

- (1) It is a pre-weakly good $(< \alpha, \mathcal{F})$ -independence relation.
- (2) It satisfies all the conditions in the definition of a *pre-good* independence relation except that:
 - (a) $K_{\mathbf{i}}$ is not required to be an AEC.
 - (b) $\text{cl}(\text{pre}(\mathbf{i}))$ is not required to have extension or uniqueness, but we still ask that $\text{pre}(\mathbf{i}^{<\omega})$ has extension.
 - (c) $\text{cl}(\text{pre}(\mathbf{i}))$ is not required to have symmetry, but we still require that $\text{pre}(\mathbf{i}^{<\omega})$ has full symmetry.
 - (d) We replace the condition on $\kappa_{\alpha_0}(\text{cl}(\text{pre}(\mathbf{i})))$ by:
 - (i) $\kappa_{<(\min(\alpha, \omega))}(\text{cl}(\text{pre}(\mathbf{i}))) = \aleph_0$.
 - (ii) For all $\alpha_0 < \alpha$, $\kappa_{\alpha_0}(\mathbf{i}) = |\alpha_0|^+ + \aleph_0$.

Theorem 8.9. Let (K, \mathbf{i}) be a λ -generator for a good $(< \alpha)$ -independence relation. Then:

- (1) K^{up} has joint embedding and no maximal models.
- (2) K^{up} is stable in every $\mu \geq \lambda$.
- (3) $\mathbf{i}' := (K, \mathbf{i})^{\text{up}} \upharpoonright (K^{\text{up}})^{\lambda^+ \text{-sat}}$ is an almost pre-good $(< \alpha, \geq \lambda^+)$ -independence relation.
- (4) If $\alpha \leq \omega$ and $\mu \geq \lambda^+$ is such that $(K^{\text{up}})^{\mu \text{-sat}}$ is an AEC with Löwenheim-Skolem number μ , then $(\mathbf{i}')^{<\alpha} \upharpoonright (K^{\text{up}})^{\mu \text{-sat}}$ is a pre-good $(< \alpha, \geq \mu)$ -independence relation.

Proof. Again, this follows from the methods of [Vasa], but we give some details. We show by induction on $\theta \geq \lambda^+$ that $\mathfrak{s}' := \text{pre}(\mathbf{i}')_{[\lambda^+, \theta]}$ is a good frame, except perhaps for symmetry and the conditions in Definition 8.8. This gives (3) (use Proposition 4.3.(7) to get symmetry, the proof of [Vasa, Lemma 5.9] to get extension for types of finite length, and Lemma 7.7 to get (2(d)i) in Definition 8.8), and (4) together with (1),(2) (use Proposition 5.8) follow.

- \mathfrak{s}' is a weakly good $(< \alpha, [\lambda^+, \theta])$ -frame: By Theorem 7.5.
- Let $\mu \geq \lambda$ be such that K_μ^{up} has joint embedding. By amalgamation, $K_{\geq \mu}^{\text{up}}$ has joint embedding. Once it is shown that K^{up} has no maximal models, it will follow that K^{up} has joint embedding (every model of size $\geq \lambda$ extends to one of size μ). Note that joint embedding is never used in any of the proofs below.
- To prove that $K_{[\lambda, \theta]}^{\text{up}}$ has no maximal models, we can assume without loss of generality that $\alpha = 2$ and (by Lemma 8.7) that $\mathfrak{i} = \mathfrak{i}_{\lambda\text{-ns}}(K)$, with $\kappa_1(\mathfrak{i}, <_{\text{univ}}) = \aleph_0$. By the induction hypothesis (and the assumption that K has no maximal models), $K_{[\lambda, \theta]}^{\text{up}}$ has no maximal models. It remains to see that K_θ^{up} has no maximal models. Assume for a contradiction that $M \in K_\theta^{\text{up}}$ is maximal. Then it is easy to check that $M \in (K^{\text{up}})_\theta^{\theta\text{-sat}}$. Build $\langle M_i : i < \theta \rangle$ increasing continuous and $a \in |M|$ such that for all $i < \theta$:
 - (1) $M_i \leq M$.
 - (2) $M_i <_{\text{univ}} M_{i+1}$.
 - (3) $M_i \in K_{< \theta}^{\text{up}}$.
 - (4) $a \notin |M_i|$.

This is enough. Let $M_\theta := \bigcup_{i < \theta} M_i$. Note that $\|M_\theta\| = \theta$ and $a \in |M| \setminus |M_\theta|$, so $M_\theta < M$. By Lemma 2.27, $M_0 <_{\text{univ}} M_\theta$. Thus there exists $f : M \xrightarrow{M_0} M_\theta$ and since M is maximal f is an isomorphism. However M is maximal whereas M witnesses that M_θ is not maximal, so M cannot be isomorphic to M_θ , a contradiction.

This is possible. Imitate the proof of [Vasa, Lemma 5.12] (this is where it is useful that the generator is nonsplitting and the local character is witnessed by $<_{\text{univ}}$).

- K^{up} is stable in all $\mu \in [\lambda^+, \theta]$: Exactly as in the proof of [Vasa, Theorem 5.6].
- \mathfrak{s}' has base monotonicity, disjointness, and uniqueness because it is weakly good. For all $\alpha_0 < \alpha$, $\kappa_{\alpha_0}(\mathfrak{i}') = |\alpha_0|^+ + \aleph_0$, $\bar{\kappa}_{\alpha_0}(\mathfrak{s}') = |\alpha_0|^+ + \lambda^{++} = \lambda^{++}$ by Lemma 6.8.

□

Remark 8.10. Our proof of no maximal models above improves on [She09, Conclusion 4.13.3], as it does not use the symmetry property.

9. CANONICITY

In [BGKV], we gave conditions under which two independence relations are the same. There we strongly relied on the extension property, but coheir and weakly good frames only have a weak version of it. In this section, we show that if we just want to show two independence relations are the same over sufficiently saturated models, then the proofs become easier and the extension property is not needed. In addition, we obtain an explicit description of the forking relation. We conclude that coheir, weakly good frames, and good frames are (in a sense made precise below) canonical. This gives further evidence that these objects are not ad-hoc and answers several questions in [BGKV]. The results of this section are also used in Section 10 to show the equivalence between superstability and strong superstability.

Lemma 9.1 (The canonicity lemma). Let K be an AEC with amalgamation and let $\lambda \geq \text{LS}(K)$ be such that K is stable in λ . Let K' be a dense sub-AC of K such that $K^{\lambda^+-\text{sat}} \subseteq K'$ and K'_λ is dense in K_λ . Let \mathbf{i}, \mathbf{i}' be $(< \alpha, \geq \lambda)$ -independence relation with $K_{\mathbf{i}} = K_{\mathbf{i}'} = K'$. Let $\alpha_0 := \min(\alpha, \lambda^+)$.

If:

- (1) $\text{pre}(\mathbf{i})$ and $\text{pre}(\mathbf{i}')$ have uniqueness.
- (2) \mathbf{i} and \mathbf{i}' have base monotonicity, the left λ -witness property, and the right λ -model-witness property.
- (3) $\bar{\kappa}_{<\alpha_0}(\mathbf{i}) = \bar{\kappa}_{<\alpha_0}(\mathbf{i}') = \lambda^+$.

Then $\text{pre}(\mathbf{i}) \upharpoonright K^{\lambda^+-\text{sat}} = \text{pre}(\mathbf{i}') \upharpoonright K^{\lambda^+-\text{sat}}$, and if in addition both \mathbf{i} and \mathbf{i}' have the right λ -witness property, then $\mathbf{i} \upharpoonright K^{\lambda^+-\text{sat}} = \mathbf{i}' \upharpoonright K^{\lambda^+-\text{sat}}$.

Moreover for $M \leq N$ in $K^{\lambda^+-\text{sat}}$, $p \in \text{gS}^{<\alpha}(N)$ does not \mathbf{i} -fork over M if and only if for all $I \subseteq \ell(p)$ with $|I| \leq \lambda$, there exists $M_0 \leq M$ in K'_λ such that p^I does not $\mathfrak{s}_{\lambda\text{-ns}}(K')$ -fork over M_0 .

Proof. By Fact 2.14, we can assume without loss of generality that K has joint embedding. If $K_{\lambda^+} = \emptyset$, there is nothing to prove so assume $K_{\lambda^+} \neq \emptyset$. Using joint embedding, it is easy to see that K_λ is nonempty and has no maximal models. By the left λ -witness property, we can assume without loss of generality that $\alpha \leq \lambda^+$, i.e. $\alpha = \alpha_0$. By Proposition 7.4, (K, \mathbf{i}) and (K, \mathbf{i}') are λ -generators for a weakly good $(< \alpha)$ -independence relation. By Lemma 7.6, $\text{pre}((K, \mathbf{i})^{\text{up}}) \upharpoonright K^{\lambda^+-\text{sat}} = \text{pre}((K, \mathbf{i}')^{\text{up}}) \upharpoonright K^{\lambda^+-\text{sat}}$.

By Lemma 7.8, for $x \in \{\mathbf{i}, \mathbf{i}'\}$, $\text{pre}((K, x)^{\text{up}}) \upharpoonright K^{\lambda^+ \text{-sat}} = \text{pre}(x) \upharpoonright K^{\lambda^+ \text{-sat}}$, so the result follows (the definition of $(K, x)_{\geq \lambda}$ and Lemma 7.6 also give the moreover part). The moreover part of lemma 7.8 says that if $x \in \{\mathbf{i}, \mathbf{i}'\}$ has the right λ -witness property, then $(K, x)^{\text{up}} \upharpoonright K^{\lambda^+ \text{-sat}} = x \upharpoonright K^{\lambda^+ \text{-sat}}$, so in case both \mathbf{i} and \mathbf{i}' have the right λ -witness property, we must have $\mathbf{i} \upharpoonright K^{\lambda^+ \text{-sat}} = \mathbf{i}' \upharpoonright K^{\lambda^+ \text{-sat}}$. \square

Remark 9.2. If K is an AEC with amalgamation, K' is a dense sub-AC of K such that $K^{\lambda^+ \text{-sat}} \subseteq K'$ and K'_λ is dense in K_λ , and \mathbf{i} is a $(\leq 1, \geq \lambda)$ -independence relation with $K_{\mathbf{i}} = K'$ and base monotonicity, uniqueness, $\bar{\kappa}_1(\mathbf{i}) = \lambda^+$, then by the proof of Proposition 4.6 and Lemma 5.8 K is stable in any $\mu \geq \text{LS}(K)$ with $\mu = \mu^\lambda$.

Theorem 9.3 (Canonicity of coheir). Let K be an AEC with amalgamation. Let $\kappa = \beth_\kappa > \text{LS}(K)$. Assume K is $(< \kappa)$ -tame and short for types of length less than α , $\alpha \geq 2$.

Let $\lambda \geq \kappa$ be such that K is stable in λ and $(\alpha_0 + 2)^{< \kappa_r} \leq \lambda$ for all $\alpha_0 < \min(\lambda^+, \alpha)$. Let \mathbf{i} be a $(< \alpha, \geq \lambda)$ -independence relation so that:

- (1) $K' := K_{\mathbf{i}}$ is a dense sub-AC of K so that $K^{\lambda^+ \text{-sat}} \subseteq K'$ and K'_λ is dense in K_λ .
- (2) $\text{pre}(\mathbf{i})$ has uniqueness.
- (3) \mathbf{i} has base monotonicity, the left λ -witness property, and the right λ -model-witness property.
- (4) $\bar{\kappa}_{< \min(\lambda^+, \alpha)}(\mathbf{i}) = \lambda^+$.

Then $\text{pre}(\mathbf{i}) \upharpoonright K^{\lambda^+ \text{-sat}} = \text{pre}(\mathbf{i}_{\kappa\text{-ch}}(K)^{< \alpha}) \upharpoonright K^{\lambda^+ \text{-sat}}$. If in addition \mathbf{i} has the right λ -witness property, then $\mathbf{i} \upharpoonright K^{\lambda^+ \text{-sat}} = \mathbf{i}_{\kappa\text{-ch}}(K)^{< \alpha} \upharpoonright K^{\lambda^+ \text{-sat}}$.

Proof. By Fact 2.14, we can assume without loss of generality that K has joint embedding. If $K_{\lambda^+} = \emptyset$, there is nothing to prove so assume $K_{\lambda^+} \neq \emptyset$. By Fact 2.13, K has arbitrarily large models so no maximal models. Let $\mathbf{i}' := \mathbf{i}_{\kappa\text{-ch}}(K)^{< \alpha}$. By the proof of Proposition 7.4, $\mathbf{i}' \upharpoonright K'$ satisfies the hypotheses of Lemma 9.1. Moreover, it has the right $(< \kappa)$ -witness property so the result follows. \square

Theorem 9.4 (Canonicity of weakly good independence relations). Let K be an AEC with amalgamation and let $\lambda \geq \text{LS}(K)$. Let K' be a dense sub-AC of K such that $K^{\lambda^+ \text{-sat}} \subseteq K'$ and K'_λ is dense in K_λ . Let \mathbf{i}, \mathbf{i}' be weakly good $(< \alpha, \geq \lambda)$ -independence relations with $K_{\mathbf{i}} = K_{\mathbf{i}'} = K'$.

Then $\text{pre}(\mathbf{i}) \upharpoonright K^{\lambda^+ \text{-sat}} = \text{pre}(\mathbf{i}') \upharpoonright K^{\lambda^+ \text{-sat}}$. If in addition both \mathbf{i} and \mathbf{i}' have the right λ -witness property, then $\mathbf{i} \upharpoonright K^{\lambda^+ \text{-sat}} = \mathbf{i}' \upharpoonright K^{\lambda^+ \text{-sat}}$.

Proof. By definition of a weakly good independence relation, K'_λ is stable in λ . Therefore by Lemma 5.8 K_λ , and hence K , is stable in λ . Now apply Lemma 9.1. \square

Theorem 9.5 (Canonicity of good independence relations). If \mathfrak{i} and \mathfrak{i}' are good $(< \alpha, \geq \lambda)$ -independence relations with the same underlying AEC K , then $\mathfrak{i} \upharpoonright K^{\lambda^+-\text{sat}} = \mathfrak{i}' \upharpoonright K^{\lambda^+-\text{sat}}$.

Proof. By Theorem 9.4 (with $K' := K$), $\text{pre}(\mathfrak{i}) \upharpoonright K^{\lambda^+-\text{sat}} = \text{pre}(\mathfrak{i}') \upharpoonright K^{\lambda^+-\text{sat}}$. Since good independence relations have extension, Lemma 4.2 implies $\mathfrak{i} \upharpoonright K^{\lambda^+-\text{sat}} = \mathfrak{i}' \upharpoonright K^{\lambda^+-\text{sat}}$. \square

Recall that [BGKV, Question 6.14] asked if two good λ -frames with the same underlying AEC should be the same. We can make progress toward this question by slightly refining our methods. Note that the results below can be further adapted to work for not necessarily type-full frames (that is for two good frames, in Shelah's sense, with the same basic types and the same underlying AEC).

Lemma 9.6. Let \mathfrak{s} and \mathfrak{s}' be good $(< \alpha, \lambda)$ -frames with the same underlying AEC K and $\alpha \leq \lambda$. Let K' be the class of λ -limit models of K (recall Definition 2.21). Then $\mathfrak{s} \upharpoonright K' = \mathfrak{s}' \upharpoonright K'$.

Proof sketch. By Remark 2.26, $I(K') = 1$. Now refine the proof of Theorem 9.5 by replacing λ^+ -saturated models by $(\lambda, |\beta|^+ + \aleph_0)$ -limit models for each $\beta < \alpha$. Everything still works since one can use the weak uniqueness and extension properties of nonsplitting (Fact 3.20.(5)). \square

Theorem 9.7 (Canonicity of categorical good λ -frames). Let \mathfrak{s} and \mathfrak{s}' be good $(< \alpha, \lambda)$ -frames with the same underlying AEC K and $\alpha \leq \lambda$. If K is categorical in λ , then $\mathfrak{s} = \mathfrak{s}'$.

Proof. By Fact 2.24, K has a limit model, and so by categoricity any model of K is limit. Now apply Lemma 9.6. \square

Remark 9.8. The proof also gives an explicit description of forking: For $M_0 \leq M$ with M_0 a limit model, $p \in \text{gS}(M)$ does not \mathfrak{s} -fork over M_0 if and only if there exists $M'_0 <_{\text{univ}} M_0$ such that p does not $\mathfrak{s}_{\lambda\text{-ns}}$ -fork over M'_0 . Note that this is the definition of forking in [Vas01].

Note that Shelah's construction of a good λ -frame in [She09, Theorem II.3.7] relies on categoricity in λ , so Theorem 9.7 establishes that the frame there is canonical. We are still unable to show that the frame built in Theorem 10.16 is canonical in general, although it will be if

λ is the categoricity cardinal or if it is weakly successful (by [BGKV, Theorem 6.13]).

10. SUPERSTABILITY

Shelah has pointed out [She09, p. 19] that superstability in abstract elementary classes suffers from schizophrenia, i.e. there are several different possible definitions that are equivalent in elementary classes but not necessarily in AECs. The existence of a good ($\geq \lambda$)-frame is a possible candidate but it is very hard to check. Instead, one would like a simple definition that implies existence of a good frame.

Shelah claims in chapter IV of his book that solvability²⁹ ([She09, Definition IV.1.4]) is such a notion, but his justification is yet to appear (in [She]). Essentially, solvability says that certain classes of EM models are superlimits. On the other hand previous work (for example [She99, SV99, GVV]) all rely on a local character property for nonsplitting. This is even made into a definition of superstability in [Gro02, Definition 7.12]. In [Vasa] we gave a similar condition and used it with tameness to build a good frame. We pointed out that categoricity in a cardinal of cofinality greater than the tameness cardinal implied the superstability condition.

We now aim to show the same conclusion under categoricity in a high-enough cardinal of arbitrary cofinality. Recall the definition of superstability implicit in [GVV, Vasa], and stated explicitly in [Gro02, Definition 7.12]:

Definition 10.1 (Superstability). An AEC K is μ -superstable if:

- (1) $\text{LS}(K) \leq \mu$.
- (2) There exists $M \in K_\mu$ such that for any $M' \in K_\mu$ there is $f : M' \rightarrow M$ with $f[M'] <_{\text{univ}} M$.
- (3) $\kappa_1(\mathfrak{s}_{\mu\text{-ns}}(K_\mu), \leq_{\text{univ}}) = \aleph_0$.

We say K is μ -superstable⁺ if $K_{\geq \mu}$ is μ -superstable, has amalgamation, and is μ -tame. We may omit μ , in which case we mean there exists a value such that the definition holds, e.g. K is *superstable* if it is μ -superstable for some μ .

Remark 10.2. Using Fact 2.24, it is easy to check that Condition (2) above is equivalent to “ K_μ is nonempty, has amalgamation, joint embedding, no maximal models, and is stable in μ ”.

²⁹One can ask whether there are any implications between this paper’s definition of superstability and Shelah’s. We leave this to future work.

Remark 10.3. While Definition 10.1 makes sense in any AEC, here we focus on tame AECs with amalgamation, and will not study what happens to Definition 10.1 without these assumptions (although this can be done, see [GVV]). In other words, we will study “superstable⁺” rather than just “superstable”.

For technical reasons, we will also use the following version that uses coheir rather than nonsplitting.

Definition 10.4. An AEC K is κ -strongly μ -superstable if:

- (1) $\text{LS}(K) < \kappa \leq \mu$.
- (2) (2) in Definition 10.1 holds.
- (3) K does not have the $(< \kappa)$ -order property of length κ .
- (4) $K_{\mu}^{\kappa\text{-sat}}$ is dense in K_{μ} .
- (5) $\kappa_1(\mathbf{i}_{\kappa\text{-ch}}(K)_{\mu}, \leq_{\text{univ}}) = \aleph_0$.

As before, we may omit some parameters and say K is κ -strongly μ -superstable⁺ if there exists $\kappa_0 < \kappa$ such that $K_{\geq \kappa_0}$ is κ -strongly μ -superstable, has amalgamation, and is $(< \kappa)$ -tame.

It is not too hard to see that a μ -superstable⁺ AEC induces a generator for a good independence relation, but what if we have a generator of some other form (assume for example that $<_{\text{univ}}$ is replaced by $<_{\mu, \delta}$ in the definition)? This is the purpose of the next definition.

Definition 10.5. Let K be an AEC.

- (1) K is (μ, \mathbf{i}) -superstable⁺ if $\text{LS}(K) \leq \mu$ and (K_{μ}, \mathbf{i}) is a μ -generator for a good (≤ 1) -independence relation.
- (2) K is κ -strongly (μ, \mathbf{i}) -superstable⁺ if:
 - (a) $\text{LS}(K) < \kappa \leq \mu$.
 - (b) There exists $\kappa_0 < \kappa$ such that $K_{\geq \kappa_0}$ has amalgamation.
 - (c) K is $(< \kappa)$ -tame.
 - (d) K does not have the $(< \kappa)$ -order property of length κ .
 - (e) K is (μ, \mathbf{i}) -superstable⁺.
 - (f) $K_{\mathbf{i}} \subseteq K_{\mu}^{\kappa\text{-sat}}$ and $\mathbf{i} = \mathbf{i}_{\kappa\text{-ch}}(K)^{\leq 1} \upharpoonright K_{\mathbf{i}}$.

The terminology is justified by the next proposition which tells us that the existence of *any* generator is equivalent to superstability. It makes checking that superstability holds easier and we will use it freely.

Proposition 10.6. Let K be an AEC.

- (1) K is μ -superstable⁺ if and only if there exists \mathbf{i} such that K is (μ, \mathbf{i}) -superstable⁺.

- (2) K is κ -strongly μ -superstable⁺ if and only if there exists \mathbf{i} such that K is κ -strongly (μ, \mathbf{i}) -superstable⁺.

Proof.

- (1) Assume first that K is μ -superstable⁺. Then one can readily check (using Proposition 7.4 and Remark 10.2) that $(K_\mu, \mathbf{i}_{\mu\text{-ns}}(K)^{\leq 1})$ is a generator for a good independence relation, where the local character axiom is witnessed by \leq_{univ} . Conversely, assume that K is (μ, \mathbf{i}) -superstable⁺. By definition, $\text{LS}(K) \leq \mu$ and by definition of a generator $K_{\geq \mu}$ has amalgamation and is μ -tame. By Lemma 8.7, $(K_\mu, \mathbf{i}_{\mu\text{-ns}}(K)^{\leq 1})$ is a μ -generator for a good (≤ 1) -independence relation, and \leq_{univ} is the ordering witnessing local character. Thus K is μ -superstable⁺.
- (2) Assume first that K is κ -strongly μ -superstable⁺. Let $\kappa_0 < \kappa$ be such that $K_{\geq \kappa_0}$ has amalgamation. Assume without loss of generality that $\kappa_0 = \text{LS}(K)$ and that $K_{\geq \kappa_0} = K$. By (the proof of) Proposition 7.4, $(K_\mu, \mathbf{i}_{\kappa\text{-ch}}(K)_\mu^{\leq 1})$ is a μ -generator for a weakly good (≤ 1) -independence relation. By the other conditions, it is actually a μ -generator for a good (≤ 1) -independence relation. Conversely, assume that K is κ -strongly (μ, \mathbf{i}) -superstable⁺. We check the last two conditions in the definition of strong superstability, the others are straightforward. We know that (K_μ, \mathbf{i}) is a generator and $\mathbf{i} = \mathbf{i}_{\kappa\text{-ch}}(K)^{\leq 1} \upharpoonright K_{\mathbf{i}}$. Thus $K_{\mathbf{i}} \subseteq K_\mu^{\kappa\text{-sat}}$ is dense in K_μ , so $K_\mu^{\kappa\text{-sat}}$ is dense in K_μ . By Lemma 6.7, $\kappa_1(\mathbf{i}_{\kappa\text{-ch}}(K)_\mu, \leq_{\text{univ}}) \leq \kappa_1(\mathbf{i}, \trianglelefteq)$ for any \trianglelefteq such that $(K_{\mathbf{i}}, \trianglelefteq)$ is a skeleton of K_μ (and hence of $K_\mu^{\kappa\text{-sat}}$). By assumption one can find such a \trianglelefteq with $\kappa_1(\mathbf{i}, \trianglelefteq) = \aleph_0$. Thus

$$\kappa_1(\mathbf{i}_{\kappa\text{-ch}}(K)_\mu, \leq_{\text{univ}}) = \aleph_0$$

□

Remark 10.7. Thus in Definitions 10.1 and 10.4, one can replace \leq_{univ} by $\leq_{\mu, \delta}$ for $1 \leq \delta < \mu^+$.

The next result gives evidence that Definition 10.1 is a reasonable definition of superstability, at least in tame AECs with amalgamation. Note that most of it already appears implicitly in [Vasa] and essentially restates Theorem 8.9.

Theorem 10.8. Assume K is a (μ, \mathbf{i}) -superstable⁺ AEC. Then:

- (1) $K_{\geq \mu}$ has joint embedding, no maximal models, and is stable in all $\lambda \geq \mu$.

- (2) Let $\lambda \geq \mu^+$ and let $\mathbf{i}' := (K_\mu, \mathbf{i})^{\text{up}} \upharpoonright K_{\geq \lambda}^{\mu^+ \text{-sat}}$.
- (a) \mathbf{i}' is an almost pre-good $(\leq 1, \geq \lambda)$ -independence relation (recall Definition 8.8).
 - (b) If in addition K is κ -strongly (μ, \mathbf{i}) -superstable $^+$, then $\text{pre}(\mathbf{i}') = \text{pre}(\mathbf{i}_{\kappa\text{-ch}}(K))^{\leq 1} \upharpoonright K_{\geq \lambda}^{\mu^+ \text{-sat}}$. That is, the frame is $(< \kappa)$ -coheir.
 - (c) If $\theta \geq \mu^+$ is such that $K' := K_{\geq \lambda}^{\theta \text{-sat}}$ is an AEC with $\text{LS}(K') = \lambda$, then $\mathbf{i}' \upharpoonright K'$ is a pre-good $(\leq 1, \geq \lambda)$ -independence relation that will be $(< \kappa)$ -coheir if K is κ -strongly (μ, \mathbf{i}) -superstable $^+$.

Proof. Theorem 8.9 gives (1) and (2a), while (2c) follows from (2a) and (2b). It remains to prove (2b). Let $\mathbf{i}'' := \mathbf{i}_{\kappa\text{-ch}}(K))^{\leq 1} \upharpoonright K_{\geq \lambda}^{\mu^+ \text{-sat}}$. By the proof of Lemma 7.8, $\downarrow \subseteq \downarrow$. Now by (2a), $\text{pre}(\mathbf{i}')$ has existence and extension and by Fact 3.17, \mathbf{i}'' has uniqueness. By [BGKV, Lemma 4.1], $\text{pre}(\mathbf{i}') = \text{pre}(\mathbf{i}'')$, as desired. \square

Remark 10.9. Let T be a complete first-order theory and let $K := (\text{Mod}(T), \preceq)$. Then this paper's definitions of superstability and strong superstability coincide with the classical definition. More precisely for all $\mu \geq |T|$, K is (strongly) μ -superstable if and only if T is stable in all $\lambda \geq \mu$.

Note also that $[\text{strong}] \mu$ -superstability $^+$ is monotonic in μ :

Proposition 10.10. If K is $[\kappa\text{-strongly}] \mu$ -superstable $^+$ and $\mu' \geq \mu$, then K is $[\kappa\text{-strongly}] \mu'$ -superstable $^+$.

Proof. Say K is (μ, \mathbf{i}) -superstable $^+$. It is clearly enough to check that K is μ' -superstable. Let $\mathbf{i}' := (K, \mathbf{i})^{\text{up}} \upharpoonright K_{\geq \mu'}^{\mu^+ \text{-sat}}$. By Theorem 10.8 and Proposition 7.4, $(K_{\mu'}, \mathbf{i}')$ is a generator for a good μ' -independence relation, so K is (μ', \mathbf{i}') -superstable. Similarly, if K is κ -strongly (μ, \mathbf{i}) -superstable $^+$ then K will be κ -strongly (μ', \mathbf{i}') -superstable. \square

Theorem 10.8.(2b) is the reason we introduced strong superstability. While it may seem like a detail, we are interested in extending our good frame to a frame for types longer than one element and using coheir to do so seems reasonable. Using the canonicity of coheir, we can show that superstability and strong superstability are equivalent if we do not care about the parameter μ :

Theorem 10.11. If K is μ -superstable $^+$ and $\kappa = \beth_\kappa > \mu$, then K is κ -strongly $(2^{<\kappa_r})^+$ -superstable $^+$.

In particular a tame AEC with amalgamation is strongly superstable if and only if it is superstable.

Proof. Let $\mu' := (2^{<\kappa_r})^+$. We show that K is κ -strongly μ' -superstable⁺. By Theorem 10.8, $K_{\geq\mu}$ has joint embedding, no maximal models and is stable in all cardinals. By definition, $K_{\geq\mu}$ also has amalgamation. Also, K is μ -tame, hence $(<\kappa)$ -tame. By Fact 2.20, K does not have the $(<\kappa)$ -order property of length κ . Moreover we have already observed that $K_{\mu'}$ is stable in μ' and has joint embedding and no maximal models. Also, $K_{\mu'}^{\kappa\text{-sat}}$ is dense in $K_{\mu'}$ by stability and the fact $\mu' > \kappa$. It remains to check that $\kappa_1(\mathbf{i}_{\kappa\text{-ch}}(K)_{\mu'}, \leq_{\text{univ}}) = \aleph_0$.

By Theorem 10.8, there is a $(\leq 1, \geq \mu^+)$ -independence relation \mathbf{i}' such that $K_{\mathbf{i}'} = K^{\mu^+\text{-sat}}$ and \mathbf{i}' is good, except that $K_{\mathbf{i}'}$ may not be an AEC. By Theorem 9.3 (with λ there standing for $2^{<\kappa_r}$ here), $\text{pre}(\mathbf{i}') \upharpoonright K^{\mu'\text{-sat}} = \text{pre}(\mathbf{i}_{\kappa\text{-ch}}(K)^{\leq 1}) \upharpoonright K^{\mu'\text{-sat}}$. By the proof of Lemma 4.5, \mathbf{i}' has the right $(<\kappa)$ -witness property for members of $K_{\geq\mu^+}$: If $M \in K_{\geq\mu^+}$, $M_0 \leq M$ is in $K^{\mu^+\text{-sat}}$, and $p \in \text{gS}(M)$, then p does not \mathbf{i} -fork over M_0 if and only if $p \upharpoonright B$ does not \mathbf{i} -fork over M_0 for all $B \subseteq |M|$ with $|B| < \kappa$. Therefore by the proof of Theorem 9.3, we actually have that for any $M \in K_{\geq\mu'}$ and $M_0 \leq M$ in $K^{\mu'\text{-sat}}$, $p \in \text{gS}(M)$ does not \mathbf{i}' -fork over M_0 if and only if p does not $\mathbf{i}_{\kappa\text{-ch}}(K)$ -fork over M_0 . In particular:

$$\kappa_1(\mathbf{i}_{\kappa\text{-ch}}(K)_{\mu'}) = \kappa_1(\mathbf{i}'_{\mu'}) = \aleph_0$$

Therefore $\kappa_1(\mathbf{i}_{\kappa\text{-ch}}(K)_{\mu'}, \leq_{\text{univ}}) = \aleph_0$, as needed. □

We now arrive to the main result of this section: categoricity implies strong superstability. We first recall several known consequences of categoricity.

Fact 10.12. Let K be an AEC with no maximal models, joint embedding, and amalgamation. Assume K is categorical in a $\lambda > \text{LS}(K)$. Then:

- (1) K is stable in all $\mu \in [\text{LS}(K), \lambda)$.
- (2) For $\text{LS}(K) \leq \mu < \text{cf}(\lambda)$, $\kappa_1(\mathfrak{s}_{\mu\text{-ns}}(K_{\mu}), \leq_{\mu, \omega}) = \aleph_0$.
- (3) Assume K does not have the weak κ -order property (see Definition 2.19) and $\text{LS}(K) < \kappa \leq \mu < \lambda$. Then:

$$\kappa_1(\mathbf{i}_{\kappa\text{-ch}}(K)_{\mu}, \leq_{\text{univ}}) = \aleph_0$$

- (4) If the model of size λ is μ -saturated for $\mu > \text{LS}(K)$, then every member of $K_{\geq \chi}$ is μ -saturated, where $\chi := \min(\lambda, \sup_{\mu_0 < \mu} h(\mu_0))$.

Proof.

- (1) Use Ehrenfeucht-Mostowski models (see for example the proof of [Bal09, Theorem 8.21]).
- (2) By [She99, Lemma 6.3].
- (3) By [BG, Theorem 6.6].
- (4) See (the proof of) [BG, Theorem 5.4].

□

The next proposition is useful to obtain joint embedding and no maximal models if we already have amalgamation.

Proposition 10.13. Let K be an AEC with amalgamation. If there exists $\lambda \geq \text{LS}(K)$ such that K_λ has joint embedding, then there exists $\chi < h(\text{LS}(K))$ such that $K_{\geq \chi}$ has joint embedding and no maximal models.

Proof. Write $\mu := h(\text{LS}(K))$. If $K_\mu = \emptyset$, then by Fact 2.13 there exists $\chi < \mu$ such that $K_{\geq \chi} = \emptyset$, so it has joint embedding and no maximal models. Now assume $K_\mu \neq \emptyset$. In particular, K has arbitrarily large models. By amalgamation, $K_{\geq \lambda}$ has joint embedding, and so no maximal models. If $\lambda < \mu$ we are done so assume $\lambda \geq \mu$. It is enough to show that there exists $\chi < \mu$ such that $K_{\geq \chi}$ has no maximal model since then any model of $K_{\geq \chi}$ embeds inside a model in $K_{\geq \lambda}$ and hence $K_{\geq \chi}$ has joint embedding.

By Fact 2.14, we can write $K = \bigcup_{i \in I} K^i$ where the K^i 's are disjoint AECs with $\text{LS}(K^i) = \text{LS}(K)$ and each K^i has joint embedding and amalgamation. Note that $|I| \leq I(K, \text{LS}(K)) \leq 2^{\text{LS}(K)}$. For $i \in I$, let χ_i be the least $\chi < \mu$ such that $K_{\geq \chi}^i = \emptyset$, or $\text{LS}(K)$ if $K_\mu^i \neq \emptyset$. Let $\chi := \sup_{i \in I} \chi_i$. Note that $\text{cf}(\mu) = (2^{\text{LS}(K)})^+ > 2^{\text{LS}(K)} \geq |I|$, so $\chi < \mu$.

Now let $M \in K_{\geq \chi}$. Let $i \in I$ be such that $M \in K^i$. M witnesses that $K_\chi^i \neq \emptyset$ so by definition of χ , K^i has arbitrarily large models. Since K^i has joint embedding, this implies that K^i has no maximal models. Therefore there exists $N \in K^i \subseteq K$ with $M < N$, as desired. □

The next theorem is implicit in [Vasa]. It is a simple consequence of Fact 10.12.(2).

Theorem 10.14. Let K be an $\text{LS}(K)$ -tame AEC with amalgamation and no maximal models. If K is categorical in a λ with $\text{cf}(\lambda) > \text{LS}(K)$, then K is $\text{LS}(K)$ -superstable⁺.

Proof. By amalgamation, categoricity, and no maximal models, K has joint embedding. By Fact 10.12.(1), K is stable in $\text{LS}(K)$. Now apply Fact 10.12.(2) and Proposition 10.6 (with Remark 10.7). \square

Corollary 10.15. Let K be an $\text{LS}(K)$ -tame AEC with amalgamation. If K is categorical in a λ with $\text{cf}(\lambda) \geq h(\text{LS}(K))$, then there exists $\mu < h(\text{LS}(K))$ such that K is μ -superstable⁺.

Proof. By Proposition 10.13, there exists $\mu < h(\text{LS}(K))$ such that $K_{\geq \mu}$ has joint embedding and no maximal models. Now apply Theorem 10.14 to $K_{\geq \mu}$. \square

We now remove the restriction on the cofinality and get strong super-stability. The downside is that $h(\text{LS}(K))$ is replaced by a fixed point of the beth function above $\text{LS}(K)$.

Theorem 10.16. Let K be an AEC with amalgamation. Let $\kappa = \beth_{\kappa} > \text{LS}(K)$ and assume K is $(< \kappa)$ -tame. If K is categorical in a $\lambda > \kappa$, then:

- (1) K is κ -strongly κ -superstable⁺.
- (2) K is stable in all cardinals $\geq h(\text{LS}(K))$.
- (3) The model of size λ is saturated.
- (4) K is categorical in κ .
- (5) For $\chi := \min(\lambda, h(\kappa))$, $\text{pre}(\mathbf{i}_{\kappa\text{-ch}}(K)_{\leq \chi}^1)$ is a good $(\leq 1, \geq \chi)$ -frame with underlying AEC $K_{\geq \chi}$.

Proof. Note that K_{λ} has joint embedding so by Proposition 10.13, there exists $\chi_0 < h(\text{LS}(K))$ such that $K_{\geq \chi_0}$ (and thus $K_{\geq \kappa}$) has joint embedding and no maximal models. By Fact 10.12.(1), $K_{\geq \chi_0}$ is stable everywhere below λ . Since $\kappa = \beth_{\kappa}$, Fact 2.20 implies that K does not have the $(< \kappa)$ -order property of length κ .

Let $\kappa \leq \mu < \lambda$. By Fact 10.12.(3), $\kappa_1(\mathbf{i}_{\kappa\text{-ch}}(K)_{\mu}, \leq_{\text{univ}}) = \aleph_0$. Now using Proposition 10.6, K is κ -strongly μ -superstable if and only if $K_{\mu}^{\kappa\text{-sat}}$ is dense in K_{μ} . If $\kappa < \mu$, then $K_{\mu}^{\kappa\text{-sat}}$ is dense in K_{μ} (by stability), so K is κ -strongly μ -superstable. However we want κ -strong κ -superstability. We proceed in several steps.

First, we show K is μ -superstable for *some* $\mu < \lambda$. If $\lambda = \kappa^+$, then this follows directly from Theorem 10.14 with $\mu = \kappa$, so assume $\lambda >$

κ^+ . Then by the previous paragraph K is κ -strongly μ -superstable for $\mu := \kappa^+$.

Second, we prove (2). We have already observed $K_{\geq \chi_0}$ is stable everywhere below λ . By Theorem 10.8, K is stable in every $\mu' \geq \mu$. In particular, it is stable in and above λ , so (2) follows.

Third, we show (3). Since K is stable in λ , we can build a λ_0^+ -saturated model of size λ for all $\lambda_0 < \lambda$. Thus the model of size λ is λ_0^+ -saturated for all $\lambda_0 < \lambda$, and hence λ -saturated.

Fourth, we prove (4). Since the model of size λ is saturated, it is κ -saturated. By Fact 10.12.(4), every model of size $\sup_{\kappa_0 < \kappa} h(\kappa_0) = \kappa$ is κ -saturated. By uniqueness of saturated models, K is categorical in κ .

Fifth, observe that since every model of size κ is saturated, $K_{\kappa}^{\kappa\text{-sat}} = K_{\kappa}$ is dense in K_{κ} . By the second paragraph above, K is κ -strongly κ -superstable so (1) holds.

Finally, we prove (5). We have seen that the model of size λ is saturated, thus κ^+ -saturated. By Fact 10.12.(4), every model of size $\geq \chi$ is κ^+ -saturated. Now use (1) with Theorem 10.8. \square

Remark 10.17. If one just wants to get strong superstability from categoricity, we suspect it should be possible to replace the $\beth_{\kappa} = \kappa$ hypothesis by something more reasonable (maybe just asking for the categoricity cardinal to be above 2^{κ}). Since we are only interested in eventual behavior here, we leave this to future work.

As a final remark, we point out that it is always possible to get a good independence relation from superstability (i.e. even without categoricity) if one is willing to restrict the class to sufficiently saturated models:

Fact 10.18 (Corollary 4.5 in [BVa]). Let K be an AEC. If K is κ -strongly μ -superstable $^+$, then whenever $\lambda > (\mu^{<\kappa_r})^+$, $K^{\lambda\text{-sat}}$ is an AEC with $\text{LS}(K^{\lambda\text{-sat}}) = \lambda$.

Corollary 10.19. Let K be an AEC. If K is κ -strongly μ -superstable $^+$, then $K^{(\mu^{<\kappa_r})^+2\text{-sat}}$ is (≤ 1) -good. Moreover the good frame is induced by $(< \kappa)$ -coheir.

Proof. Combine Theorem 10.8.(2c) and Fact 10.18. \square

Remark 10.20. Let K be an AEC in $\lambda := \text{LS}(K)$ with amalgamation, joint embedding, and no maximal models. If $K^{\lambda\text{-sat}}$ is a nonempty AEC in λ , then the saturated model is superlimit (see [She09, Definition

1.13]). Thus we even obtain a good frame in the sense of [She09, Chapter II].

11. DOMINATION

Our next aim is to take a sufficiently nice good λ -frame (for types of length 1) and show that it can be extended to types of length $\leq \lambda$. To do this, we will give conditions under which a good λ -frame is *weakly successful* (a key technical property of [She09, Chapter II], see Definition 11.4), and even ω -successful (Definition 11.20).

The hypotheses we will work with are:

Hypothesis 11.1.

- (1) $\mathbf{i} = (K, \perp)$ is a $(< \infty, \geq \mu)$ -independence relation.
- (2) $\mathfrak{s} := \text{pre}(\mathbf{i}^{\leq 1})$ is a type-full good $(\geq \mu)$ -frame.
- (3) $\lambda > \mu$ is a cardinal.
- (4) For all $n < \omega$:
 - (a) $K^{\lambda^{+n}\text{-sat}}$ is an AEC³⁰ with Löwenheim-Skolem number λ^{+n} .
 - (b) $\kappa_{\lambda^{+n}}(\mathbf{i}) = \lambda^{+n+1}$.
- (5) \mathbf{i} has base monotonicity, $\text{pre}(\mathbf{i})$ has uniqueness.
- (6) \mathbf{i} has the left and right $(\leq \mu)$ -model-witness properties.

Remark 11.2. We could have given more local hypotheses (e.g. by replacing ∞ by θ or only assuming (4) for n below some fixed $m < \omega$) and made some of the required properties more precise (this is part of what should be done to improve “short” to “diagonally tame” in the main theorem, see the discussion in Section 15).

The key is that we assume there is *already* an independence notion for longer types. However, it is potentially quite weak compared to what we want. The next fact shows that our hypotheses are reasonable.

Fact 11.3. Assume K^0 is a fully $(< \kappa)$ -tame and short κ -strongly μ_0 -superstable AEC with amalgamation. Then for any $\mu \geq (\mu_0^{< \kappa_r})^{+2}$ and any $\lambda > \mu$ with $\lambda = \lambda^{< \kappa_r}$, Hypothesis 11.1 holds for $K := (K^0)^{\mu\text{-sat}}$ and $\mathbf{i} := \mathbf{i}_{\kappa\text{-ch}}(K^0) \upharpoonright K$.

Proof. By Fact 10.18, for any $\mu' \geq \mu$, $K^{\mu'\text{-sat}}$ is an AEC with $\text{LS}(K^{\mu'\text{-sat}}) = \mu'$. By Theorem 10.8.(2c), $(< \kappa)$ -coheir induces a good $(\geq \mu)$ -frame for μ -saturated models. The other conditions follow directly from the definition of strong superstability and the properties of coheir (Fact 3.17).

³⁰Thus we have a superlimit of size λ^{+n} , see Remark 10.20.

For example, the local character condition holds because $\lambda^{<\kappa_r} = \lambda$ implies $(\lambda^{+n})^{<\kappa_r} = \lambda^{+n}$ for any $n < \omega$. \square

The next technical property is of great importance in Chapter II and III of [She09]. The definition below follows [JS13, Definition 4.1.5] (but as usual, we work only with type-full frames).

Definition 11.4. Let \mathfrak{t} be a type-full good $\lambda_{\mathfrak{t}}$ -frame.

- (1) For $M_0 \leq M_\ell$ in K , $\ell = 1, 2$, an *amalgam of M_1 and M_2 over M_0* is a triple (f_1, f_2, N) such that $N \in K_{\mathfrak{t}}$ and $f_\ell : M_\ell \xrightarrow{M_0} N$.
- (2) Let (f_1^x, f_2^x, N^x) , $x = a, b$ be amalgams of M_1 and M_2 over M_0 . We say (f_1^a, f_2^a, N^a) and (f_1^b, f_2^b, N^b) are *equivalent over M_0* if there exists $N_* \in K_{\mathfrak{t}}$ and $f^x : N^x \rightarrow N_*$ such that $f^b \circ f_1^b = f^a \circ f_1^a$ and $f^b \circ f_2^b = f^a \circ f_2^a$, namely, the following commutes:

$$\begin{array}{ccccc}
 & & N^b & \xrightarrow{f^b} & N_* \\
 & \nearrow f_1^b & \uparrow f_2^b & & \uparrow f^a \\
 M_1 & \xrightarrow{f_1^a} & N^a & & \\
 \uparrow & & \uparrow & \nearrow f_2^a & \\
 M_0 & \longrightarrow & M_2 & &
 \end{array}$$

Note that being “equivalent over M_0 ” is an equivalence relation ([JS13, Proposition 4.3]).

- (3) Let $K_{\mathfrak{t}}^{3,\text{uq}}$ be the set of triples (a, M, N) such that $M \leq N$ are in K , $a \in |N| \setminus |M|$ and for any $M_1 \geq M$ in K , there exists a unique (up to equivalence over M_0) amalgam (f_1, f_2, N_1) of N and M_1 over M such that $\text{gtp}(f_1(a)/f_2[M_1]; N_1)$ does not fork over M . We call the elements of $K_{\mathfrak{t}}^{3,\text{uq}}$ *uniqueness triples*.
- (4) $K_{\mathfrak{t}}^{3,\text{uq}}$ has the *existence property* if for any $M \in K_{\mathfrak{t}}$ and any nonalgebraic $p \in \text{gS}(M)$, one can write $p = \text{gtp}(a/M; N)$ with $(a, M, N) \in K_{\mathfrak{t}}^{3,\text{uq}}$. We also talk about the *existence property for uniqueness triples*.
- (5) \mathfrak{s} is *weakly successful* if $K_{\mathfrak{t}}^{3,\text{uq}}$ has the existence property.

The uniqueness triples can be seen as describing a version of domination. They were introduced by Shelah for the purpose of starting with a good λ -frame and extending it to a good λ^+ -frame. The idea is to first extend the good λ -frame to a forking notion for types of models of size λ (and really this is what interests us here, since tameness already

gives us a good λ^+ -frame). Now, since we already have an independence notion for longer types, we can follow [MS90, Definition 4.21] and give a more explicit version of domination that is exactly as in the first-order case.

Definition 11.5 (Domination). Fix $N \in K$. For $M \leq N$, $B, C \subseteq |N|$, B *dominates* C over M in N if for any $N' \geq N$ and any $D \subseteq |N'|$, $B \downarrow_M^{N'} D$ implies $B \cup C \downarrow_M^{N'} D$.

We say that B *model-dominates* C over M in N if for any $N' \geq N$ and any $M \leq N'_0 \leq N'$, $B \downarrow_M^{N'} N'_0$ implies $B \cup C \downarrow_M^{N'} N'_0$.

Model-domination turns out to be the technical variation we need, but of course if \mathbf{i} has extension, then it is equivalent to domination. We start with two easy ambient monotonicity properties:

Lemma 11.6. Let $M \leq N$. Let $B, C \subseteq |N|$ and assume B [model-]dominates C over M in N . Then:

- (1) If $N' \geq N$, then B [model-]dominates C over M in N' .
- (2) If $M \leq N_0 \leq N$ contains $B \cup C$, then B [model-]dominates C over M in N_0 .

Proof. We only do the proofs for the non-model variation but of course the model variation is completely similar.

- (1) By definition of domination.
- (2) Let $N' \geq N_0$ and $D \subseteq |N'|$ be given such that $B \downarrow_M^{N'} D$. By amalgamation, there exists $N'' \geq N$ and $f : N' \xrightarrow[N_0]{N'} N''$. By invariance, $B \downarrow_M^{N''} f[D]$. By definition of domination, $B \cup C \downarrow_M^{f[N']}\ f[D]$. By invariance again, $B \cup C \downarrow_M^{N'} D$, as desired.

□

The next result is key for us: it ties domination with the notion of uniqueness triples:

Lemma 11.7. Assume $M_0 \leq M_1$ are in K_λ , and $a \in M_1$ model-dominates M_1 over M_0 (in M_1). Then $(a, M_0, M_1) \in K_{\mathfrak{s}_\lambda}^{3, \text{uq}}$.

Proof. Let $M_2 \geq M_0$ be in K_λ . First, we need to show that there exists (b, M_2, N) such that $\text{gtp}(b/M_2; N)$ extends $\text{gtp}(a/M_0; M_1)$ and $\text{gtp}(b/M_2; N)$ does not fork over M_0 . This holds by the extension property of good frames.

Second, we need to show that any such amalgam is unique: Let (f_1^x, f_2^x, N^x) , $x \in \{a, b\}$ be amalgams of M_1 and M_2 over M_0 such that $f_1^x(a) \downarrow_{M_0}^{N^x} f_2^x[M_2]$.

We want to show that the two amalgams are equivalent: we want $N_* \in K_\lambda$ and $f^x : N^x \rightarrow N_*$ such that $f^b \circ f_1^b = f^a \circ f_1^a$ and $f^b \circ f_2^b = f^a \circ f_2^a$, namely, the following commutes:

$$\begin{array}{ccccc}
 & & N^b & \xrightarrow{f^b} & N_* \\
 & \nearrow f_1^b & \uparrow f_2^b & & \uparrow f^a \\
 M_1 & \xrightarrow{f_1^a} & N^a & & \\
 \uparrow & & \uparrow & \nearrow f_2^a & \\
 M_0 & \longrightarrow & M_2 & &
 \end{array}$$

For $x = a, b$, rename f_2^x to the identity to get amalgams $((f_1^x)', \text{id}_{M_2}, (N^x)')$ of M_1 and M_2 over M_0 . For $x = a, b$, the amalgams $((f_1^x)', \text{id}_{M_2}, (N^x)')$ and (f_1^x, f_2^x, N^x) are equivalent over M_0 , hence we can assume without loss of generality that the renaming has already been done and $f_2^x = \text{id}_{M_2}$.

Thus we know that $f_1^x(a) \downarrow_{M_0}^{N^x} M_2$ for $x = a, b$. By domination, $f_1^x[M_1] \downarrow_{M_0}^{N^x} M_2$.

Let \bar{M}_1 be an enumeration of M_1 . Using amalgamation, we can obtain the following diagram:

$$\begin{array}{ccc}
 N^a & \xrightarrow{g^a} & N' \\
 f_1^a \uparrow & & \uparrow g^b \\
 M_1 & \xrightarrow{f_1^b} & N^b
 \end{array}$$

This shows $\text{gtp}(f_1^a(\bar{M}_1)/M_0; N^a) = \text{gtp}(f_1^b(\bar{M}_1)/M_0; N^b)$. By uniqueness, $\text{gtp}(f_1^a(\bar{M}_1)/M_2; N^a) = \text{gtp}(f_1^b(\bar{M}_1)/M_2; N^b)$. Let N_* and $f^x : N^x \rightarrow N_*$ witness the equality. Since $f_2^x = \text{id}_{M_2}$, $f^b \circ f_2^b = f^b \upharpoonright M_2 = \text{id}_{M_2} = f^a \circ f_2^a$. Moreover, $(f^b \circ f_1^b)(\bar{M}_1) = f^b(f_1^b(\bar{M}_1)) = f^a(f_1^a(\bar{M}_1))$ by definition, so $f^b \circ f_1^b = f^a \circ f_1^a$. This completes the proof. \square

Remark 11.8. The converse holds if \mathbf{i} has left extension.

Remark 11.9. The relationship of uniqueness triples with domination is already mentioned in [JS13, Proposition 4.1.7], although the definition of domination there is different.

Thus to prove the existence property for uniqueness triples, it will be enough to imitate the proof of [MS90, Proposition 4.22], which gives conditions under which the hypothesis of Lemma 11.7 holds. We first show that we can work inside a local monster model.

Lemma 11.10. Let $M \leq N$ and $B \subseteq |N|$. Let $\mathfrak{C} \geq N$ be $\|N\|^{+}$ -saturated. Then B model-dominates N over M in \mathfrak{C} if and only if for any $M' \leq \mathfrak{C}$ with $M \leq M'$, $B \underset{M}{\downarrow}^{\mathfrak{C}} M'$ implies $N \underset{M}{\downarrow}^{\mathfrak{C}} M'$. Moreover if \mathbf{i} has the right $(\leq \mu)$ -witness property, we get an analogous result for domination instead of model-domination.

Proof. We prove the non-trivial direction for model-domination. The proof of the moreover part for domination is similar. Assume $\mathfrak{C}' \geq \mathfrak{C}$ and $M \leq M' \leq \mathfrak{C}'$ is such that $B \underset{M}{\downarrow}^{\mathfrak{C}'} M'$. We want to show that

$N \underset{M}{\downarrow}^{\mathfrak{C}'} M'$. Suppose not. Then we can use the $(\leq \mu)$ -model-witness property to assume without loss of generality that $\|M'\| \leq \mu + \|M\|$, and so we can find $N \leq N' \leq \mathfrak{C}'$ containing M' with $\|N'\| = \|N\|$ and $B \underset{M}{\downarrow}^{N'} M'$, $N \not\underset{M}{\downarrow}^{N'} M'$. By homogeneity, find $f : N' \rightarrow \mathfrak{C}$. By invariance, $B \underset{M}{\downarrow}^{f[N']} f[M']$ but $N \not\underset{M}{\downarrow}^{f[N']} f[M']$. By monotonicity, $B \underset{M}{\downarrow}^{\mathfrak{C}} f[M']$ but $N \not\underset{M}{\downarrow}^{\mathfrak{C}} f[M']$, a contradiction. \square

Lemma 11.11 (Lemma 4.20 in [MS90]). Let $\langle M_i : i < \lambda^+ \rangle, \langle N_i : i < \lambda^+ \rangle$ be increasing in K_λ such that $M_i \leq N_i$ for all $i < \lambda^+$. Let $M_{\lambda^+} := \bigcup_{i < \lambda^+} M_i, N_{\lambda^+} := \bigcup_{i < \lambda^+} N_i$.

Then there exists $i < \lambda^+$ such that $N_i \underset{M_i}{\downarrow}^{N_{\lambda^+}} M_{\lambda^+}$.

Proof. For each $i < \lambda^+$, let $j_i < \lambda^+$ be least such that $N_i \underset{M_{j_i}}{\downarrow}^{N_{\lambda^+}} M_{\lambda^+}$ (exists since $\kappa_\lambda(\mathbf{i}) = \lambda^+$). Let i^* be such that $j_i < i^*$ for all $i < i^*$ and

$\text{cf}(i^*) \geq \mu^+$. By definition of j_i and base monotonicity we have that for all $i < i^*$, $N_i \underset{M_{i^*}}{\overset{N_{\lambda^+}}{\downarrow}} M_{\lambda^+}$. By the left $(\leq \mu)$ -model-witness property,

$$N_{i^*} \underset{M_{i^*}}{\overset{N_{\lambda^+}}{\downarrow}} M_{\lambda^+}. \quad \square$$

Lemma 11.12 (Proposition 4.22 in [MS90]). Let $M \in K_\lambda$ be saturated. Let $\mathfrak{C} \geq M$ be saturated of size λ^+ . Work inside \mathfrak{C} . Write $A \underset{M}{\downarrow} B$ for $A \underset{M}{\overset{\mathfrak{C}}{\downarrow}} B$.

- There exists a saturated $N \leq \mathfrak{C}$ in K_λ such that $M \leq N$, N contains a , and a model-dominates N over M (in \mathfrak{C}).
- In fact, if $M^* \leq M$ is in $K_{<\lambda}$, $a \underset{M^*}{\downarrow} M$, and $r \in \text{gS}^{\leq \lambda}(M^*a)$, then N can be chosen so that it realizes r .

Proof. Since $\bar{\kappa}_1(\mathfrak{s}) = \mu^+ \leq \lambda$, it suffices to prove the second part. Assume it fails.

Claim: For any saturated $M' \geq M$ in K_λ , if $a \underset{M}{\downarrow} M'$, then the second part fails with M' replacing M .

Proof of claim: By transitivity, $a \underset{M^*}{\downarrow} M'$. By uniqueness of saturated models, there exists $f : M' \cong_{M^*} M$, which we can extend to an automorphism of \mathfrak{C} . Thus we also have $f(a) \underset{M^*}{\downarrow} M$. By uniqueness, we can assume without loss of generality that f fixes a as well. Since the second part above is invariant under applying f^{-1} , the result follows.

We now construct increasing continuous chains $\langle M_i : i \leq \lambda^+ \rangle$, $\langle N_i : i \leq \lambda^+ \rangle$ such that for all $i < \lambda^+$:

- (1) $M_0 = M$.
- (2) $M_i \leq N_i$.
- (3) $M_i \in K_\lambda$ is saturated.
- (4) $a \underset{M_0}{\downarrow} M_i$.
- (5) $N_i \not\underset{M_i}{\downarrow} M_{i+1}$.

This is enough: the sequences contradict Lemma 11.11. This is possible: take $M_0 = M$, and N_0 any saturated model of size λ containing M_0 and a and realizing r . At limits, take unions (we are using that $K^{\lambda\text{-sat}}$ is an AEC). Now assume everything up to i has been constructed. By the

claim, the second part above fails for M_i , so in particular N_i cannot be model-dominated by a over M_i . Thus (implicitly using Lemma 11.10) there exists $M'_i \geq M_i$ with $a \perp_{M_i} M'_i$ and $N_i \not\perp_{M_i} M'_i$. By the model-witness property, we can assume without loss of generality that $\|M'_i\| \leq \lambda$, so using extension and transitivity, we can find $M_{i+1} \in K_\lambda$ saturated containing M'_i so that $a \perp_{M_i} M_{i+1}$. By monotonicity we still have $N_i \not\perp_{M_i} M_{i+1}$. Let $N_{i+1} \in K_\lambda$ be any saturated model containing N_i and M_{i+1} . \square

Theorem 11.13. $\mathfrak{s}_\lambda \upharpoonright K_\lambda^{\lambda\text{-sat}}$ is a *weakly successful* type-full good λ -frame.

Proof. Since \mathfrak{s}_λ is a type-full good frame, $\mathfrak{s}_\lambda \upharpoonright K_\lambda^{\lambda\text{-sat}}$ also is. To show it is weakly successful, we want to prove the existence property for uniqueness triples. So let $M \in K_\lambda^{\lambda\text{-sat}}$ and $p \in \text{gS}(M)$ be nonalgebraic. Say $p = \text{gtp}(a/M; N')$. Let \mathfrak{C} be a monster model with $N' \leq \mathfrak{C}$. By Lemma 11.12, there exists $N \leq \mathfrak{C}$ in $K_\lambda^{\lambda\text{-sat}}$ such that $M \leq N$, $a \in |N|$, and a dominates N over M in \mathfrak{C} . By Lemma 11.6, a dominates N over M in N . By Lemma 11.7, $(a, M, N) \in K_{\mathfrak{s}_\lambda \upharpoonright K_\lambda^{\lambda\text{-sat}}}^{3, \text{uq}}$. Now, $p = \text{gtp}(a/M; N') = \text{gtp}(a/M; \mathfrak{C}) = \text{gtp}(a/M; N)$, as desired. \square

The term “weakly successful” suggests that there must exist a definition of “successful”. This is indeed the case:

Definition 11.14 (Definition 10.1.1 in [JS13]). A type-full good λ_t -frame \mathfrak{t} is *successful* if it is weakly successful and $\leq_{\lambda_t^+}^{\text{NF}}$ has smoothness: whenever $\langle N_i : i \leq \delta \rangle$ is a $\leq_{\lambda_t^+}^{\text{NF}}$ -increasing continuous chain of saturated models in $(K_t^{\text{up}})_{\lambda_t^+}$, $N \in (K_t^{\text{up}})_{\lambda_t^+}$ is saturated and $i < \delta$ implies $N_i \leq_{\lambda_t^+}^{\text{NF}} N$, then $N_\delta \leq_{\lambda_t^+}^{\text{NF}} N$.

We will not define $\leq_{\lambda_t^+}^{\text{NF}}$ (the interested reader can consult e.g. [JS13, Definition 6.14]). The only fact about it we will need is:

Fact 11.15 (Theorem 4.1 in [Jarb]). If \mathfrak{t} is a weakly successful type-full good λ_t -frame, $(K_t^{\text{up}})_{[\lambda_t, \lambda_t^+]}$ has amalgamation and is λ_t -tame, then $\leq \upharpoonright (K_t^{\text{up}})_{\lambda_t^+}^{\lambda_t^+ \text{-sat}} = \leq_{\lambda_t^+}^{\text{NF}}$.

Corollary 11.16. $\mathfrak{s}_\lambda \upharpoonright K_\lambda^{\lambda\text{-sat}}$ is a *successful* type-full good λ -frame.

Proof. By Theorem 11.13, $\mathfrak{s}_\lambda \upharpoonright K_\lambda^{\lambda\text{-sat}}$ is weakly successful. To show it is successful, it is enough (by Fact 11.15), to see that \leq has smoothness. But this holds since K is an AEC. \square

For a good λ_t -frame \mathfrak{t} , Shelah also defines a λ_t^+ -frame \mathfrak{t}^+ ([She09, Definition III.1.7]). He then goes on to show:

Fact 11.17 (Claim III.1.9 in [She09]). If \mathfrak{t} is a successful good λ_t -frame, then \mathfrak{t}^+ is a good³¹ λ_t^+ -frame.

Remark 11.18. This does *not* use the weak continuum hypothesis.

Note that in our case, it is easy to check that:

Fact 11.19. $(\mathfrak{s}_\lambda)^+ = \mathfrak{s}_{\lambda^+} \upharpoonright K_{\lambda^+}^{\lambda^+-\text{sat}}$.

Definition 11.20 (Definition III.1.12 in [She09]). Let \mathfrak{t} be a pre- λ_t -frame.

- (1) By induction on $n < \omega$, define \mathfrak{t}^{+n} as follows:
 - (a) $\mathfrak{t}^{+0} = \mathfrak{t}$.
 - (b) $\mathfrak{t}^{+(n+1)} = (\mathfrak{t}^{+n})^+$.
- (2) By induction on $n < \omega$, define “ \mathfrak{t} is n -successful” as follows:
 - (a) \mathfrak{t} is 0-successful if and only if it is a good λ -frame.
 - (b) \mathfrak{t} is $(n+1)$ -successful if and only if it is a successful good λ -frame and \mathfrak{t}^+ is n -successful.
- (3) \mathfrak{t} is ω -successful if it is n -successful for all $n < \omega$.

Thus by Fact 11.17, \mathfrak{t} is 1-successful if and only if it is a successful good λ_t -frame. More generally a good λ_t -frame \mathfrak{t} is n -successful if and only if \mathfrak{t}^{+m} is a successful good λ_t^{+m} -frame for all $m < n$.

Theorem 11.21. $\mathfrak{s}_\lambda \upharpoonright K_\lambda^{\lambda\text{-sat}}$ is an ω -successful type-full good λ -frame.

Proof. By induction on $n < \omega$, simply observing that we can replace λ by λ^{+n} in Corollary 11.16. \square

We emphasize again that we did *not* use the weak continuum hypothesis (as Shelah does in [She09, Chapter II]). We pay for this by using tameness (in Fact 11.3). Note that all the results of [She09, Chapter III] apply to our ω -successful good frame.

Recall that part of Shelah’s point is that ω -successful good λ -frames extend to $(\geq \lambda)$ -frames. However this is secondary for us (since tameness already implies that a frame extends to larger models, see [Bon14a, BVb]). Really, we want to extend the good frame to longer *types*. We show it is possible in the next section.

³¹Shelah proves that \mathfrak{t}^+ is actually good⁺. There is no reason to define what this means here.

12. A FULLY GOOD LONG FRAME

Hypothesis 12.1. $\mathfrak{s} = (K, \perp)$ is a weakly successful type-full good λ -frame.

This is reasonable since the previous section showed us how to build such a frame. Our goal is to extend \mathfrak{s} to obtain a fully good $(\leq \lambda, \lambda)$ -independence relation.

Fact 12.2 (Conclusion II.6.34 in [She09]). There exists a relation $\text{NF} \subseteq {}^4K$ satisfying:

- (1) $\text{NF}(M_0, M_1, M_2, M_3)$ implies $M_0 \leq M_\ell \leq M_3$ are in K for $\ell = 1, 2$.
- (2) $\text{NF}(M_0, M_1, M_2, M_3)$ and $a \in |M_1| \setminus |M_2|$ implies $\text{gtp}(a/M_2; M_3)$ does not \mathfrak{s} -fork over M_0 .
- (3) Invariance: NF is preserved under isomorphisms.
- (4) Monotonicity: If $\text{NF}(M_0, M_1, M_2, M_3)$:
 - (a) If $M_0 \leq M'_\ell \leq M_\ell$ for $\ell = 1, 2$, then $\text{NF}(M_0, M'_1, M'_2, M_3)$.
 - (b) If $M'_3 \leq M_3$ contains $|M_1| \cup |M_2|$, then $\text{NF}(M_0, M_1, M_2, M'_3)$.
 - (c) If $M'_3 \geq M_3$, then $\text{NF}(M_0, M_1, M_2, M'_3)$.
- (5) Symmetry: $\text{NF}(M_0, M_1, M_2, M_3)$ if and only if $\text{NF}(M_0, M_2, M_1, M_3)$.
- (6) Long transitivity: If $\langle M_i : i \leq \alpha \rangle, \langle N_i : i \leq \alpha \rangle$ are increasing continuous and $\text{NF}(M_i, N_i, M_{i+1}, N_{i+1})$ for all $i < \alpha$, then $\text{NF}(M_0, N_0, M_\alpha, N_\alpha)$.
- (7) Independent amalgamation: If $M_0 \leq M_\ell$, $\ell = 1, 2$, then for some $M_3 \in K$, $f_\ell : M_\ell \xrightarrow{M_0} M_3$, we have $\text{NF}(M_0, f_1[M_1], f_2[M_2], M_3)$.
- (8) Uniqueness: If $\text{NF}(M_0^\ell, M_1^\ell, M_2^\ell, M_3^\ell)$, $\ell = 1, 2$, $f_i : M_i^1 \cong M_i^2$ for $i = 0, 1, 2$, and $f_0 \subseteq f_1$, $f_0 \subseteq f_2$, then $f_1 \cup f_2$ can be extended to $f_3 : M_3^1 \rightarrow M_4^2$, for some M_4^2 with $M_3^2 \leq M_4^2$.

Notation 12.3. We write $M_1 \underset{M_0}{\overset{M_3}{\perp}} M_2$ instead of $\text{NF}(M_0, M_1, M_2, M_3)$.

If \bar{a} is a sequence, we write $\bar{a} \underset{M_0}{\overset{M_3}{\perp}} M_2$ for $\text{ran}(\bar{a}) \underset{M_0}{\overset{M_3}{\perp}} M_2$, and similarly if sequences appear at other places.

Remark 12.4. Shelah's definition of NF ([She09, Definition II.6.12]) is very complicated. It is somewhat simplified in [JS13].

Remark 12.5. Shelah calls such an NF a *nonforking relation which respects \mathfrak{s}* ([She09, Definition II.6.1]). While there are similarities with this paper's definition of a good $(\leq \lambda)$ -frame, note that NF is only defined for types of models while we would like to make it into a relation for arbitrary types of length at most λ .

We start by showing that uniqueness is really the same as the uniqueness property stated for frames. We drop Hypothesis 12.1 for the next lemma.

Lemma 12.6. Let K be an AEC in λ and assume K has amalgamation. The following are equivalent for a relation $\text{NF} \subseteq {}^4K$ satisfying (1), (3), (4) of Fact 12.2:

- (1) Uniqueness in the sense of Fact 12.2.(8).
- (2) Uniqueness in the sense of frames: If $A \underset{M_0}{\downarrow}^N M_1$ and $A' \underset{M_0}{\downarrow}^{N'} M_1$ for models A and A' , \bar{a} and \bar{a}' are enumerations of A and A' respectively, $p := \text{gtp}(\bar{a}/M_1; N)$, $q := \text{gtp}(\bar{a}'/M_1; N')$, and $p \upharpoonright M_0 = q \upharpoonright M_0$, then $p = q$.

Proof.

- (1) implies (2): Since $p \upharpoonright M_0 = q \upharpoonright M_0$, there exists $N'' \geq N'$ and $f : N \xrightarrow{M_0} N''$ such that $f(\bar{a}) = \bar{a}'$. Therefore by invariance, $\bar{a}' \underset{M_0}{\downarrow}^{N''} f[M_1]$. Let $f_0 := \text{id}_{M_0}$, $f_1 := f^{-1} \upharpoonright f[M_1]$, $f_2 := \text{id}_{A'}$. By uniqueness, there exists $N''' \geq N''$, $g \supseteq f_1 \cup f_2$, $g : N'' \rightarrow N'''$. Consider the map $h := g \circ f : N \rightarrow N'''$. Then $g \upharpoonright M_1 = \text{id}_{M_1}$ and $h(\bar{a}) = g(\bar{a}') = \bar{a}'$, so h witnesses $p = q$.
- (2) implies (1): By some renaming, it is enough to prove that whenever $M_2 \underset{M_0}{\downarrow}^N M_1$ and $M_2 \underset{M_0}{\downarrow}^{N'} M_1$, there exists $N'' \geq N'$ and $f : N' \xrightarrow{|M_1| \cup |M_2|} N''$. Let \bar{a} be an enumeration of M_2 . Let $p := \text{gtp}(\bar{a}/M_1; N)$, $q := \text{gtp}(\bar{a}/M_1; N')$. We have that $p \upharpoonright M_0 = \text{gtp}(\bar{a}/M_1; M_2) = q \upharpoonright M_0$. Thus $p = q$, so there exists $N'' \geq N'$ and $f : N \xrightarrow{M_1} N''$ such that $f(\bar{a}) = \bar{a}$. In other words, f fixes M_2 , so is the desired map.

□

We now extend NF to take sets on the left hand side. This step is already made by Shelah in [She09, Claim III.9.6], for singletons rather than arbitrary sets. We check that Shelah's proofs still work.

Definition 12.7. Define $\text{NF}'(M_0, A, M, N)$ to hold if and only if $M_0 \leq M \leq N$ are in K , $A \subseteq |N|$, and there exists $N' \geq N$, $N_A \geq M$ with

$N_A \leq N'$ and $N_A \underset{M_0}{\downarrow}^{N'} M$. We abuse notation and also write $A \underset{M_0}{\downarrow}^N M$ instead of $\text{NF}'(M_0, A, M, N)$. We let $\mathfrak{t} := (K, \downarrow)$.

Remark 12.8. Compare with the definition of cl (Definition 3.8).

Proposition 12.9.

- (1) If $M_0 \leq M_\ell \leq M_3$, $\ell = 1, 2$, then $\text{NF}(M_0, M_1, M_2, M_3)$ if and only if $\text{NF}'(M_0, M_1, M_2, M_3)$.
- (2) \mathfrak{t} is a (type-full) pre- $(\leq \lambda, \lambda, \lambda)$ -frame.
- (3) \mathfrak{t} has base monotonicity, full symmetry, uniqueness, existence, and extension.

Proof. Exactly as in [She09, Claim III.9.6]. \square

We now turn to local character. The key is:

Fact 12.10 (Claim III.1.17.2 in [She09]). Given $\langle M_i : i \leq \delta \rangle$ increasing continuous, we can build $\langle N_i : i \leq \delta \rangle$ increasing continuous such that

for all $i \leq j \leq \delta$, $N_i \underset{M_i}{\downarrow}^{N_j} M_j$ and $M_\delta <_{\text{univ}} N_\delta$.

Lemma 12.11. For all $\alpha \leq \lambda$, $\kappa_\alpha(\mathfrak{t}) = |\alpha|^+ + \aleph_0$.

Proof. Let $\langle M_i : i \leq \delta + 1 \rangle$ be increasing continuous with $\delta = \text{cf}(\delta) > |\alpha|$. Let $A \subseteq |M_{\delta+1}|$ have size $\leq \alpha$. Let $\langle N_i : i \leq \delta \rangle$ be as given by Fact 12.10. By universality, we can assume without loss of generality that $M_{\delta+1} \leq N_\delta$. Thus $A \subseteq |N_\delta|$ and by the cofinality hypothesis, there exists $i < \delta$ such that $A \subseteq |N_i|$. In particular, $A \underset{M_i}{\downarrow}^{N_\delta} M_\delta$, so $A \underset{M_i}{\downarrow}^{M_{\delta+1}} M_\delta$, as needed. \square

Remark 12.12. In [JS13] (and later in [JS12, Jara, Jarb]), the authors have considered *semi-good* λ -frames, where the stability condition is replaced by almost stability ($|\text{gS}(M)| \leq \lambda^+$ for all $M \in K_\lambda$), and an hypothesis called the conjugation property is often added. Many of the above results carry through in that setup but we do not know if Lemma 12.11 would also hold.

We come to the last property: disjointness. The situation is a bit murky: At first glance, Fact 12.2.(2) seems to give it to us for free (since we are assuming \mathfrak{s} has disjointness), but unfortunately we are assuming $a \notin |M_2|$ there. We will obtain it with the additional hypothesis of categoricity in λ (this is reasonable since if the frame has

a superlimit, see Remark 10.20, one can always restrict oneself to the class generated by the superlimit). Note that disjointness is never used in a crucial way in this paper (but it is always nice to have, as it implies for example disjoint amalgamation when combined with independent amalgamation).

Lemma 12.13. If K is categorical in λ , then \mathfrak{t} has disjointness and $\mathfrak{t}^{\leq 1} = \mathfrak{s}$.

Proof. We have shown that $\mathfrak{t}^{\leq 1}$ has all the properties of a good frame except perhaps disjointness so by the proof of Theorem 9.7 (which never relied on disjointness), $\mathfrak{s} = \mathfrak{t}^{\leq 1}$. Since \mathfrak{s} has disjointness, $\mathfrak{t}^{\leq 1}$ also does, and therefore \mathfrak{t} has disjointness. \square

What about continuity for chains? The long transitivity property seems to suggest we can say something, and indeed we can:

Fact 12.14. Assume $\lambda = \lambda_0^{+3}$ and there exists an ω -successful good λ_0 -frame \mathfrak{s}' such that $\mathfrak{s} = (\mathfrak{s}')^{+3}$.

Assume δ is a limit ordinal and $\langle M_i^\ell : i \leq \delta \rangle$ is increasing continuous in K_λ , $\ell \leq 3$. If $M_i^1 \downarrow_{M_i^0}^{M_i^3} M_i^2$ for each $i < \delta$, then $M_\delta^1 \downarrow_{M_\delta^0}^{M_\delta^3} M_\delta^2$.

Proof. By [She09, Claim III.12.2], all the hypotheses at the beginning of each section of Chapter III in the book hold for \mathfrak{s} . Now apply Claim III.8.19 in the book. \square

Remark 12.15. Where does the hypothesis $\lambda = \lambda_0^{+3}$ come from? Shelah's analysis in chapter III of his book proceeds on the following lines: starting with an ω -successful frames \mathfrak{s} , we want to show \mathfrak{s} has nice properties like existence of prime triples, weak orthogonality being orthogonality, etc. They are hard to show in general, however it turns out \mathfrak{s}^+ has some nicer properties than \mathfrak{s} (for example, $K_{\mathfrak{s}^+}$ is always categorical)... In general, $\mathfrak{s}^{+(n+1)}$ has even nicer properties than \mathfrak{s}^{+n} ; and Shelah shows that the frame has all the nice properties he wants after going up three successors.

We obtain:

Theorem 12.16.

- (1) If K is categorical in λ , then \mathfrak{t} is a good $(\leq \lambda, \lambda)$ -frame.
- (2) If $\lambda = \lambda_0^{+3}$ and there exists an ω -successful good λ_0 -frame \mathfrak{s}' such that $\mathfrak{s} = (\mathfrak{s}')^{+3}$, then \mathfrak{t} is a fully good $(\leq \lambda, \lambda)$ -frame.

Proof. \mathfrak{t} is good by Proposition 12.9, Lemma 12.11, and Lemma 12.13. The second part follows from Fact 12.14 (note that by definition of the successor frame, K will be categorical in λ in that case). \square

Remark 12.17. In [BVb, Corollary 6.10], it is shown that λ -tameness and amalgamation imply that a good λ -frame extends to a good $(< \infty, \lambda)$ -frame. However, the definition of a good frame there is not the same as it does *not* assume that the frame is type-full. Thus the conclusion of Theorem 12.16 is much stronger.

13. EXTENDING THE BASE AND RIGHT HAND SIDE

Hypothesis 13.1.

- (1) $\mathfrak{i} = (K, \perp)$ is a fully good $(\leq \lambda, \lambda)$ -independence relation.
- (2) $K' := K^{\text{up}}$ has amalgamation and is λ -tame for types of length less than λ^+ .

In this section, we give conditions under which \mathfrak{i} becomes a fully good $(\leq \lambda, \geq \lambda)$ -independence relation. In the next section, we will make the left hand side bigger and get a fully good $(< \infty, \geq \lambda)$ -independence relation.

Recall that extending a $(\leq 1, \lambda)$ -frame to bigger models was investigated in [She09, Chapter II] and [Bon14a, BVb]. Here, most of the arguments are similar but the longer types cause some additional difficulties (e.g. in the proof of local character).

Notation 13.2. Let $\mathfrak{i}' := \mathfrak{i}^{\text{up}}$ (recall Definition 6.3). Write $\mathfrak{s} := \text{pre}(\mathfrak{i})$, $\mathfrak{s}' := \text{pre}(\mathfrak{i}')$. We abuse notation and also denote $\perp_{\mathfrak{i}'}$ by \perp .

We want to investigate when the properties of \mathfrak{i} carry over to \mathfrak{i}' .

Lemma 13.3.

- (1) \mathfrak{i}' is a $(< \infty, \geq \lambda)$ -independence relation.
- (2) K' has joint embedding, no maximal models, and is stable in all cardinals.
- (3) \mathfrak{i}' has base monotonicity, transitivity, uniqueness, and disjointness.
- (4) \mathfrak{i}' has full model continuity.

Proof.

- (1) By Proposition 6.5.

- (2) By [BVb, Theorem 1.1], $(\mathfrak{s}')^{\leq 1}$ is a good frame, so in particular K' has joint embedding, no maximal models, and is stable in all cardinals.
- (3) See [She09, Claim II.2.11] for base monotonicity and transitivity. Disjointness is straightforward from the definition of \mathfrak{i}' , and uniqueness follows from the tameness hypothesis and the definition of \mathfrak{i}' .
- (4) Assume $\langle M_i^\ell : i \leq \delta \rangle$ is increasing continuous in K' , $\ell \leq 3$, δ is regular, $M_i^0 \leq M_i^\ell \leq M_i^3$ for $\ell = 1, 2$, $i < \delta$, and $M_i^1 \underset{M_i^0}{\overset{M_i^3}{\downarrow}} M_i^2$ for all $i < \delta$. Let $N := M_\delta^3$. By ambient monotonicity, $M_i^1 \underset{M_i^0}{\overset{N}{\downarrow}} M_i^2$ for all $i < \delta$. We want to see that $M_\delta^1 \underset{M_\delta^0}{\overset{N}{\downarrow}} M_\delta^2$. Since $\|M_\delta^1\| < \theta$, M_δ^1 and M_δ^0 are in K . Thus it is enough to show that for all $M' \leq M_\delta^2$ in K with $M_\delta^0 \leq M'$, $M_\delta^1 \underset{M_\delta^0}{\overset{N}{\downarrow}} M'$. Fix such an M' . We consider two cases:

- Case 1: $\delta < \theta$: Then we can find $\langle M'_i : i \leq \delta \rangle$ increasing continuous in K such that $M'_\delta = M'$ and for all $i < \delta$, $M_\delta^0 \leq M'_i \leq M_i^2$. By monotonicity, for all $i < \delta$, $M_i^1 \underset{M_i^0}{\overset{N}{\downarrow}} M'_i$.

By full model continuity in K , $M_\delta^1 \underset{M_\delta^0}{\overset{N}{\downarrow}} M'$, as desired.

- Case 2: $\delta \geq \theta$: Since $M_\delta^0, M_\delta^1 \in K$, we can assume without loss of generality that $M_\delta^0 = M_0^0$, $M_\delta^1 = M_\delta^1$. Since δ is regular, there exists $i < \delta$ such that $M' \leq M_i^2$. By assumption, $M_0^1 \underset{M_0^0}{\overset{N}{\downarrow}} M_i^2$, so by monotonicity, $M_0^1 \underset{M_0^0}{\overset{N}{\downarrow}} M'$, as needed.

□

We now turn to local character.

Lemma 13.4. Assume $\langle M_i : i \leq \delta \rangle$ is increasing continuous, $p \in \text{gS}^\alpha(M_\delta)$, $\alpha < \lambda^+$ a cardinal and $\delta = \text{cf}(\delta) > \alpha$.

- (1) If $\alpha < \lambda$, then there exists $i < \delta$ such that p does not fork over M_i .

- (2) If $\alpha = \lambda$ and \mathbf{i} has the left ($< \text{cf}(\lambda)$)-witness property, then there exists $i < \delta$ such that p does not fork over M_i .

Proof.

- (1) As in the proof of Lemma 6.8.(2b) Note that weak chain local character holds for free because $\alpha < \lambda$ and $\kappa_\alpha(\mathbf{i}) = \alpha^+ + \aleph_0$ by assumption.
- (2) By the proof of Lemma 6.8.(2b) again, it is enough to see that \mathbf{i} has weak chain local character: Let $\langle M_i : i < \lambda^+ \rangle$ be increasing in K and let $M_{\lambda^+} := \bigcup_{i < \lambda^+} M_i$. Let $p \in \text{gS}^\lambda(M_{\lambda^+})$. We will show that there exists $i < \lambda^+$ such that p does not fork over M_i . Say $p = \text{gtp}(\bar{a}/M_{\lambda^+}; N)$ and let $A := \text{ran}(\bar{a})$. Write $A = \bigcup_{j < \text{cf}(\lambda)} A_j$ with $\langle A_j : j < \text{cf}(\lambda) \rangle$ increasing continuous and $|A_j| < \lambda$. By the first part for each $j < \text{cf}(\lambda)$ there exists $i_j < \lambda^+$ such that $A_j \downarrow_{M_{i_j}}^N M_{\lambda^+}$. Let $i := \sup_{j < \text{cf}(\lambda)} i_j$. We claim that $A \downarrow_{M_i}^N M_{\lambda^+}$. By the ($< \text{cf}(\lambda)$)-witness property and the definition of \mathbf{i}' (here we use that $M_i \in K$), it is enough to show this for all $B \subseteq A$ of size less than $\text{cf}(\lambda)$. But any such B is contained in an A_j , and so the result follows from base monotonicity.

□

Lemma 13.5. Assume \mathbf{i}' has existence. Then \mathbf{i}' has independent amalgamation.

Proof. As in, for example, [Bon14a, Theorem 5.3], using full model continuity. □

Putting everything together, we obtain:

Theorem 13.6. If K is ($< \text{cf}(\lambda)$)-tame and short for types of length less than λ^+ , then \mathbf{i}' is a fully pre-good ($\leq \lambda, \geq \lambda$)-independence relation.

Proof. We want to show that \mathbf{s}' is fully good. The basic properties are proven in Lemma 13.3. By Lemma 4.5, \mathbf{i} has the left ($< \text{cf}(\lambda)$)-witness property. Thus by Lemma 13.4, for any $\alpha < \lambda^+$, $\kappa_\alpha(\mathbf{i}') = |\alpha|^+ + \aleph_0$. In particular, \mathbf{i}' has existence, and thus by the definition of \mathbf{i}' and transitivity in \mathbf{i} , $\bar{\kappa}_\alpha(\mathbf{i}') = \lambda^+ = |\alpha|^+ + \lambda^+$. Finally by Lemma 13.5,

\mathfrak{i}' has independent amalgamation and so by Proposition 4.3.(3), \mathfrak{i}' has extension. \square

14. EXTENDING THE LEFT HAND SIDE

We now enlarge the left hand side of the independence relation built in the previous section.

Hypothesis 14.1.

- (1) $\mathfrak{i} = (K, \perp)$ is a fully good $(\leq \lambda, \geq \lambda)$ -independence relation.
- (2) K is fully λ -tame and short.

Definition 14.2. Define $\mathfrak{i}^{\text{long}} = (K, \perp^{\text{long}})$ by setting $\perp^{\text{long}}(M_0, A, B, N)$ if and only if for all $A_0 \subseteq A$ of size less than λ^+ , $A_0 \underset{M_0}{\overset{N}{\perp}} B$.

Remark 14.3. The idea is the same as for [BVb, Definition 4.3]: we extend the frame to have longer types. The difference is that $\mathfrak{i}^{\text{long}}$ is type-full.

Remark 14.4. We could also have defined extension to types of length less than θ for θ a cardinal or ∞ but this complicates the notation and we have no use for it here.

Notation 14.5. Write $\mathfrak{i}' := \mathfrak{i}^{\text{long}}$. We abuse notation and also write \perp for \perp^{long} .

Lemma 14.6.

- (1) \mathfrak{i}' is a $(< \infty, \geq \lambda)$ -independence relation.
- (2) K has joint embedding, no maximal models, and is stable in all cardinals.
- (3) \mathfrak{i}' has base monotonicity, transitivity, disjointness, existence, symmetry, the left λ -witness property, and uniqueness.

Proof.

- (1) Straightforward.
- (2) Because \mathfrak{i} is good.
- (3) Base monotonicity, transitivity, disjointness, existence, and the left λ -witness property are straightforward. Now by Lemma 4.5, \mathfrak{i} has the right λ -witness property, and so symmetry follows easily. Uniqueness is by the shortness hypothesis.

\square

Lemma 14.7. Assume there exists a regular $\kappa \leq \lambda$ such that \mathbf{i} has the left ($< \kappa$)-model-witness property. Then \mathbf{i}' has full model continuity.

Proof. Let $\langle M_i^\ell : i \leq \delta \rangle$, $\ell \leq 3$ be increasing continuous in K such that $M_i^0 \leq M_i^\ell \leq M_i^3$, $\ell = 1, 2$, and $M_i^1 \underset{M_i^0}{\overset{M_i^3}{\downarrow}} M_i^2$. Without loss of generality,

δ is regular. Let $N := M_\delta^3$. We want to show that $M_\delta^1 \underset{M_\delta^0}{\overset{N}{\downarrow}} M_\delta^2$. Let

$A \subseteq |M_\delta^1|$ have size less than λ^+ . Write $\mu := |A|$. By monotonicity, assume without loss of generality that $\lambda + \kappa \leq \mu$. We show that $A \underset{M_\delta^0}{\overset{N}{\downarrow}} M_\delta^2$, which is enough by definition of \mathbf{i}' . We consider two cases.

- Case 1: $\delta > \mu$: By local character in \mathbf{i} there exists $i < \delta$ such that $A \underset{M_i^2}{\overset{N}{\downarrow}} M_\delta^2$. By right transitivity, $A \underset{M_i^0}{\overset{N}{\downarrow}} M_\delta^2$, so by base monotonicity, $A \underset{M_\delta^0}{\overset{N}{\downarrow}} M_\delta^2$.
- Case 2: $\delta \leq \mu$: For $i \leq \delta$, let $A_i := A \cap |M_i^1|$. Build $\langle N_i : i \leq \delta \rangle$, $\langle N_i^0 : i \leq \delta \rangle$ increasing continuous in $K_{\leq \mu}$ such that for all $i < \delta$:
 - (1) $A_i \subseteq |N_i|$.
 - (2) $N_i \leq M_i^1$, $A \subseteq |N_i|$.
 - (3) $N_i^0 \leq M_i^0$, $N_i^0 \leq N_i$.
 - (4) $N_i \underset{N_i^0}{\overset{N}{\downarrow}} M_i^2$.

This is possible. Fix $i \leq \delta$ and assume N_j, N_j^0 have already been constructed for $j < i$. If i is limit, take unions. Otherwise,

recall that we are assuming $M_i^1 \underset{M_i^0}{\overset{N}{\downarrow}} M_i^2$. By Lemma 4.7 (with

$A_i \cup \bigcup_{j < i} |N_j|$ standing for A there, this is where we use the ($< \kappa$)-model-witness property), we can find $N_i^0 \leq M_i^0$ and $N_i \leq M_i^1$ in $K_{\leq \mu}$ such that $N_i^0 \leq N_i$, $N_i \underset{N_i^0}{\overset{N}{\downarrow}} M_i^2$, $A_i \subseteq |N_i|$, $N_j \leq N_i$

for all $j < i$, and $N_j^0 \leq N_i^0$ for all $j < i$. Thus they are as desired.

This is enough. Note that $A_\delta = A$, so $A \subseteq |N_\delta|$. By full model continuity in \mathbf{i} , $N_\delta \underset{N_\delta^0}{\overset{N}{\downarrow}} M_\delta^2$. By monotonicity, $A \underset{M_\delta^0}{\overset{N}{\downarrow}} M_\delta^2$, as desired.

□

Lemma 14.8. Assume there exists a regular $\kappa \leq \lambda$ such that \mathfrak{i} has the left ($< \kappa$)-model-witness property. Then for all cardinals μ :

- (1) $\bar{\kappa}_\mu(\mathfrak{i}') = \lambda^+ + \mu^+$.
- (2) $\kappa_\mu(\mathfrak{i}') = \aleph_0 + \mu^+$.

Proof. By Lemma 14.7, \mathfrak{i}' has full model continuity. By Lemma 4.8, (1) holds. For (2), if $\mu \leq \lambda$, this holds because \mathfrak{i} is good and if $\mu > \lambda$, this follows from Proposition 4.3.(5) and (1). □

We now turn to proving extension. The proof is significantly more complicated than in the previous section. We attempt to explain why and how our proof goes. Of course, it suffices to show independent amalgamation (Proposition 4.3.(3)). We work by induction on the size of the models but land in trouble when all models have the same size. Suppose for example that we want to amalgamate $M^0 \leq M^\ell$, $\ell = 1, 2$ that are all in K_{λ^+} . If M^1 (or, by symmetry, M^2) had smaller size, we could use local character to assume without loss of generality that M^0 is in K_λ and then imitate the usual directed system argument (as in for example [Bon14a, Theorem 5.3]).

Here however it seems we have to take at least two resolutions at once so we fix $\langle M_i^\ell : i < \lambda^+ \rangle$, $\ell = 0, 1$, satisfying the usual conditions. Letting $p := \text{gtp}(M^1/M^0; M^1)$ and its resolution $p_i := \text{gtp}(M_i^1/M_i^0; M^1)$, it is natural to build $\langle q_i : i < \lambda^+ \rangle$ such that q_i is the nonforking extension of p_i to M^2 . If everything works, we can take the direct limit of the q_i s and get the desired nonforking extension of p . However with what we have said so far it is not clear that q_{i+1} is even an extension of q_i ! In the usual argument, this is the case since both p_i and p_{i+1} do not fork over the same domain but we cannot expect it here. Thus

we require in addition that $M_i^1 \underset{M_i^0}{\downarrow}^{M^1} M^0$ and this turns out to be enough

for successor steps. To achieve this extra requirement, we use Lemma 4.7. Unfortunately, we also do not know how to go through limit steps without making one extra locality hypothesis:

Definition 14.9 (Type-locality).

- (1) Let δ be a limit ordinal, and let $\bar{p} := \langle p_i : i < \delta \rangle$ be an increasing chain of Galois types, where for $i < \delta$, $p_i \in \text{gS}^{\alpha_i}(M)$ and $\langle \alpha_i : i \leq \delta \rangle$ are increasing continuous. We say \bar{p} is *type-local* if whenever $p, q \in \text{gS}^{\alpha_\delta}(M)$ are such that $p^{\alpha_i} = q^{\alpha_i} = p_i$ for all $i < \delta$, then $p = q$.

- (2) We say K is *type-local* if every \bar{p} as above is type-local.
- (3) We say K is *densely type-local above λ* if for every $\lambda_0 > \lambda$, $M \in K_{\lambda_0}$, $p \in \text{gS}^{\lambda_0}(M)$, there exists $\langle N_i : i \leq \delta \rangle$ such that:
 - (a) $\delta = \text{cf}(\lambda_0)$.
 - (b) For all $i < \delta$, $N_i \in K_{<\lambda_0}$.
 - (c) $\langle N_i : i \leq \delta \rangle$ is increasing continuous.
 - (d) $N_\delta \geq M$ is in K_{λ_0} .
 - (e) Letting $q_i := \text{gtp}(N_i/M; N_\delta)$ (seen as a member of $\text{gS}^{\alpha_i}(M)$, where of course $\langle \alpha_i : i \leq \delta \rangle$ are increasing continuous), we have that q_δ extends p and $\langle q_j : j < i \rangle$ is type-local for all limit $i \leq \delta$.

We say K is *densely type-local* if it is densely type-local above λ for some λ .

Intuitively, type-locality is a “dual” to locality (see [Bal09, Definition 11.4]) in the same sense that type-shortness is a dual to tameness. We suspect that dense type-locality should hold in our context, see the discussion in Section 15 for more. The following lemma says that increasing the elements in the resolution of the type preserves type-locality.

Lemma 14.10. Let δ be a limit ordinal. Assume $\bar{p} := \langle p_i : i < \delta \rangle$ is an increasing chain of Galois types, $p_i \in \text{gS}^{\alpha_i}(M)$ and $\langle \alpha_i : i \leq \delta \rangle$ are increasing continuous. Assume \bar{p} is type-local and assume $p_\delta \in \text{gS}^{\alpha_\delta}(M)$ is such that $p^{\alpha_i} = p_i$ for all $i < \delta$. Say $p = \text{gtp}(\bar{a}_\delta/M; N)$ and let $\bar{a}_i := \bar{a}_\delta \upharpoonright \alpha_i$ (so $p_i = \text{gtp}(\bar{a}_i/M; N)$).

Assume $\langle \bar{b}_i : i \leq \delta \rangle$ are increasing continuous sequences such that $\bar{a}_\delta = \bar{b}_\delta$ and \bar{a}_i is an initial segment of \bar{b}_i for all $i < \delta$. Let $q_i := \text{gtp}(\bar{b}_i/M; N)$. Then $\bar{q} := \langle q_i : i < \delta \rangle$ is type-local.

Proof. Say \bar{b}_i is of type β_i . So $\langle \beta_i : i \leq \delta \rangle$ is increasing continuous and $\alpha_\delta = \beta_\delta$.

If $q \in \text{gS}^{\beta_\delta}(M)$ is such that $q^{\beta_i} = q_i$ for all $i < \delta$, then $q^{\alpha_i} = (q_i)^{\alpha_i} = p_i$ for all $i < \delta$ so by type-locality of \bar{p} , $p = q$, as desired. \square

Before proving Lemma 14.13, let us make precise what was meant above by “direct limit” of a chain of types. It is known that (under some set-theoretic hypotheses) there exists AECs where some chains of Galois types do not have an upper bound, see [BS08, Theorem 3.3]. However a *coherent* chain of types (see below) always has an upper bound. We adapt Definition 5.1 in [Bon14a] (which is implicit already in [She01, Claim 0.32.2] or [GV06a, Lemma 2.12]) to our purpose.

Definition 14.11. Let δ be an ordinal. An increasing chain of types $\langle p_i : i < \delta \rangle$ is said to be *coherent* if there exists a sequence $\langle (\bar{a}_i, M_i, N_i) : i < \delta \rangle$ and maps $f_{i,j} : N_i \rightarrow N_j$, $i \leq j < \delta$, so that for all $i \leq j \leq k < \delta$:

- (1) $f_{j,k} \circ f_{i,j} = f_{i,k}$.
- (2) $\text{gtp}(\bar{a}_i/M_i; N_i) = p_i$.
- (3) $\langle M_i : i < \delta \rangle$ and $\langle N_i : i < \delta \rangle$ are increasing.
- (4) $M_i \leq N_i$, $\bar{a}_i \in {}^{<\infty}N_i$.
- (5) $f_{i,j}$ fixes M_i .
- (6) $f_{i,j}(\bar{a}_i)$ is an initial segment of \bar{a}_j .

We call the sequence and maps above a *witnessing sequence* to the coherence of the p_i 's.

Given a witnessing sequence $\langle (\bar{a}_i, M_i, N_i) : i < \delta \rangle$ with maps $f_{i,j} : N_i \rightarrow N_j$, we can let N_δ be the direct limit of the system $\langle N_i, f_{i,j} : i \leq j < \delta \rangle$, $M_\delta := \bigcup_{i < \delta} M_i$, and $\bar{a}_\delta := \bigcup_{i < \delta} f_{i,\delta}(\bar{a}_i)$ (where $f_{i,\delta} : N_i \rightarrow N_\delta$ is the canonical embedding). Then $p := \text{gtp}(\bar{a}_\delta/M_\delta; N_\delta)$ extends each p_i . Note that p depends on the witness but we sometimes abuse language and talk about “the” direct limit (where really some witnessing sequence is fixed in the background).

Finally, note that full model continuity also applies to coherent sequences. More precisely:

Proposition 14.12. Assume \mathfrak{i} has full model continuity. Let $\langle (\bar{a}_i, M_i, N_i) : i < \delta \rangle$, $\langle f_{i,j} : N_i \rightarrow N_j, i \leq j < \delta \rangle$ be witnesses to the coherence of $p_i := \text{gtp}(\bar{a}_i/M_i; N_i)$. Assume that for each $i < \delta$, \bar{a}_i enumerates a model M'_i and that $\langle M_i^0 : i < \delta \rangle$ are increasing such that $M_i^0 \leq M_i$, $M_i^0 \leq M'_i$, and p_i does not fork over M_i^0 . Let p be the direct limit of the p_i s (according to the witnessing sequence). Then p does not fork over $M_\delta^0 := \bigcup_{i < \delta} M_i^0$.

Proof. Use full model continuity inside the direct limit. □

Lemma 14.13. Assume K is densely type-local above λ , and assume there exists a regular $\kappa \leq \lambda$ such that K is fully $(< \kappa)$ -tame. Then \mathfrak{i}' has extension.

Proof. By Lemmas 14.7 and 14.8, \mathfrak{i}' has full model continuity and the local character properties. Let $\lambda_0 \geq \lambda$ be a cardinal. We prove by induction on λ_0 that \mathfrak{i}' has extension for base models in K_{λ_0} . By Proposition 4.3.(3), it is enough to prove independent amalgamation.

Let $M^0 \leq M^\ell$, $\ell = 1, 2$ be in K with $\|M^0\| = \lambda_0$. We want to find $q \in \text{gS}^{\lambda_0}(M^2)$ a nonforking extension of $p := \text{gtp}(M^1/M^0; M^1)$. Let $\lambda_\ell := \|M^\ell\|$ for $\ell = 1, 2$.

Assume we know the result when $\lambda_0 = \lambda_1 = \lambda_2$. Then we can work by induction on (λ_1, λ_2) : if they are both λ_0 , the result holds by assumption. If not, we can assume by symmetry that $\lambda_1 \leq \lambda_2$, find an increasing continuous resolution of M^2 , $\langle M_i^2 \in K_{<\lambda_2} : i < \lambda_2 \rangle$ and do a directed system argument as in [Bon14a, Theorem 5.3] (using full model continuity and the induction hypothesis).

Now assume that $\lambda_0 = \lambda_1 = \lambda_2$. If $\lambda_0 = \lambda$, we get the result by extension in \mathbf{i} , so assume $\lambda_0 > \lambda$. Let $\delta := \text{cf}(\lambda_0)$. By dense type-locality, we can assume (extending M^1 if necessary) that there exists $\langle N_i : i \leq \delta \rangle$ an increasing continuous resolution of M^1 with $N_i \in K_{<\lambda_0}$ for $i < \delta$ so that $\langle \text{gtp}(N_j/M^0; M^1) : j < i \rangle$ is type-local for all limit $i \leq \delta$.

Step 1. Fix increasing continuous $\langle M_i^\ell : i \leq \delta \rangle$ for $\ell < 2$ such that for all $i < \delta$, $\ell < 2$:

- (1) $M^\ell = M_\delta^\ell$.
- (2) $M_i^\ell \in K_{<\lambda_0}$.
- (3) $N_i \leq M_i^1$.
- (4) $M_i^0 \leq M_i^1$.
- (5) $M_i^1 \underset{M_i^0}{\downarrow}^{M^1} M^0$.

This is possible by repeated applications of Lemma 4.7 (as in the proof of Lemma 14.7), starting with $M^1 \underset{M^0}{\downarrow}^{M^1} M^0$ which holds by existence.

Step 2. Fix enumerations of M_i^1 of order type α_i such that $\langle \alpha_i : i \leq \delta \rangle$ is increasing continuous, $\alpha_\delta = \lambda_0$ and $i < j$ implies that M_i' appears as the initial segment up to α_i of the enumeration of M_j' . For $i \leq \delta$, let $p_i := \text{gtp}(M_i^1/M_i^0; M^1)$ (seen as an element of $\text{gS}^{\alpha_i}(M_i^0)$). We want to find $q \in \text{gS}^{\lambda_0}(M^2)$ extending $p = p_\delta$ and not forking over M^0 . Note that since for all $j < \delta$, $N_j \leq M_j^1$, we have by Lemma 14.10 that $\langle \text{gtp}(M_j^1/M^0; M^1) : j < i \rangle$ is type-local for all limit $i \leq \delta$.

Build an increasing, coherent $\langle q_i : i \leq \delta \rangle$ such that for all $i \leq \delta$,


- (1) $q_i \in \text{gS}^{\alpha_i}(M^2)$.
- (2) $q_i \upharpoonright M_i^0 = p_i$.
- (3) q_i does not fork over M_i^0 .

This is enough: then q_δ is an extension of $p = p_\delta$ that does not fork over $M_\delta^0 = M^0$.

This is possible: We work by induction on $i \leq \delta$. While we do not make it explicit, the sequence witnessing the coherence is also built inductively in the natural way (see also [Bon14a, Proposition 5.2]): at base and successor steps, we use the definition of Galois types. At limit steps, we take direct limits.

Now fix $i \leq \delta$ and assume everything has been defined for $j < i$.

- **Base step:** When $i = 0$, let $q_0 \in \text{gS}^{\alpha_0}(M^2)$ be the nonforking extension of p_0 to M_0^2 (exists by extension below λ_0).
- **Successor step:** When $i = j+1$, $j < \delta$, let q_i be the nonforking extension (of length α_i) of p_i to M^2 . We have to check that q_i indeed extends q_j (i.e. $q_i^{\alpha_j} = q_j$). Note that $q_j \upharpoonright M^0$ does not fork over M_j^0 so by step 1 and uniqueness, $q_j \upharpoonright M^0 = \text{gtp}(M_j^1/M_j^0; M^1)$. In particular, $q_j \upharpoonright M_i^0 = \text{gtp}(M_j^1/M_i^0; M^1)$. Since q_i extends p_i , $q_i \upharpoonright M_i^0 = \text{gtp}(M_i^1/M_i^0; M^1)$ so $q_i^{\alpha_j} \upharpoonright M_i^0 = \text{gtp}(M_j^1/M_i^0; M^1) = q_j \upharpoonright M_i^0$. By base monotonicity, q_j does not fork over M_i^0 so by uniqueness $q_i^{\alpha_j} = q_j$. A picture is below.



- **Limit step:** Assume i is limit. Let q_i be the direct limit of the coherent sequence $\langle q_j : j < i \rangle$. Note that $q_i \in \text{gS}^{\alpha_i}(M^2)$ and by Proposition 14.12, q_i does not fork over M_i^0 . It remains to see $q_i \upharpoonright M_i^0 = p_i$.

For $j < i$, let $p'_j \in \text{gS}^{\alpha_j}(M_i^0)$ be the nonforking extension of p_j to M_i^0 . By step 1, $p'_j = \text{gtp}(M_j^1/M_i^0; M^1)$. Thus $\langle p'_j : j < i \rangle$ is type-local. By an argument similar to the successor step above, we have that for all $j < i$, $p_i^{\alpha_j} = p'_j$. Moreover, for all $j < i$, $q_i^{\alpha_j} \upharpoonright M_j^0 = p_j$ and q_j does not fork over M_j^0 so by uniqueness, $q_i^{\alpha_j} \upharpoonright M_i^0 = (q_i \upharpoonright M_i^0)^{\alpha_j} = p'_j$. By type-locality, it follows that $q_i \upharpoonright M_i^0 = p_i$, as desired.

□

Putting everything together, we get:

Theorem 14.14. If:

- (1) For some regular $\kappa \leq \lambda$, K is fully $(< \kappa)$ -tame.
- (2) K is densely type-local above λ .

Then i' is a fully good $(< \infty, \geq \lambda)$ -independence relation.

Proof. Lemma 14.6 gives most of the properties of a good independence relation. By Lemma 4.5 and symmetry, i has the left $(< \kappa)$ -model-witness property. By Lemma 14.7, i' has full model continuity. By Lemma 14.8, it has the local character properties. By Lemma 14.13, i' has extension. \square

We suspect that dense type-locality is not necessary, at least when i comes from our construction (see the proof of Theorem 15.1). For example, by the proof below, it would be enough to see that $\text{pre}(i_\mu^{\leq 1})$ is weakly successful for all $\mu \geq \lambda$. We delay a full investigation to a future work. For now, here is what we can say without dense type-locality:

Theorem 14.15. Assume that for some regular $\kappa \leq \lambda$, K is fully $(< \kappa)$ -tame. Then:

- (1) i' is a fully good independence relation, except perhaps for the extension property.
- (2) Assume that³² for all $\mu \geq \lambda$, $(i')_{\geq \mu}$ satisfies Hypothesis 11.1. Then i' has the extension property when the base is saturated.

Proof. The first part has been observed in the proof of Theorem 14.14. To see the second part, let $\mu \geq \lambda$. By Theorem 11.13 and Theorem 12.16, there exists a good $(\leq \mu, \mu)$ -independence relation i'' with underlying class $K_\mu^{\mu\text{-sat}}$. Using the witness properties and the methods of [BGKV] we have that $(i')^{\leq \mu} \upharpoonright K_\mu^{\mu\text{-sat}} = i''$. By the proof of Lemma 14.13, i' has extension when the base model is in $K_\mu^{\mu\text{-sat}}$. \square

15. THE MAIN THEOREMS

Recall (Definition 8.4) that an AEC K is fully good if there is a fully good independence relation with underlying class K . Intuitively, a fully good independence relation is one that satisfies all the basic properties of forking in a superstable first-order theory. We are finally ready to show that densely type-local fully tame and short superstable classes are fully good, at least on a class of sufficiently saturated models³³:

³²If for example i' is constructed as in the proof of Theorem 15.1, this will be the case.

³³The number 7 in (1) is possibly the largest natural number ever used in a statement about abstract elementary classes!

Theorem 15.1. Let K be a fully $(< \kappa)$ -tame and short AEC with amalgamation. Assume that K is densely type-local above κ .

- (1) If K is μ -superstable, $\kappa = \beth_\kappa > \mu$, and $\lambda := (\mu^{<\kappa_r})^{+7}$, then $K^{\lambda\text{-sat}}$ is fully good.
- (2) If K is κ -strongly μ -superstable and $\lambda := (\mu^{<\kappa_r})^{+6}$, then $K^{\lambda\text{-sat}}$ is fully good.
- (3) If K has amalgamation, $\kappa = \beth_\kappa > \text{LS}(K)$, and K is categorical in a $\mu > \lambda_0 := (\kappa^{<\kappa_r})^{+5}$, then $K_{\geq \lambda}$ is fully good, where $\lambda := \min(\mu, h(\lambda_0))$.

Proof.

- (1) By Theorem 10.11 and Proposition 10.10, K is κ -strongly $(2^{<\kappa_r})^+$ -superstable. Now apply (2).
- (2) By Fact 11.3, Hypothesis 11.1 holds for $\mu' := (\mu^{<\kappa_r})^{+2}$, λ there standing for $(\mu')^+$ here, and $K' := K^{\mu'\text{-sat}}$. By Theorem 11.21, there is an ω -successful type-full good $(\mu')^+$ -frame \mathfrak{s} on $K^{(\mu')^+\text{-sat}}$. By Theorem 12.16, \mathfrak{s}^{+3} induces a fully good $(\leq \lambda, \lambda)$ -independence relation \mathbf{i} on $K^{(\mu')^{+4}\text{-sat}} = K^{\lambda\text{-sat}}$. By Theorem 13.6, $\mathbf{i}' := \text{cl}(\text{pre}(\mathbf{i}_{\geq \lambda}))$ is a fully good $(\leq \lambda, \geq \lambda)$ -independence relation on $K^{\lambda\text{-sat}}$. By Theorem 14.14, $(\mathbf{i}')^{\text{long}}$ is a fully good $(< \infty, \geq \lambda)$ -independence relation on $K^{\lambda\text{-sat}}$. Thus $K^{\lambda\text{-sat}}$ is fully good.
- (3) By Theorem 10.16, K is κ -strongly κ -superstable. By (2), $K^{\lambda_0^+\text{-sat}}$ is fully good. By Fact 10.12.(4), all the models in $K_{\geq \lambda}$ are λ_0^+ -saturated, hence $K_{\geq \lambda}^{\lambda_0^+\text{-sat}} = K_{\geq \lambda}$ is fully good.

□

We now discuss the necessity of the hypotheses of the above theorem. It is easy to see that a fully good AEC is superstable⁺. Moreover, the existence of a relation \perp with disjointness and independent amalgamation directly implies disjoint amalgamation. An interesting question is whether there is a general framework in which to study independence without assuming amalgamation, but this is out of the scope of this paper. To justify full tameness and shortness, one can ask:

Question 15.2. Let K be a fully good AEC. Is K fully tame and short?

If the answer is positive, we believe the proof to be nontrivial. We suspect however that the shortness hypothesis of our main theorem can

be weakened to a condition that easily holds in all fully good classes. In fact, we propose the following:

Definition 15.3. An AEC K is *diagonally $(< \kappa)$ -tame* if for any $\kappa' \geq \kappa$, K is $(< \kappa')$ -tame for types of length less than κ' . K is *diagonally κ -tame* if it is diagonally $(< \kappa^+)$ -tame. K is *diagonally tame* if it is diagonally $(< \kappa)$ -tame for some κ .

It is easy to check that if \mathbf{i} is a good $(< \infty, \geq \lambda)$ -independence relation, then $K_{\mathbf{i}}$ is diagonally λ -tame. Thus we suspect the answer to the following should be positive:

Question 15.4. In Theorem 15.1, can “fully $(< \kappa)$ -tame and short” be replaced by “diagonally $(< \kappa)$ -tame”?

Finally, we believe the dense type-locality hypothesis can be removed³⁴. Indeed, chapter III of [She09] has many results on getting models “generated” by independent sequences. Since independent sequences exhibit a lot of finite character (see also [BVb]), we suspect the following conjecture should be true. Note that if it holds, then one can remove the dense type-locality hypothesis from Theorem 15.1.

Conjecture 15.5. Let K be a fully $(< \kappa)$ -tame and short abstract elementary class. If K is κ -strongly μ -superstable and $\lambda := (\mu^{< \kappa r})^{+6}$, then $K^{\lambda\text{-sat}}$ is densely type-local above λ .

We end with some results that do not need dense type-locality. Note that we can replace categoricity by superstability or strong superstability as in the proof of Theorem 15.1.

Theorem 15.6. Let K be a fully $(< \kappa)$ -tame and short AEC with amalgamation. Let λ, μ be cardinals such that:

$$\text{LS}(K) < \kappa = \beth_{\kappa} < \lambda = \beth_{\lambda} \leq \mu$$

Assume further that $\text{cf}(\lambda) \geq \kappa$. If K is categorical in μ , then:

- (1) There exists an ω -successful type-full good λ -frame \mathfrak{s} with $K_{\mathfrak{s}} = K_{\lambda}$. Furthermore, the frame is induced by $(< \kappa)$ -coheir: $\mathfrak{s} = \text{pre}(\mathbf{i}_{\kappa\text{-ch}}(K)_{\lambda}^{\leq 1})$.
- (2) K_{λ} is $(\leq \lambda, \lambda)$ -good.
- (3) $K^{\lambda^{+3}\text{-sat}}$ is fully $(\leq \lambda^{+3})$ -good.

³⁴In fact, the result was initially announced without this hypothesis but Will Boney found a mistake in our proof of Lemma 14.13. This is the only place where type-locality is used

- (4) $K^{\lambda+3\text{-sat}}$ is fully good, except it may not have extension. Moreover it has extension over saturated models. In particular, if $K^{\lambda+3\text{-sat}}$ is categorical in all cardinals, then it is fully good.

Proof. By cardinal arithmetic, $\lambda = \lambda^{<\kappa_r}$. By Fact 11.3 and Theorem 11.21, there is an ω -successful type-full good λ -frame \mathfrak{s} with $K_{\mathfrak{s}} = K_{\lambda}^{\lambda\text{-sat}}$. Now (by Theorem 10.16 if $\mu > \lambda$), K is categorical in λ . Thus $K^{\lambda\text{-sat}} = K_{\geq\lambda}$. Theorem 12.16 and Theorem 13.6 give the next two parts. Theorem 14.15 gives the last part. \square

REFERENCES

- [Bal09] John T. Baldwin, *Categoricity*, University Lecture Series, vol. 50, American Mathematical Society, 2009.
- [BG] Will Boney and Rami Grossberg, *Forking in short and tame AECs*, Preprint. URL: <http://arxiv.org/abs/1306.6562v7>.
- [BGKV] Will Boney, Rami Grossberg, Alexei Kolesnikov, and Sebastien Vasey, *Canonical forking in AECs*, Preprint. URL: <http://arxiv.org/abs/1404.1494v2>.
- [BKV06] John T. Baldwin, David Kueker, and Monica VanDieren, *Upward stability transfer for tame abstract elementary classes*, Notre Dame Journal of Formal Logic **47** (2006), no. 2, 291–298.
- [BL03] Steven Buechler and Olivier Lessmann, *Simple homogeneous models*, Journal of the American Mathematical Society **16** (2003), no. 1, 91–121.
- [Bon14a] Will Boney, *Tameness and extending frames*, Journal of Mathematical Logic **14** (2014), no. 2, 1450007.
- [Bon14b] ———, *Tameness from large cardinal axioms*, The Journal of Symbolic Logic **79** (2014), no. 4, 1092–1119.
- [BS08] John T. Baldwin and Saharon Shelah, *Examples of non-locality*, The Journal of Symbolic Logic **73** (2008), 765–782.
- [BVa] Will Boney and Sebastien Vasey, *Chains of saturated models in AECs*, Preprint. URL: <http://arxiv.org/abs/1503.08781v1>.
- [BVb] ———, *Tameness and frames revisited*, Preprint. URL: <http://arxiv.org/abs/1406.5980v4>.
- [Gro] Rami Grossberg, *A course in model theory I*, A book in preparation.
- [Gro02] ———, *Classification theory for abstract elementary classes*, Contemporary Mathematics **302** (2002), 165–204.
- [GV06a] Rami Grossberg and Monica VanDieren, *Categoricity from one successor cardinal in tame abstract elementary classes*, Journal of Mathematical Logic **6** (2006), 181–201.
- [GV06b] ———, *Galois-stability for tame abstract elementary classes*, Journal of Mathematical Logic **6** (2006), no. 1, 25–49.
- [GV06c] ———, *Shelah’s categoricity conjecture from a successor for tame abstract elementary classes*, The Journal of Symbolic Logic **71** (2006), no. 2, 553–568.

- [GVV] Rami Grossberg, Monica VanDieren, and Andrés Villaveces, *Uniqueness of limit models in classes with amalgamation*, Preprint. URL: <http://arxiv.org/abs/math/0509338v3>.
- [HK06] Tapani Hyttinen and Meeri Kesälä, *Independence in finitary abstract elementary classes*, *Annals of Pure and Applied Logic* **143** (2006), 103–138.
- [HL02] Tapani Hyttinen and Olivier Lessmann, *A rank for the class of elementary submodels of a superstable homogeneous model*, *The Journal of Symbolic Logic* **67** (2002), no. 4, 1469–1482.
- [Jara] Adi Jarden, *Primeness triples in non-forking frames*, Preprint. Available online from <http://ariel.academia.edu/AdiJarden>.
- [Jarb] ———, *Tameness, uniqueness triples, and amalgamation*, Preprint. URL: <http://arxiv.org/abs/1408.3383v2>.
- [JS12] Adi Jarden and Alon Sitton, *Independence, dimension and continuity in non-forking frames*, *The Journal of Symbolic Logic* **78** (2012), no. 2, 602–632.
- [JS13] Adi Jarden and Saharon Shelah, *Non-forking frames in abstract elementary classes*, *Annals of Pure and Applied Logic* **164** (2013), 135–191.
- [Lie11] Michael J. Lieberman, *A topology for Galois types in abstract elementary classes*, *Mathematical Logic Quarterly* **57** (2011), no. 2, 204–216.
- [LP79] Daniel Lascar and Bruno Poizat, *An introduction to forking*, *The Journal of Symbolic Logic* **44** (1979), no. 3, 330–350.
- [MS90] Michael Makkai and Saharon Shelah, *Categoricity of theories in $L_{\kappa, \omega}$, with κ a compact cardinal*, *Annals of Pure and Applied Logic* **47** (1990), 41–97.
- [She] Saharon Shelah, *Eventual categoricity spectrum and frames*, Paper number 842 on Shelah’s publication list. Preliminary draft from Oct. 3, 2014 (obtained from the author).
- [She78] ———, *Classification theory and the number of non-isomorphic models*, *Studies in logic and the foundations of mathematics*, vol. 92, North-Holland, 1978.
- [She90] ———, *Classification theory and the number of non-isomorphic models*, 2nd ed., *Studies in logic and the foundations of mathematics*, vol. 92, North-Holland, 1990.
- [She99] ———, *Categoricity for abstract classes with amalgamation*, *Annals of Pure and Applied Logic* **98** (1999), no. 1, 261–294.
- [She00] ———, *On what I do not understand (and have something to say), model theory*, *Math. Japonica* **51** (2000), 329–377.
- [She01] ———, *Categoricity of an abstract elementary class in two successive cardinals*, *Israel Journal of Mathematics* **126** (2001), 29–128.
- [She09] ———, *Classification theory for abstract elementary classes*, *Studies in Logic: Mathematical logic and foundations*, vol. 18, College Publications, 2009.
- [SV99] Saharon Shelah and Andrés Villaveces, *Toward categoricity for classes with no maximal models*, *Annals of Pure and Applied Logic* **97** (1999), 1–25.
- [Van06] Monica VanDieren, *Categoricity in abstract elementary classes with no maximal models*, *Annals of Pure and Applied Logic* **141** (2006), 108–147.

- [Van13] ———, *Erratum to "Categoricity in abstract elementary classes with no maximal models"* [*Ann. Pure Appl. Logic* 141 (2006) 108-147], *Annals of Pure and Applied Logic* **164** (2013), no. 2, 131–133.
- [Vasa] Sebastien Vasey, *Forking and superstability in tame AECs*, Preprint. URL: <http://arxiv.org/abs/1405.7443v2>.
- [Vasb] ———, *Infinitary stability theory*, Preprint. URL: <http://arxiv.org/abs/1412.3313v3>.

E-mail address: `sebv@cmu.edu`

URL: <http://math.cmu.edu/~svasey/>

DEPARTMENT OF MATHEMATICAL SCIENCES, CARNEGIE MELLON UNIVERSITY,
PITTSBURGH, PENNSYLVANIA, USA