

Why has active immunotherapy not worked in lung cancer?

A. Thomas¹, G. Giaccone²

¹Thoracic and GI Oncology Branch, National Cancer Institute, Bethesda, Maryland, USA

²Lombardi Comprehensive Cancer Center, Georgetown University, Washington DC, USA

Address for correspondence: Prof. Giuseppe Giaccone, Lombardi Comprehensive Cancer Center, Georgetown University, Research Building Room W503, 3970 Reservoir Road NW, Washington DC 20057. Phone: 001-240-6877072. Email: gg496@georgetown.edu.

Key Message: "This paper reviews the phase III trial results of antigen-specific immunotherapeutic approaches in NSCLC and explore in-depth the potential reasons behind their failure and discuss strategies for the future."

abstract

Vaccines which rely on active-specific stimulation of the host immune system have the potential to trigger durable antitumor responses with minimal toxicity. However in non-small cell lung cancer (NSCLC), several large phase III trials of vaccines reported within the last year have yielded disappointing results. Compared with placebo, belagenpumatucel-L (an allogenic tumor cell vaccine), tecemotide (a peptide vaccine targeting MUC-1) and melanoma associated antigen-A3 (a protein-based vaccine) did not improve outcomes in NSCLC. The lack of clinically significant outcomes, despite their ability to prime and expand tumor antigen-specific T cells could at least partly be attributed to the inability of vaccine-induced T-cell responses to overcome the tumoral mechanisms of immune escape which limit the clonal expansion of T cells following vaccination. A number of such mechanisms have been recognized including reduced antigen presentation, antigenic loss, cytokines, immunosuppressive cells and immune checkpoints. Strategies aimed at modulating the immune checkpoints have shown promise and are on the verge of revolutionizing the therapeutic landscape of metastatic NSCLC. Overcoming immune tolerance and improving the activation of antitumor T cells via combinatorial approaches may represent a new and more promising therapeutic application for active immunotherapies in NSCLC.

keywords:

active immunotherapy; vaccines; non-small cell lung cancer; immune checkpoint; tumor-mediated immunosuppression

introduction

The significant and durable responses induced by antibodies blocking the programmed cell death-1 (PD-1) checkpoint have led to a renewed interest in immunotherapy for non-small cell lung cancer (NSCLC).[1, 2] These results are particularly encouraging given the many unsuccessful attempts at immunotherapy in NSCLC over the last several years. In general these have included active immunotherapies which rely on the ability of the patient's own immune system to mount an immune response specific to tumor-associated antigens, passive immunotherapy which uses exogenous lymphocytes or antibodies to mediate an immune response, and non-specific immune stimulation which should be effective regardless of the tumor antigen which stimulates the immune response.[3, 4]

Active-specific stimulation of the host immune system has the potential to cause durable antitumor responses with minimal toxicity. This promise of antigen-specific immunotherapy has borne out in prostate cancer where the use of sipuleucel-T, an autologous active cellular immunotherapy prolonged overall survival (OS) among men with metastatic castration-resistant prostate cancer [5]. However in NSCLC, several agents whose large phase III trial results have been reported within the last year have yielded no significant benefit. Given the dire need for better therapies and the cost of drug development, it is imperative to try to understand these failures. In this article, we will review the phase III trial results of recently reported antigen-specific immunotherapeutic approaches in NSCLC, explore the potential reasons behind their failure and discuss strategies for the future.

antigen-specific immunotherapeutic approaches in NSCLC

Belagenpumatucel-L

Belagenpumatucel-L (Lucanix) is an allogeneic tumor cell vaccine, which consists of four irradiated NSCLC cell lines that have been modified with transforming growth factor beta-2

(TGF- β 2) antisense gene plasmid. TGF- β inhibits T cell, B cell, and dendritic cell activation, induces immunosuppressive T regulatory (Treg) cells and inhibits immune effector cell activation.[6] In a phase II study of patients with low volume disease, belagenpumatucel-L was well tolerated, induced antibody-mediated response to vaccine human leukocyte antigens (HLA), and demonstrated a dose-dependent improvement in survival and response.[7]

A phase III trial compared the efficacy of belagenpumatucel-L with placebo as a maintenance therapy in patients with stages IIIA (T3, N2 only), IIIB and IV NSCLC without progression after up to six cycles of first-line platinum-based chemotherapy (which had to be completed 4-17 weeks prior to randomization).[8] Belagenpumatucel-L (2.5×10^7 cells/injection intradermally) or placebo were administered every month for 18 months followed by additional two quarterly injections. The primary endpoint was OS. Maintenance belagenpumatucel-L did not result in improvement in OS over placebo [median OS 20.3 months with belagenpumatucel-L ($n=270$) and 17.8 months with placebo ($n=262$); $p=0.59$]. Of interest, however, in a pre-planned subgroup analysis, among patients who received prior radiation therapy and enrolled within 12 weeks, belagenpumatucel-L resulted in significantly improved OS [median OS 40.1 months with belagenpumatucel-L ($n=43$) and 10.3 months with placebo ($n=36$); $p=0.014$].

Tecemotide

Tecemotide (Liposomal BLP25; L-BLP25) is a peptide vaccine, which targets the exposed core peptide of MUC-1, a membrane associated glycoprotein differentially over-expressed and aberrantly glycosylated in cancer cells [9, 10]. Tecemotide consists of the MUC1-derived 25-aminoacid BLP25 lipopeptide, the immunoadjuvant monophosphoryl lipid A, and three liposome-forming lipids. Tecemotide was well tolerated and induced T-cell responses to MUC1 in phase I and II studies.[11-13]

A phase III trial compared the efficacy of tecemotide with placebo (2:1 randomization) as a maintenance therapy in patients with unresectable stage III NSCLC who had responded to or had stable disease after primary chemoradiotherapy (which had to be completed within 4–12 weeks prior to randomization).[14] One dose of cyclophosphamide (300 mg/m² intravenously, maximum dose 600 mg) or placebo was administered prior to treatment. Eight consecutive weekly subcutaneous injections of tecemotide or placebo were followed in the absence of progressive disease by maintenance tecemotide or placebo every 6 weeks until disease progression. The primary endpoint was OS. Maintenance tecemotide did not result in improvement in OS over placebo [median OS 25.6 months with tecemotide ($n=829$) and 22.3 months with placebo ($n=410$) (HR 0.88, 0.75-1.03; $p=0.123$)]. In a pre-planned subgroup analysis, however, among patients who received concurrent chemoradiotherapy, OS was significantly longer with tecemotide than placebo [median OS 30.8 months (95% CI, 25.6-36.8) with tecemotide ($n=538$) and 20.6 months (95% CI, 17.4-23.9) with placebo ($n=268$)]. However, in patients who received previous sequential chemoradiotherapy, OS was worse in patients in the tecemotide group [median OS 19.4 months (95% CI, 17.6-23.1; $n=291$) and 24.6 months (95% CI, 18.8–33.0) with placebo ($n=142$) (HR 1.12, 0.87-1.44; $p=0.38$)]. Based on these results, an ongoing trial is studying the effect of tecemotide or placebo on OS of patients with unresectable stage III NSCLC with either stable disease or objective response following primary concurrent chemo-radiotherapy (ClinicalTrials.gov Identifier: NCT02049151).

Melanoma associated antigen-A3 vaccine

Melanoma associated antigen-A3 (MAGE-A3) vaccine is a protein-based vaccine consisting of the recombinant antigen ProtD-MAGE-A3/His (a fusion protein containing Protein D, a lipoprotein present on the surface of haemophilus influenzae B, MAGE-A3 protein, and a polyhistidine tail) and a proprietary immunological adjuvant. Melanoma associated antigens (MAGE) are tumor-specific shared antigens which are differentially over-expressed in many

cancers including NSCLC. In a phase II trial of patients with completely resected, MAGE-A3-expressing early stage NSCLC, humoral and cellular immune responses to MAGE-A3 and statistically non-significant improvements in disease-free intervals were observed.[15, 16]

A phase III trial compared the efficacy of MAGE-A3 vaccine with placebo (2:1 randomization) in patients with completely resected MAGE-A3-expressing stage IB, II or IIIA NSCLC. Up to four cycles of adjuvant chemotherapy could be administered at the investigators' discretion. Thirteen doses of the vaccine were administered intramuscularly over 27 months. The primary objectives were disease-free survival (DFS) in the overall population and in those who did not receive adjuvant chemotherapy (co-primary endpoints). The trial enrolled 2312 MAGE-A3-positive patients (33% of patients screened had MAGE-A3 expressing tumors). The study was terminated by an independent data monitoring committee as MAGE-A3 vaccine did not significantly extend DFS compared with placebo either in the overall MAGE-A3 positive population or in those MAGE-A3-positive patients who did not receive chemotherapy.[17]

considerations for active immunotherapy in NSCLC

While a number of factors are important in clinical translation of successful active immunotherapy (Figure 1), we will discuss some which are more relevant in the context of the above described negative large phase III trials.

Humoral and Cellular Immune Dysregulation in Lung Cancer

In the first step of an adaptive immune response, effector T cells recognize antigenic peptides of tumor cells presented by antigen-presenting cells (APC) in the context of major histocompatibility complex (MHC) class I or class II molecules expressed on the APC surface. Additional co-stimulatory signals mediated through constitutively expressed co-stimulatory molecules on the T cell and the APC are also necessary for T cell activation. The presence of both signals trigger intracellular events resulting in the activation and interleukin (IL)-2-

dependent clonal proliferation of T cells. Expansion of T cells in sufficient numbers results in recognition and elimination of tumor cells. However immune responses are dysregulated in cancer.

A number of mechanisms are employed by tumors to escape the host immune response and promote immune tolerance. These are perhaps the most important hurdles that need to be overcome for successful antigen-specific immunotherapy in NSCLC. The better understood immune resistance mechanisms in NSCLC are outlined in Figure 2.

Suppression of antigen-presenting machinery is one of several mechanisms of immune escape. Multiple molecular mechanisms can lead to altered HLA expression within lung cancer. These include deficiencies in expression of antigen-processing genes [18-21], and haplotype loss of HLA class I antigens [22-24] In small retrospective studies, absence HLA class I expression was associated with poor prognosis suggesting that down-regulation of HLA class I expression may play a critical role in immune surveillance of patients with NSCLC.[25, 26] The reversibility of some of the aberrations in antigen processing by interferon (IFN)-gamma indicates that it is possible to overcome the suppression of antigen presenting machinery and may be of therapeutic relevance.[27, 28] Considering the critical role of antigen presentation in immune recognition of tumor cells, these mechanisms may be of potential therapeutic importance.

In addition to reduced antigen presentation, immune inhibitory cytokines secreted by the tumor cells can impair T-cell survival and help them avoid T cell-mediated immune responses. Soluble factors derived from NSCLC cell line supernatants have been described to markedly enhanced apoptosis of activated T cells [29]. Transforming growth factor beta (TGF- β) enables tumor evasion of immune surveillance through various mechanisms most of which converge on the impairment of tumor cell killing by immune effector cells. [30] In addition to inhibiting proliferation and differentiation of normal bronchial epithelial cells, TGF- β mediates conversion of

CD4+CD25⁻ T cells to Tregs [31, 32]. Serum TGF- β levels are elevated in patients with lung cancer compared with normal individuals. Elevated plasma levels of TGF- β confer a poorer prognosis for patients with lung cancer [33]. IL-10 is a potent immunosuppressive cytokine that promotes lung cancer growth by suppressing T-cell and macrophage function and enabling tumors to escape immune detection [34-36]

Yet another mechanism of immunosuppression involves immune checkpoints which are molecules expressed on the surface of T lymphocytes and modulates the immune response to antigens via inhibitory or stimulatory signaling to T cells. Two most extensively studied immune inhibitory checkpoints in NSCLC are cytotoxic T-lymphocyte antigen-4 (CTLA-4) and PD-1. Activation of both receptors causes down-regulation and inhibition of immune responses. PD-1 functions primarily in peripheral tissues where T cells may encounter the immunosuppressive PD-1 ligands, PD-L1 (B7-H1) and PD-L2 (B7-DC), which are expressed by tumor cells, stromal cells, or both.[37] CTLA-4 mediates immune inhibitory signals which are distinct from PD-1.[38] Clinical trial results of antibody-mediated blocking of CTLA-4 and PD-1 pathways indicate that this strategy is feasible and effective in NSCLC.[1,39]

A number of cells in the tumor microenvironment including Tregs, myeloid-derived suppressor cells (MDSCs) and tumor-associated macrophages have immunosuppressive properties. Tumor-infiltrating lymphocytes (TILs) which are CD4+CD25⁺, the activated phenotype of Tregs, mediates potent inhibition of autologous T cell proliferation and prevents the host from mounting an immune response to tumor antigens [40]. Tregs of a similar phenotype (CD4+CD25⁺) with marked immunosuppressive activity are elevated in peripheral blood of NSCLC patients [41]. MDSCs are a heterogeneous population of cells of myeloid origin that are characterized by their immature state and ability to suppress T-cell responses [42]. In lung cancer, antibody mediated MDSC depletion increased APC activity and augmented the activity of effector T cells leading to

reduced tumor growth and enhanced therapeutic vaccination responses [43]. The prognostic significance of MDSCs in the tumor microenvironment is not established in NSCLC.

A number of metabolic enzymes including those associated with the catabolism of the amino acids arginine and tryptophan are associated with the suppressive activity of myeloid cells.

Indoleamine 2,3-dioxygenase-1 (IDO1) is an enzyme which is expressed by a subset of dendritic cells that catalyzes the degradation of the amino acid tryptophan to kynurenine. IDO1 is thought to be an important regulator of the immunosuppressive mechanisms responsible for tumor escape from host immune surveillance and blockade of IDO activity increases the ability of tumor-bearing mice to reject tumors.[44]

In summary, a number of mechanisms including reduced antigen presentation, antigenic loss, cytokines, immune checkpoints, immunosuppressive cells, and enzymes are employed by tumors to escape the host immune response and promote immune tolerance.

Trial Design

With the benefit of hindsight, the negative results of these large phase III trials (with a combined accrual of over 4000) should come as no surprise. All three trials were initiated based on results of negative or at best inconclusive phase II data (Table 1) and post hoc analysis of small subgroups which showed positive results.

For example, a randomized, open label, phase II trial failed to show significant improvement in OS of patients who received tecemotide over those who received best supportive care. In the small subset of patients with stage IIIB-LR (loco-regional) disease ($n=65$), those who received tecemotide had a 17.3 month improvement in median OS (30.6 versus 13.3 months) [12]

In another instance, the phase III trial of belagenpumatucel-L was initiated based on a dose-related improvement in survival and response in the phase II trial. However the phase II trial

itself had small numbers of patients in the individual treatment arms (about 20 patients each in the three cohorts) who had low volume disease. Furthermore the phase II trial did not have a control arm.[7, 8]

In a third instance, the randomized, placebo-controlled phase II trial of MAGE-A3 vaccine which led to the larger phase III trial, had a limited sample size. With 182 patients, and an estimated power of 50% to detect a difference of 10% in absolute recurrence after 30 months, the study was unlikely to demonstrate improvements in efficacy. A related issue, highlighted by the phase II to III transition of this drug is the lack of adequate follow up. Trends of activity observed in earlier analysis were not confirmed with more mature follow up data [15, 16]. A number of factors including commercial pressures and misguided enthusiasm of investigators based on early trends may explain these failures.

While it is true that investigators would not initiate a trial if they did not think it had a reasonable chance of a statistically significant and clinically meaningful benefit, some have argued that the investigators frequently use overly optimistic assumptions of treatment benefits.[45]

Unfortunately, this may have been true in the transition from phase II to phase III trials of antigen-specific immunotherapies in NSCLC.

future of antigen-specific immunotherapy in NSCLC

The failure of vaccines in NSCLC, despite their ability to prime and expand tumor antigen-specific T cells, could at least partly be attributed to the inability of vaccine-induced T-cell responses to overcome the tumoral mechanisms of immune escape. These mechanisms probably limit the clonal expansion of T cells following vaccination.

Many of the immunosuppressive mechanisms discussed above are potentially amenable to therapeutic modulation. Low doses of cyclophosphamide have been shown to selectively decrease circulating Tregs and suppress their inhibitory functions leading to a restoration of

peripheral T cell proliferation and innate killing activity [46]. Other drugs including chemotherapies and signal transduction inhibitors have also been shown to selectively target immunosuppressive cells in the tumor microenvironment.[47,48] Metabolic enzymes and cytokines involved in the induction of tumor immune tolerance can also be inhibited pharmacologically [43, 48]. MDSC differentiation can be blocked in a number of ways including cyclooxygenase inhibitors, which prevent the production of prostaglandin.[50]

Recent studies have demonstrated that immune checkpoints can be successfully modulated [1, 2]. An anti-PD-1 antibody, nivolumab was evaluated in a phase I trial in patients with advanced previously treated cancers.[1] Doses of 1, 3, and 10 mg/kg were administered intravenously once every 2 weeks with immune response assessment every eight weeks. In the NSCLC expansion cohort, across all doses and histologies (squamous and non-squamous), the ORR was 17% (22 of 129 patients) and median response duration 17 months.[51] Median OS was 9.2 to 14.9 months and 1-year OS rates 32 to 56%. In March 2015, nivolumab was approved by the FDA for use in patients with metastatic squamous cell lung cancer with progression on or after platinum-based chemotherapy. Its efficacy was established in a phase III, open-label, study that randomized previously treated patients ($n=272$) with advanced squamous cell lung cancer to receive nivolumab 3 mg/kg intravenously every two weeks or docetaxel 75 mg/m² intravenously every three weeks. Overall survival, the primary end point of the trial, was prolonged by 3.2 months at the median in patients who received nivolumab compared with those who received docetaxel. Several other agents targeting PD-1 pathway are in clinical development, including pembrolizumab (MK-3475, anti PD1), MEDI4736 (anti-PDL1), BMS-936559 (anti-PDL1) and MPDL-3280 (anti-PDL1). Despite the promise of immune checkpoint inhibitors, it is clear that responses are limited, restricted presumably to patients with a preexisting tumor-reactive T-cell response. Investigations of ways to select patients (e.g. PDL-1 expression in the tumor or infiltrating immune cells or both) are underway. There is growing

interest in modulating the multiple immune inhibitory and co-stimulatory pathways in the tumor microenvironment by combining inhibitors of the PD-1 pathway with other immune checkpoints antibodies, including antagonist antibodies to KIR, LAG-3 and CTLA-4.

Antigen-specific vaccines offer an opportunity to potentially extend the responses with immune checkpoint inhibitors to a greater percentage of patients. A recent study showed that tumors resistant to anti PD-1 antibodies could be eradicated by combining them with vaccines containing tumor-specific peptides with high MHC-binding affinity [52]. In the study, melanomas that contained a high percentage of dysfunctional endogenous PD-1+ tumor-specific CD8+ T cells were treated with a PD-1 inhibitor and an exogenous tumor-specific antigen using attenuated *Salmonella Typhimurium*. The combination rescued the endogenous tumor-specific CD8+ T-cell response and resulted in tumor regressions. A combinatorial strategy of vaccines and immune checkpoint inhibitors could rescue T cells which become dysfunctional after infiltrating long-established suppressive tumors, thereby overcoming one of the major obstacles to clinical benefit from vaccines. Most of these strategies are still in pre-clinical evaluation in NSCLC. While there is strong rationale to combine vaccines with other immunomodulatory strategies, important considerations in clinical testing of these combinations include determining the sequence of administration of drugs, and metrics of response assessment.

While the above discussed approaches aim to overcome tumor-mediated immunosuppression, other approaches seek to enhance cellular immune responses through a number of different mechanisms. These include induction of immunogenic cell death with radiotherapy [53] and combination with adoptive T cell transfer to prime T cells and amplify anti-tumor T cell responses.[54] Immunogenic cell death is different from apoptotic cell death in the generation of specific molecular signals that are sensed by APC which stimulate their maturation and ability to cross-present tumor-derived antigens to T cells.[55] In addition to immunogenic cell death, radiation causes MHC I upregulation, and release of antigens which are taken up by dendritic

cells and presented to T cells that in turn migrate back to the tumor and provide local control, thus serving as an intrinsic vaccine priming adaptive immunity.[56] The ongoing process of killing of tumor cells by cytotoxic T lymphocytes sustains release of more tumor antigens and possibly promotes antigenic spread, i.e. the activation of a broader T-cell repertoire. Antigenic spread has been reported in some patients with prostate cancer who were treated with the combination of a vaccine and local radiotherapy.[57]

Possible beneficial effects observed in subsets of patients on active immunotherapy trials indicates the need for better patient selection.[8, 14] While it is generally believed that these therapies are most active in patients with minimal volume of disease, no predictive markers have been identified to date. Better measures are needed to assess tumor-specific immune responses and understand the relationship between immune induction and clinical responses. The failure of phase III trials which were initiated based on “promising” phase II trials also indicate the need to temper our optimism, particularly when making the expensive leap from phase II to phase III trials.

Finally, a better understanding of the immune dysregulation specific to NSCLC is needed. The immune evasion mechanisms in lung cancer are likely different from other tumors [58] due to the pro-inflammatory and immunosuppressive effects of tobacco smoke. Chronic inhalation of cigarette smoke is known to alter a wide range of immunological functions, including innate and adaptive immune responses.[59] In the context of active immunotherapy, the effect of smoking on T cell responsiveness and proliferative capacity are important considerations. In animal models, chronic exposure to cigarette smoke affects T-cell responsiveness and decreases T-cell proliferative and T-cell dependent antibody responses.[60] Yet there are limited data on the effects of cigarette smoke on immune dysregulation in lung cancer patients. Challenges to this field of study include the multipartite nature of cigarette smoke and the significant variability in smoking patterns which makes it difficult to study its effect in experimental systems.[61] Recent

data indicating that smoking-associated NSCLC may respond better to immune checkpoint blockade [62] also suggests the distinctive influence of tobacco smoke on the tumor microenvironment. To our knowledge clinical reports of active specific immunostimulatory agents have not assessed the effect if any of smoking on the clinical or immune outcomes. Heterogeneity within NSCLC, between the primary tumor and metastatic sites and between tumors from different patients is well described.[63] However, our understanding of the association between oncogenes and immune escape and the differential influences of different oncogenic drivers on the immune milieu are still preliminary.[64] A study of PD-L1 expression by immunohistochemistry in surgically resected NSCLC samples showed a significant association between PD-L1 expression and the presence of *EGFR* mutations independent of other clinical factors studied.[65] In preclinical models, *EGFR* mutation-positive NSCLC may preferentially use PD-1/PD-L1-mediated mechanisms to evade immune surveillance.[64] In mouse models of lung cancer, tumors with different oncogenic drivers were characterized by distinct immune infiltrates.[66] Taken together, these data suggests the potentially distinctive effects on the immune microenvironment in individual genetic subsets of NSCLC. Further understanding of how NSCLCs with different genetic backgrounds shape the tumor immune milieu will help refine the use of active specific immunotherapy in NSCLC.

In conclusion, despite their ability to prime and expand tumor antigen-specific T cells, large phase III trials of several active specific immunostimulatory agents have yielded disappointing results in NSCLC. Several important issues need to be addressed to fully harness the therapeutic potential of antitumor immune responses induced by active immunotherapy. Strategies aimed at overcoming immune tolerance and improving the activation of antitumor T cells via combinatorial approaches may represent a new and more promising therapeutic application for active immunotherapies in NSCLC.

funding

This work was supported by the Intramural program, National Cancer Institute, National Institutes of Health. No grant numbers apply.

disclosure

The authors have declared no conflicts of interest

references

1. Topalian SL, Hodi FS, Brahmer JR et al. Safety, activity, and immune correlates of anti-PD-1 antibody in cancer. *N Engl J Med* 2012; 366: 2443-2454.
2. Brahmer JR, Drake CG, Wollner I et al. Phase I study of single-agent anti-programmed death-1 (MDX-1106) in refractory solid tumors: safety, clinical activity, pharmacodynamics, and immunologic correlates. *J Clin Oncol* 2010; 28: 3167-3175.
3. Thomas A, Hassan R. Immunotherapies for non-small-cell lung cancer and mesothelioma. *Lancet Oncol* 2012; 13: e301-310.
4. Shepherd FA, Douillard JY, Blumenschein GR, Jr. Immunotherapy for non-small cell lung cancer: novel approaches to improve patient outcome. *J Thorac Oncol* 2011; 6: 1763-1773.
5. Kantoff PW, Higano CS, Shore ND et al. Sipuleucel-T immunotherapy for castration-resistant prostate cancer. *N Engl J Med* 2010; 363: 411-422.
6. Ikushima H, Miyazono K. TGFbeta signalling: a complex web in cancer progression. *Nat Rev Cancer* 2010; 10: 415-424.
7. Nemunaitis J, Dillman RO, Schwarzenberger PO et al. Phase II study of belagenpumatucel-L, a transforming growth factor beta-2 antisense gene-modified allogeneic tumor cell vaccine in non-small-cell lung cancer. *J Clin Oncol* 2006; 24: 4721-4730.
8. Giaccone LB, Nemunaitis J, Juhász RR, van den Heuvel MM, Lal R, Dunlop EC, and Fakhrai H. A phase III study of belagenpumatucel-L therapeutic tumor cell vaccine for non-small cell lung cancer (NSCLC). *ESMO* 2013.
9. North S, Butts C. Vaccination with BLP25 liposome vaccine to treat non-small cell lung and prostate cancers. *Expert Rev Vaccines* 2005; 4: 249-257.
10. Sangha R, Butts C. L-BLP25: a peptide vaccine strategy in non small cell lung cancer. *Clin Cancer Res* 2007; 13: s4652-4654.

11. Butts C, Murray N, Maksymiuk A et al. Randomized phase IIB trial of BLP25 liposome vaccine in stage IIIB and IV non-small-cell lung cancer. *J Clin Oncol* 2005; 23: 6674-6681.
12. Butts C, Maksymiuk A, Goss G et al. Updated survival analysis in patients with stage IIIB or IV non-small-cell lung cancer receiving BLP25 liposome vaccine (L-BLP25): phase IIB randomized, multicenter, open-label trial. *J Cancer Res Clin Oncol* 2011; 137: 1337-1342.
13. Butts C, Murray RN, Smith CJ et al. A multicenter open-label study to assess the safety of a new formulation of BLP25 liposome vaccine in patients with unresectable stage III non-small-cell lung cancer. *Clin Lung Cancer* 2010; 11: 391-395.
14. Butts C, Socinski MA, Mitchell PL et al. Tecemotide (L-BLP25) versus placebo after chemoradiotherapy for stage III non-small-cell lung cancer (START): a randomised, double-blind, phase 3 trial. *Lancet Oncol* 2014; 15: 59-68.
15. Vansteenkiste J ZM, Linder A et al. Final results of a multi-center, double-blind, randomized, placebo-controlled phase II study to assess the efficacy of MAGE-A3 immunotherapeutic as adjuvant therapy in stage IB/II non-small cell lung cancer (NSCLC). *J Clin Oncol* 2007; 25 (suppl 18): abstr 7554. 2007.
16. Vansteenkiste J, Zielinski M, Linder A, et al. Adjuvant MAGE-A3 immunotherapy in resected non-small-cell lung cancer: phase II randomized study results. *J Clin Oncol* 2013; 31: 2396-403.
16. Update on phase III clinical trial of investigational MAGE-A3 antigen-specific cancer immunotherapeutic in non-small cell lung cancer. GSK Press Release. April 2 2014.
18. Chen HL, Gabrilovich D, Tampe R et al. A functionally defective allele of TAP1 results in loss of MHC class I antigen presentation in a human lung cancer. *Nat Genet* 1996; 13: 210-213.
19. Singal DP, Ye M, Qiu X. Molecular basis for lack of expression of HLA class I antigens in human small-cell lung carcinoma cell lines. *Int J Cancer* 1996; 68: 629-636.

20. Johnsen A, France J, Sy MS, Harding CV. Down-regulation of the transporter for antigen presentation, proteasome subunits, and class I major histocompatibility complex in tumor cell lines. *Cancer Res* 1998; 58: 3660-3667.
21. Restifo NP, Esquivel F, Kawakami Y et al. Identification of human cancers deficient in antigen processing. *J Exp Med* 1993; 177: 265-272.
22. Baba T, Hanagiri T, Ichiki Y et al. Lack and restoration of sensitivity of lung cancer cells to cellular attack with special reference to expression of human leukocyte antigen class I and/or major histocompatibility complex class I chain related molecules A/B. *Cancer Science* 2007; 98: 1795-1802.
23. Hiraki A, Kaneshige T, Kiura K et al. Loss of HLA haplotype in lung cancer cell lines: implications for immunosurveillance of altered HLA class I/II phenotypes in lung cancer. *Clin Cancer Res* 1999; 5: 933-936.
24. Sugaya M, Takenoyama M, Osaki T et al. Establishment of 15 cancer cell lines from patients with lung cancer and the potential tools for immunotherapy. *Chest* 2002; 122: 282-288.
25. Passlick B, Izbicki JR, Simmel S et al. Expression of major histocompatibility class I and class II antigens and intercellular adhesion molecule-1 on operable non-small cell lung carcinomas: frequency and prognostic significance. *Eur J Cancer* 1994; 30A: 376-381.
26. Kikuchi E, Yamazaki K, Torigoe T et al. HLA class I antigen expression is associated with a favorable prognosis in early stage non-small cell lung cancer. *Cancer Science* 2007; 98: 1424-1430.
27. Traversari C, Meazza R, Coppolecchia M et al. IFN-gamma gene transfer restores HLA-class I expression and MAGE-3 antigen presentation to CTL in HLA-deficient small cell lung cancer. *Gene Therapy* 1997; 4: 1029-1035.
28. Restifo NP, Esquivel F, Kawakami Y et al. Identification of Human Cancers Deficient in Antigen Processing. *Journal of Experimental Medicine* 1993; 177: 265-272.

29. Batra RK, Lin A, Sharma S et al. Non-small cell lung cancer-derived soluble mediators enhance apoptosis in activated T lymphocytes through an I kappa B kinase-dependent mechanism. *Cancer Research* 2003; 63: 642-646.
30. Torre-Amione G, Beauchamp RD, Koeppen H et al. A highly immunogenic tumor transfected with a murine transforming growth factor type beta 1 cDNA escapes immune surveillance. *Proc Natl Acad Sci U S A* 1990; 87: 1486-1490.
31. Masui T, Wakefield LM, Lechner JF et al. Type-Beta Transforming Growth-Factor Is the Primary Differentiation-Inducing Serum Factor for Normal Human Bronchial Epithelial-Cells. *Proceedings of the National Academy of Sciences of the United States of America* 1986; 83: 2438-2442.
32. Liu VC, Wong LY, Jang T et al. Tumor evasion of the immune system by converting CD4(+) CD25(-) T cells into CD4(+) CD25(+) T regulatory cells: Role of tumor-derived TGF-beta. *Journal of Immunology* 2007; 178: 2883-2892.
33. Hasegawa Y, Takanashi S, Kanehira Y et al. Transforming growth factor-beta1 level correlates with angiogenesis, tumor progression, and prognosis in patients with nonsmall cell lung carcinoma. *Cancer* 2001; 91: 964-971.
34. Sharma S, Stolina M, Lin Y et al. T cell-derived IL-10 promotes lung cancer growth by suppressing both T cell and APC function. *Journal of Immunology* 1999; 163: 5020-5028.
35. Spagnoli GC, Juretic A, Schultzhater E et al. On the Relative Roles of Interleukin-2 and Interleukin-10 in the Generation of Lymphokine-Activated Killer-Cell Activity. *Cellular Immunology* 1993; 146: 391-405.
36. Huang M, Sharma S, Mao JT, Dubinett SM. Non-small cell lung cancer-derived soluble mediators and prostaglandin E2 enhance peripheral blood lymphocyte IL-10 transcription and protein production. *J Immunol* 1996; 157: 5512-5520.
37. Topalian SL, Drake CG, Pardoll DM. Targeting the PD-1/B7-H1(PD-L1) pathway to activate anti-tumor immunity. *Curr Opin Immunol* 2012; 24: 207-212.

38. Nishimura H, Nose M, Hiai H et al. Development of lupus-like autoimmune diseases by disruption of the PD-1 gene encoding an ITIM motif-carrying immunoreceptor. *Immunity* 1999; 11: 141-151.
39. Lynch TJ, Bondarenko I, Luft A et al. Ipilimumab in combination with paclitaxel and carboplatin as first-line treatment in stage IIIB/IV non-small-cell lung cancer: results from a randomized, double-blind, multicenter phase II study. *J Clin Oncol* 2012; 30: 2046-2054.
40. Woo EY, Yeh H, Chu CS et al. Cutting edge: Regulatory T cells from lung cancer patients directly inhibit autologous T cell proliferation. *Journal of Immunology* 2002; 168: 4272-4276.
41. Wolf AM, Wolf D, Steurer M et al. Increase of regulatory T cells in the peripheral blood of cancer patients. *Clinical Cancer Research* 2003; 9: 606-612.
42. Gabrilovich DI, Nagaraj S. Myeloid-derived suppressor cells as regulators of the immune system. *Nature Reviews Immunology* 2009; 9: 162-174.
43. Srivastava MK, Zhu L, Harris-White M et al. Myeloid Suppressor Cell Depletion Augments Antitumor Activity in Lung Cancer. *Plos One* 2012; 7.
44. Liu X, Shin N, Koblish HK et al. Selective inhibition of IDO1 effectively regulates mediators of antitumor immunity. *Blood* 2010; 115: 3520-3530.
45. Gan HK, You B, Pond GR, Chen EX. Assumptions of expected benefits in randomized phase III trials evaluating systemic treatments for cancer. *J Natl Cancer Inst* 2012; 104: 590-598.
46. Sistigu A, Viaud S, Chaput N et al. Immunomodulatory effects of cyclophosphamide and implementations for vaccine design. *Semin Immunopathol* 2011; 33: 369-383.
47. Finke JH, Rini B, Ireland J et al. Sunitinib reverses type-1 immune suppression and decreases T-regulatory cells in renal cell carcinoma patients. *Clin Cancer Res* 2008; 14: 6674-6682.


48. Locher C, Conforti R, Aymeric L et al. Desirable cell death during anticancer chemotherapy. *Ann N Y Acad Sci* 2010; 1209: 99-108.
49. Garrison K, Hahn T, Lee WC et al. The small molecule TGF-beta signaling inhibitor SM16 synergizes with agonistic OX40 antibody to suppress established mammary tumors and reduce spontaneous metastasis. *Cancer Immunol Immunother* 2012; 61: 511-521.
50. Sinha P, Clements VK, Fulton AM, Ostrand-Rosenberg S. Prostaglandin E2 promotes tumor progression by inducing myeloid-derived suppressor cells. *Cancer Res* 2007; 67: 4507-4513.
51. Brahmer JR HL, Gandhi G et al. . Nivolumab (anti-PD-1, BMS-936558, ONO-4538) in patients (pts) with advanced non-small-cell lung cancer (NSCLC): Survival and clinical activity by subgroup analysis. *J Clin Oncol* 32:5s, 2014 (suppl; abstr 8112[^]) 2014.
52. Binder DC, Engels B, Arina A, et al. Antigen-specific bacterial vaccine combined with anti-PD-L1 rescues dysfunctional endogenous T cells to reject long-established cancer. *Cancer Immunol Res* 2013; 1: 123-33.
53. Golden EB, Demaria S, Schiff PB, et al. An Abscopal Response to Radiation and Ipilimumab in a Patient with Metastatic Non-Small Cell Lung Cancer. *Cancer Immunology Research* 2013; 1: 365-372.
54. Kandalaf LE, Tanyi J, Chiang C, et al. Autologous whole-tumor antigen vaccination in combination with adoptive T cell therapy for patients with recurrent ovarian cancer. *Cancer Research* 2013; 73.
55. Tesniere A, Panaretakis T, Kepp O, et al. Molecular characteristics of immunogenic cancer cell death. *Cell Death Differ* 2008; 15: 3-12.
56. Tang C, Wang X, Soh H, et al. Combining radiation and immunotherapy: a new systemic therapy for solid tumors? *Cancer Immunol Res* 2014; 2: 831-8.



57. Nesslinger NJ, Ng A, Tsang KY, et al. A viral vaccine encoding prostate-specific antigen induces antigen spreading to a common set of self-proteins in prostate cancer patients. *Clin Cancer Res* 2010; 16: 4046-56.
58. Prado-Garcia H, Romero-Garcia S, Aguilar-Cazares D et al. Tumor-induced CD8+ T-cell dysfunction in lung cancer patients. *Clin Dev Immunol* 2012; 2012: 741741.
59. Sopori M. Effects of cigarette smoke on the immune system. *Nat Rev Immunol*. 2002; 2: 372-7.
60. Kalra R, Singh SP, Savage SM, et al. Effects of cigarette smoke on immune response: chronic exposure to cigarette smoke impairs antigen-mediated signaling in T cells and depletes IP3-sensitive Ca(2+) stores. *J Pharmacol Exp Ther* 2000; 293: 166-71.
61. Stämpfli MR, Anderson GP. How cigarette smoke skews immune responses to promote infection, lung disease and cancer. *Nat Rev Immunol*. 2009; 9: 377-84.
62. Herbst RS, Soria JC, Kowanetz M, et al. Predictive correlates of response to the anti-PD-L1 antibody MPDL3280A in cancer patients. *Nature* 2014; 515: 563-7.
63. Zhang J, Fujimoto J, Zhang J, et al. Intratumor heterogeneity in localized lung adenocarcinomas delineated by multiregion sequencing. *Science*. 2014; 346: 256-9.
64. Akbay EA, Koyama S, Carretero J et al. Activation of the PD-1 Pathway Contributes to Immune Escape in EGFR-Driven Lung Tumors. *Cancer Discovery* 2013; 3: 1355-1363.
65. Azuma K, Ota K, Kawahara A, et al. Association of PD-L1 overexpression with activating EGFR mutations in surgically resected nonsmall-cell lung cancer. *Ann Oncol* 2014; 25: 1935-40.
66. Xu C, Fillmore CM, Koyama S, et al. Loss of Lkb1 and Pten leads to lung squamous cell carcinoma with elevated PD-L1 expression. *Cancer Cell* 2014; 25: 590-604.

figure legends

Figure 1: Important considerations in clinical translation of successful active immunotherapy.

Figure 2: Mechanisms of humoral and cellular immune dysregulation in lung cancer. Tumor antigens are presented by antigen-presenting cells in the context of major histocompatibility complex class I or class II molecules are recognized by the T cell receptors (TCR). Additional co-stimulatory signals are mediated through constitutively expressed co-stimulatory molecules on the T cell and the APC (for example, B7-CD28) are also necessary for T cell activation. The presence of both signals trigger intracellular events resulting in the activation and interleukin (IL)-2-dependent clonal proliferation of T cells. Some of the mechanisms employed by tumors to escape the host immune response and promote immune tolerance are represented 1. Suppression of antigen presenting machinery, 2. Soluble factors released by the tumor (examples include interleukin 10, and transforming growth factor β), 3. Tumor infiltrating T lymphocytes, 4. Myeloid derived suppressor cells and 5. The immunosuppressive effects of tobacco smoke.





Immunotherapy



- Patient selection
- Stage of disease to intervene
- Integration with chemotherapy, radiation, other immunotherapies
- Overcoming humoral and cellular immune dysregulation
- Trial designs
- End-points
- Assessment of response
- Regulatory issues




Table 1. Phase II and phase III studies of selected antigen-specific immunotherapeutic approaches in non-small cell lung cancer

Investigational agent	Phase of study	n	Patients	Primary end-point	Primary end-point outcome		Significance of differences between treatment group and control group
					Treatment group	Control group	
Tecemotide	Randomized phase II (Butts, Maksymiuk et al. 2011)	171	IIIB or IV NSCLC SD or OR after first line chemotherapy or chemoradiation	OS	17.2 m	13 m	NS
	Randomized, double-blind placebo-controlled phase III (Butts, Socinski et al. 2014)	1513	IIIA (T3, N2 only), IIIB and IV SD or OR after first-line chemotherapy or chemoradiation	OS	25.6 m	22.3 m	NS
Belagenpumatucel-L	Randomized, dose-variable phase II (Nemunaitis, Dillman et al. 2006)	75	II, IIIA, IIIB, and IV; low tumor burden Completed conventional therapy	OS	Dose-related improvements in survival in three-treatment arms*	NA	No control arm
	Randomized, double-blind placebo-controlled phase III (G Giaccone, E Juhász et al. 2013)	532	IIIA (T3, N2 only), IIIB and IV SD or OR after primary platinum based chemoradiotherapy	OS	20.3	17.8	NS
Melanoma associated antigen-A3 vaccine	Randomized phase II (Vansteenkiste J 2007)	182	Completely resected IB/II MAGE-A3-expressing tumor	DFI	HR 0.74 (95% CI 0.44–1.20) p=0.107**	NA	NS
	Randomized, double-blind placebo-controlled phase III (release 2014)	2312	completely resected IB, II or IIIA MAGE-A3-expressing tumor	DFS	Not available	Not available	NS
*Three doses (1.25, 2.5, or 5.0 × 10 ⁷ cells/injection) of belagenpumatucel-L were studied in 3 cohorts of 25, 26 and 24 patients each							
** HR in favor of the MAGE-A3 group							