

CTRI Dataset and Description

CTRI Field	Description
Public title of study	<p>Title intended for the lay public in easily understood language.</p> <p>Example: A clinical trial to study the effects of two drugs, ramipril and candesartan in patients with high blood pressure and type 2 diabetes mellitus.</p>
Scientific title of study Acronym, if any	<p>Scientific title of the study as it appears in the protocol submitted for funding and ethical review. Include trial acronym if available.</p> <p>Example: A randomized double-blind placebo controlled crossover clinical trial to compare the safety and efficacy of ramipril and candesartan in hypertensive patients with type 2 diabetes mellitus.</p> <p>Acronym RACE</p>
Secondary IDs, if any	<p>Secondary ID is any number that is associated with a clinical trial, such as Protocol Number or any other Trial Registry Number, if registered in another Registry, such as ClinicalTrials.gov, ACTR, ISRCTN etc. There is no limit on the number of Secondary ID numbers that can be provided.</p> <p>In case of a multi-country trial, the trial may have already been registered in another registry such as the www.ClinicalTrials.gov. However, the trial, if also being conducted in India needs to be registered in the CTRI as well. In this case, the ClinicalTrials.gov identifying number would be this trial's Secondary ID number.</p> <p>Universal Trial Number UTN (earlier known as UTRN) may be obtained from http://apps.who.int/trialsearch/utn.aspx Please quote the obtained UTN number under SECONDARY ID. Currently obtaining the UTN is not mandatory</p> <p>If there are no secondary IDs select NIL from the drop down list and type in NIL in the corresponding box.</p>

Principal Investigator's name and address	Details should include name, official address, affiliation and designation, contact telephone and fax numbers and email ID. For a multi-center study, enter the contact information for the lead Principal Investigator (PI) or overall Trial Coordinator. Designated person must be from India (for trials being conducted in India). This is not a mandatory field.
Contact person (Scientific Query)	Details should include name, official address, affiliation and designation, email address, telephone number, Fax No and postal address, and affiliation of the local person (in case of multi-country trial) to contact for scientific queries about the trial (local principal investigator, medical contact of sponsor). May or may not be the same as the PI.
Contact person (Public Query)	Details should include name, official address, affiliation and designation, email address, telephone number, Fax No and postal address of the contact who will respond to general queries, including information about current recruitment status. This may or may not be the same as the contact person for scientific queries.
Source/s of monetary or material support	Major source/s of monetary or material or infrastructural support for the trial (e.g., funding agency, foundation, company, hospital, university, etc).
Primary sponsor	<p>Name and address of the individual, organization, group or other legal person taking responsibility for securing the arrangements to initiate and/or manage a study (including arrangements to ensure that the study design meets appropriate standards and to ensure appropriate conduct and reporting).</p> <p>The Primary Sponsor is responsible for ensuring that the trial is properly registered. The Primary Sponsor may or may not be the main source of funding.</p> <p>In commercial trials, the primary sponsor is normally the main applicant for regulatory authorization to begin the study. It may or may not be the main source of funding.</p> <p>In investigator initiated trials, the principal investigator is the primary sponsor, though the affiliated institution may be the main source of funding, and acknowledged under "Source/s of Monetary or Material Support".</p>

Secondary Sponsor	<p>Name and address of additional individuals, organizations or other legal persons, if any, that have agreed with the primary sponsor to take on responsibilities of sponsorship.</p> <p>A secondary sponsor may have –</p> <ul style="list-style-type: none"> Agreed to take on all the responsibilities of sponsorship jointly with the primary sponsor; To form a group with the primary sponsor in which the responsibilities of sponsorship are allocated among the members of the group; To act as the sponsor’s legal representative in relation to some or all of the trial sites; or to take responsibility for the accuracy of trial registration information submitted.
Countries of recruitment	<p>Select from drop down list, countries from which participants are intended to be, or have been recruited.</p> <p>E.g.: India - for trials conducted only in India; India, USA, France - for multi-country trials (as the case may be)</p>
Site/s of study	<p>List all site/s within India including the site address as well as the complete address, email, telephone number and Fax No of responsible contact person at each site (This individual should be a medically qualified person and to whom the EC approval is addressed, i.e. the PI; in case a separate person is mentioned, the PI should also be mentioned in any of the other contact person details (“PI or Overall trial coordinator, Contact Person (Scientific query or Public query)).</p> <p>For PMS trials with hundreds of trial sites, site details may be ‘copy-pasted’ in the Brief summary, specifying the few initiated sites under “Site/s of study”</p>
Name of Ethics Committee and approval status	<p>Provide name of Ethics Committee (EC) from whom approval has been sought; for multi-centre trials, add names of all ECs from whom approval has been sought; also provide approval status, i.e. submitted for approval or approved with date.</p> <p>As per the DCGI notice File No. ECR/Misc/Indt EC/007/2013 dated 30/07/2013, no clinical trial shall be approved by IEC(Independent Ethics Committee). Accordingly select ‘NO’ for IEC. Therefore henceforth all clinical trials should be approved by Institutional Ethics Committee.</p>

	Mention EC approval status of each site separately even if it is "under review" and/or from the same IEC (please mention the city from which the IEC functions).
Regulatory clearance obtained from DCGI	Mention whether approval has been taken from Drugs Controller General (India) [DCGI] or not. If DCGI has been notified, the same should be selected. It is the responsibility of the Sponsor to ascertain whether or not DCGI approval is required for a particular trial.
Health condition/problem studied	State the primary health condition(s) or problem(s) studied. If the study is conducted in healthy human volunteers belonging to the target population of the intervention (e.g., preventative or screening interventions), enter the particular health condition(s) or problem(s) being prevented or screened Example: Type 2 Diabetes Mellitus; Hypertension
Study type	Please indicate if trial part of post-graduation thesis Please select whether the trial is an Interventional trial, Observational trial or Post marketing surveillance Interventional Trial: An interventional trial is one that prospectively assigns human participants or groups of humans to one or more health-related intervention to evaluate the effect on outcomes. Choose the intervention that is best suited for the trial, more than one option may be selected according to the intervention/s being used; e.g. Drug & Ayurveda Observational Trial An observational trial is one where no experimental intervention or treatment is given to human participants. In this type of trial, the investigator only observes the effect of a risk factor, diagnostic test, or treatment on a particular outcome. Choose the intervention that is best suited for the trial. PMS: Post marketing surveillance study

	<p>Choose a Study Design from the list provided</p> <p>Examples:</p> <p>Single arm trial</p> <p>Non-randomized, placebo controlled trial</p> <p>Non-randomized, active controlled trial</p> <p>Non-randomized, multiple arm trial</p> <p>Randomized parallel group trial</p> <p>Randomized, parallel group, placebo controlled trial</p> <p>Randomized, parallel group, active controlled trial</p> <p>Randomized, parallel group, multiple arm trial</p> <p>Randomized, crossover trial</p> <p>Cluster randomized trial</p> <p>Randomized factorial trial</p>
Intervention and comparator agent	<p>Enter the specific name of the intervention/s and the comparator/control/s being studied. Use the International Non-Proprietary Name if possible (not brand/trade names). For an unregistered drug, the generic name, chemical name, or company serial number is acceptable. If the intervention consists of several separate treatments, list them all in one line separated by commas (e.g., "low-fat diet, exercise").</p> <p>The control intervention/s is/are the interventions against which the study intervention is evaluated (e.g., placebo, no treatment, active control). If an active control is used, be sure to enter in the name/s of that intervention, or enter "placebo" or "no treatment" as applicable.</p> <p>For each intervention, describe other intervention details as applicable (dose, duration, mode of administration, etc).</p> <p>Example:</p> <p>Ramipril</p> <p>2.5 mg OD for 12 months</p> <p>Candesartan</p> <p>16 mg OD for 12 months</p>

	For observational trials, NIL may be mentioned with trial details mentioned in the Brief Summary.
Inclusion/ Exclusion criteria	<p>Inclusion and exclusion criteria for participant selection, including age and sex. Age and sex to be mentioned in specific boxes.</p> <p>Example: Inclusion criteria</p> <p>Adult males or females with a diagnosis of type 2 diabetes mellitus and hypertension</p> <p>Hypertension defined as systolic blood pressure of 140 mmHg or diastolic blood pressure of 90 mmHg</p> <p>Diabetes defined as those patients with fasting glucose levels of ≥ 126 mg/dl or random blood glucose $>$ or $= 200$ mg/dl, HbA1c $>$ or $= 6.5\%$, 2 h blood glucose on 75 g oral glucose tolerance test (OGTT) $>$ or $= 200$ mg/dl, or current treatment with hypoglycemic therapy).</p> <p>Exclusion Criteria: A history of coronary heart disease or stroke, serum creatinine ≥ 1.5 mg/dl, albuminuria ≥ 40 μg/min, and use of lipid-lowering drugs, aspirin, or other antihypertensive agents.</p> <p>Please separate each criteria by using the "Enter" button</p>
Method of generating randomization sequence	<p>The method used to generate the random allocation sequence.</p> <p>The main purpose of randomization is to eliminate selection bias and balance known or unknown confounding factors in order to create a control group that is as similar as possible to the treatment group.</p> <p>Methods for randomly assigning participants to groups, which limits bias, include the use of a table of random numbers and a computer program that generates random numbers.</p> <p>Methods of assignment that are prone to bias include alternating assignment or assignment by date of birth or hospital admission number.</p>

	<p>Example:</p> <p>Coin toss, lottery, toss of dice, shuffling cards etc</p> <p>Random number table</p> <p>Computer generated randomization</p> <p>Permuted block randomization, fixed</p> <p>Permuted block randomization, variable</p> <p>Stratified randomization</p> <p>Stratified block randomization</p> <p>Adaptive randomization, such as minimization</p> <p>Other, describe</p>
Method of allocation concealment	<p>Concealment of the randomization sequence is critical to prevent selection bias. Adequate allocation concealment is a pre-requisite for adequate blinding.</p> <p>Adequate allocation concealment methods include:</p> <ul style="list-style-type: none"> centralized (e.g. allocation by a central office unaware of subject characteristics) pharmacy-controlled randomization pre-numbered or coded identical containers which are administered serially to participants on-site computer system combined with allocations kept in a locked unreadable computer file sequentially numbered, sealed, opaque envelopes <p>Allocation concealment that is prone to bias include</p> <ul style="list-style-type: none"> alternation case record numbers dates of birth or day of the week an open list of random numbers and any procedure that is entirely transparent before allocation
Blinding/masking	<p>Blinding refers to methods used to prevent participants and investigators from knowing what interventions are being used to reduce bias. Open trials do not use blinding. Masking refers to the methods used to camouflage interventions to achieve blinding.</p>

	<p>Examples:</p> <ul style="list-style-type: none"> Open label Participant blinded Investigator blinded Outcome assessor blinded Double blind double dummy Participant and Investigator blinded Participant and outcome assessor blinded Participant, investigator and outcome assessor blinded Participant, investigator, outcome assessor and data-entry operator/statistician blinded
<p>Primary outcome/s</p>	<p>Outcomes are events, variables, or experiences that are measured because it is believed that they may be influenced by the intervention. The primary outcome could be the outcome used in sample size calculations, or the main outcome/s used to determine the effects of the intervention/s.</p> <p>Enter the names of all primary outcomes in the trial as well as the pre-specified timepoint/s of primary interest. Be as specific as possible with the metric used (e.g., "% with Beck Depression Score > 10 "rather than just "depression").</p> <p>Examples</p> <p>Outcome Name: all-cause mortality, Time-points: 5 years; or</p> <p>Outcome Name: Mean Beck Depression Score, Time-point: 18 weeks.</p>
<p>Secondary outcome/s</p>	<p>Secondary outcomes are events, variables, or experiences that are of secondary interest or that are measured at time-points of secondary interest. A secondary outcome may involve the same event, variable, or experience as the primary outcome, but measured at time-points other than those of primary interest (e.g., Primary outcome: all-cause mortality at 5 years; Secondary outcome: all-cause mortality at 1 year, 3 years), or may involve a different event, variable, or experience altogether (e.g., Primary outcome: all-cause mortality at 5 years; Secondary outcome: hospitalization rate at 5 years).</p>

	Enter the name and time-point(s) for all secondary outcomes of clinical and/or scientific importance.
Target sample size	<p>Total number of participants that the trial plans to enroll. For global/multi-country trials, enter both Total sample size and Target sample size from India. This is a numbers only field.</p> <p>Example Target sample size 120 India 500 Total</p> <p>For trials being conducted only in India, target sample should be same under both columns</p> <p>Target sample size 120 India 120 Total</p>
Phase of trial	<p>Phases of investigation, usually applied to a drug trial</p> <p><u>Phase 1</u>: includes initial study to determine the metabolism and pharmacologic actions of drugs in humans, the side effects associated with increasing doses, and to gain early evidence of effectiveness; may include healthy participants and/or patients (such as those testing anticancer or anti-HIV drugs) . Trials are often dose ranging trials which are done to determine the maximum dose of a new medication that can be safely given to a patient.</p> <p><u>Phase 1 / Phase 2</u>: for trials that are at a combined stage of phases 1 and 2</p> <p><u>Phase 2</u>: includes controlled clinical study conducted to evaluate/test the effectiveness of a new drug/medication or intervention for a particular indication or indications in patients with the disease or condition being studied and to determine the common short-term side effects and risks</p> <p><u>Phase 2 / Phase 3</u>: for trials that are at a combined stage of phases 2 and 3</p> <p><u>Phase 3</u>: includes expanded controlled and uncontrolled trials after preliminary evidence suggesting effectiveness of the drug has been obtained, and are intended to gather additional information to evaluate the overall benefit-risk relationship of a new drug/medication or intervention, including possible</p>

	<p>adverse reactions. It is also to provide an adequate basis for physician labeling</p> <p><u>Phase 3 /Phase 4:</u> For trials that are at a combined stage of phases 3 and 4</p> <p><u>Phase 4:</u> Studies (other than routine surveillance) performed after drug is marketed and is related to the approved indication. Trials are done to monitor the toxicity, risks, utility, benefits and optimal use after the efficacy of the drug/medication or intervention has been proven.</p> <p><u>N/A (Not applicable):</u> This selection is for a non-drug trial, BA/BE trial.</p> <p><u>Post marketing surveillance:</u> Routine surveillance trials after marketing approval</p> <p>Example Phase 3</p>
Date of first enrollment	<p>Select anticipated or actual date of enrollment of the first participant from the calendar.</p> <p>For global/multi-country trials, both global trial start date as well as start date in India should be mentioned.</p> <p>Example</p> <p>date of first enrollment 02/05/2009 India</p> <p style="padding-left: 150px;">15/06/2010 Global</p>
Estimated duration of trial	<p>Specify the expected time duration of trial, starting from enrollment of first patient to final submission of report.</p>
Recruitment status of trial	<p>Indicate status of trial. For global/multi-country trials enter status of global arm as well as Indian arm</p> <ul style="list-style-type: none"> ○ Not Yet Recruiting: Yet to initiate patient enrolment ○ Open to Recruitment: Participants are currently being recruited and enrolled ○ Suspended: There is a temporary halt in recruitment and enrolment but potentially will resume ○ Completed: Closed to recruitment of participants and data analysis complete

	<p>Closed to recruitment of participants: Follow- up continuing</p> <p>Other(Terminated): Recruiting or enrolling participants has halted and will not resume</p> <p><i>Additional Note:</i> Once a study is completed or terminated, the registered trial status must be updated accordingly. Subsequently two additional data set fields will be displayed which are to be filled as appropriate.</p> <p>Date of Study Completion (India) Date of Study Completion (Global)</p> <p>The date of study completion is the date of last visit of last trial participant (even if trial was terminated). For a detailed definition visit WHO's ICTRP (https://prsinfo.clinicaltrials.gov/definitions.html#PrimaryCompletionDate).</p> <p>For trials which are being conducted only in India, Date of Study Completion (Global) would be Not Applicable.</p> <p>AND</p> <p>Final Enrolment numbers achieved (Total) Final Enrolment numbers achieved (India)</p> <p>For global i.e. multi-country trials, both global and India final and actual enrolled number of patients should be mentioned. For trials which are being conducted only in India, the same number should be mentioned in both the above sections.</p>
Brief Summary	<p>Short description of the primary purpose of the protocol, including a brief statement of the study hypothesis. Include publication/s details (link/reference), if any.</p> <p>Example: This study is a randomized, double blind, parallel group, multi-centre trial comparing the safety and efficacy of Ramipril 2.5 mg daily and Candesartan 16 mg daily for 12 months in 500 patients with diabetes and hypertension that will be conducted in five centers in India, three in France and five in USA. The primary outcome measures will be all-cause mortality at five years and Mean Beck Depression Score at 18 weeks. The</p>

	secondary outcomes will be all-cause mortality at 6 months and 1 year; and Mean glycosylated hemoglobin A1C at 4 and 8 weeks.
--	---

For any clarifications please contact –

Clinical Trials Registry – India
National Institute of Medical Statistics
Indian Council of Medical Research
Ansari Nagar
New Delhi-110029
India
Tel: 011-26589635; 011-26588803
Email: ctri@gov.in